

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 December 2002 (19.12.2002)

PCT

(10) International Publication Number
WO 02/100403 A1

(51) International Patent Classification⁷: **A61K 31/42**,
C07D 263/32, A61P 3/10, C07D 231/12, A61K 31/422,
31/426, 31/427, C07D 233/58, 277/24, 277/24, 277/38,
C07C 271/24, 271/22, 233/87, 233/63, C07D 413/12,
417/12, 413/14, 413/04, 413/10, 417/14, 417/04, 213/81,
307/83

Fishers, IN 46038 (US). **SMITH, Daryl, Lynn** [US/US];
7656 Madden Lane, Fishers, IN 46038 (US). **VANCE,**
Jennifer, Ann [US/US]; 132 El Bosque, San Diego, CA
95134 (US). **WARSHAWSKY, Alan, M.** [US/US]; 13828
Spring Mill Ponds Circle, Carmel, IN 46032 (US).

(21) International Application Number: PCT/US02/15143

(74) Agents: **JANG, Soonhee** et al.; Eli Lilly and Company,
Lilly Corporate Center, Indianapolis, IN 46285 (US).

(22) International Filing Date: 24 May 2002 (24.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/296,701 7 June 2001 (07.06.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT (util-
ity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ (utility model), CZ, DE, DK
(utility model), DK, DM, DZ, EC, EE (utility model), EE,
ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,
NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility
model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **ELI
LILLY AND COMPANY** [US/US]; Lilly Corporate
Center, Indianapolis, IN 46285 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GOSSETT, Lynn,**
Stacy [US/US]; 7345 Donegal Drive, Indianapolis, IN
46217 (US). **GREEN, Jonathan, Edward** [US/US];
1019 East 96th Street, Apartment D, Indianapolis, IN
46240 (US). **HENRY, James, Robert** [US/US]; 7307
Elm Ridge Drive, Indianapolis, IN 46236 (US). **JONES,**
Winton, Dennis, Junior [US/US]; 1227 East 126th Street,
Carmel, IN 46033 (US). **MATTHEWS, Donald, Paul**
[US/US]; 7736 Wawasee Drive, Indianapolis, IN 46250
(US). **SHEN, Quan, Rong** [CN/US]; 10565 Harlow Drive,

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: MODULATORS OF PEROXISOME PROLIFERATOR ACTIVATED RECEPTORS (PPAR)

(57) Abstract: The present invention is directed to a compound of formula I, and pharmaceutically acceptable salts, solvates, hy-
drates or stereoisomer thereof, which are useful in treating Syndrome X, Type II diabetes, hyperglycemia, hyperlipidemia, obesity,
coagulopathy, hypertension, arteriosclerosis, and other disorders related to Syndrome X as well as cardiovascular diseases.

WO 02/100403 A1

MODULATORS OF PEROXISOME PROLIFERATOR ACTIVATED RECEPTORS (PPAR)

5

FIELD OF THE INVENTION

The present invention relates to a compound of peroxisome proliferator activated receptor (PPAR) agonists, which are useful for the treatment and/or prevention of disorders modulated by a PPAR.

10

BACKGROUND OF THE INVENTION

The peroxisome proliferator activated receptors (PPARs) are members of the nuclear receptor gene family that are activated by fatty acids and fatty acid metabolites. The PPARs belong to the subset of nuclear receptors that function as heterodimers with the 9-*cis* retinoic acid receptor (RXR). Three subtypes, designated PPAR α , PPAR γ and PPAR δ , are found in species ranging from *Xenopus* to humans.

15

PPAR α is the main subtype in the liver and has facilitated analysis of the mechanism by which peroxisome proliferators exert their pleiotropic effects. PPAR α is activated by a number of medium and long-chain fatty acids, and it is involved in stimulating β -oxidation of fatty acids. PPAR α is also involved with the activity of fibrates and fatty acids in rodents and humans. Fibrates such as clofibrate, fenofibrate, bezafibrate, ciprofibrate, beclofibrate and etofibrate, as well as gemfibrozil, produce a substantial reduction in plasma triglycerides along with moderate reduction in low-density lipoprotein (LDL) cholesterol, and they are used particularly for the treatment of hypertriglyceridemia.

20

25

PPAR γ is the main subtype in adipose tissue and involved in activating the program of adipocyte differentiation. PPAR γ is not involved in stimulating peroxisome proliferation in the liver. There are two isomers of PPAR γ : PPAR γ 1 and PPAR γ 2, which differ only in that PPAR γ 2 contains an additional 28 amino acids present at the amino terminus. The DNA sequences for the PPAR γ receptors are described in Elbrecht, et al., BBRC 224;431-437 (1996). Although peroxisome proliferators, including the fibrates and fatty acids, activate the transcriptional activity of PPAR's, only prostaglandin J₂ derivatives have been identified as natural ligands for PPAR γ , which also binds the anti-diabetic agents thiazolidinediones with high affinity. The physiological functions of

30

35

5 PPAR α and PPAR γ in lipid and carbohydrate metabolism were uncovered once it was recognized that they were the receptors for the fibrate and glitazone drugs, respectively.

PPAR α and PPAR γ receptors have been implicated in diabetes mellitus, cardiovascular disease, obesity, and gastrointestinal disease, such as inflammatory bowel disease and other inflammation related illnesses. Such inflammation related illnesses
10 include, but are not limited to Alzheimer's disease, Crohn's disease, rheumatoid arthritis, psoriasis, and ischemia reprofusion injury.

By contrast, PPAR δ (also referred to as PPAR β and NUC1) is not reported to be receptor for any known class of drug molecules, and its role in mammalian physiology has remained undefined. The human nuclear receptor gene PPAR δ (hPPAR δ) has been
15 cloned from a human osteosarcoma cell cDNA library and is fully described in A. Schmidt et al., *Molecular Endocrinology*, 6:1634-1641 (1992).

Diabetes is a disease in which a mammal's ability to regulate glucose levels in the blood is impaired because the mammal has a reduced ability to convert glucose to glycogen for storage in muscle and liver cells. In Type I diabetes, this reduced
20 ability to store glucose is caused by reduced insulin production. "Type II Diabetes" or "non-insulin dependent diabetes mellitus" (NIDDM) is the form of diabetes, which is due to a profound resistance to insulin stimulating or regulatory effect on glucose and lipid metabolism in the main insulin-sensitive tissues, muscle, liver and adipose tissue. This resistance to insulin responsiveness results in insufficient insulin activation of glucose
25 uptake, oxidation and storage in muscle and inadequate insulin repression of lipolysis in adipose tissue and of glucose production and secretion in liver. When these cells become desensitized to insulin, the body tries to compensate by producing abnormally high levels of insulin and hyperinsulemia results. Hyperinsulemia is associated with hypertension and elevated body weight. Since insulin is involved in promoting the cellular uptake of
30 glucose, amino acids and triglycerides from the blood by insulin sensitive cells, insulin insensitivity can result in elevated levels of triglycerides and LDL (known as the "bad" cholesterol) which are risk factors in cardiovascular diseases. The constellation of symptoms which includes hyperinsulemia combined with hypertension, elevated body weight, elevated triglycerides and elevated LDL is known as Syndrome X.

35 Hyperlipidemia is a condition which is characterized by an abnormal increase in serum lipids, such as cholesterol, triglycerides and phospholipids. These lipids

5 do not circulate freely in solution in plasma, but are bound to proteins and transported as macromolecular complexes called lipoproteins. One form of hyperlipidemia is hypercholesterolemia, characterized by the existence of elevated LDL cholesterol levels. The initial treatment for hypercholesterolemia is often a diet low in fat and cholesterol coupled with appropriate physical exercise. Drug intervention is initiated if LDL-
10 lowering goals are not met by diet and exercise alone. It is desirable to lower elevated levels of LDL cholesterol and increase levels of HDL cholesterol. Generally, it has been found that increased levels of HDL are associated with lower risk for coronary heart disease (CHD). See Gordon, et al., *Am. J. Med.*, 62, 707-714 (1977); Stampfer, et al., *N. England J. Med.*, 325, 373- 381 (1991); and Kannel, et al., *Ann. Internal Med.*, 90, 85-91
15 (1979). An example of an HDL raising agent is nicotinic acid, but the quantities needed to achieve HDL elevation are associated with undesirable effects, such as flushing.

There are several treatments currently available for treating diabetes mellitus but these treatments still remain unsatisfactory and have limitations. While physical exercise and reduction in dietary intake of calories will improve the diabetic
20 condition, compliance with this approach can be poor because of sedentary lifestyles and excess food consumption, in particular high fat-containing food. Therefore, treatment with hypoglycemics, such as sulfonylureas (e.g., chlorpropamide, tolbutamide, tolazamide and acetohexamide) and biguanides (e.g. phenformin and metformin) are often necessary as the disease progresses. Sulfonylureas stimulate the β cells of the
25 pancreas to secrete more insulin as the disease progresses. However, the response of the β cells eventually fails and treatment with insulin injections is necessary. In addition, both sulfonylurea treatment and insulin injection have the life threatening side effect of hypoglycemic coma, and thus patients using these treatments must carefully control dosage.

30 It has been well established that improved glycemic control in patients with diabetes (Type I and Type II) is accompanied by decreased microvascular complications (DCCT and UKPDS). Due to difficulty in maintaining adequate glycemic control over time in patients with Type II diabetes, the use of insulin sensitizers in the therapy of Type II diabetes is growing. There is also a growing body of evidence that
35 PPAR γ agonist, insulin sensitizer, may have benefits in the treatment of Type II diabetes beyond their effects in improving glycemic control.


-4-

5 In the last decade a class of compounds known as thiazolidinediones (e.g. U.S. Pat. Nos. 5,089,514; 4,342,771; 4,367,234; 4,340,605; and 5,306,726) have emerged as effective anidiabetic agents that have been shown to increase the sensitivity of insulin sensitive tissues, such as skeletal muscle, liver and adipose, to insulin. Increasing insulin sensitivity rather than the amount of insulin in the blood reduces the likelihood of hypoglycemic coma. Although thiazolidinediones have been shown to increase insulin sensitivity by binding to PPAR γ receptors, this treatment also produces unwanted side effects such as weight gain and, for troglitazone, liver toxicity.

10 In view of the above, there exists a need for new pharmaceutical agents which modulate these receptors to prevent, treat and/or alleviate these diseases or conditions while ameliorating side effects of current treatments.

SUMMARY OF THE INVENTION

The present invention relates to compound of novel peroxisome proliferator activated receptor agonists having a structural formula 1,



20 and pharmaceutically acceptable salts, solvates, hydrates or stereoisomers thereof, wherein:


n^1 is 2, 3, 4 or 5;


V is a bond or O;

25 X is CH₂ or O;

p is 0 or 1;

m is 1-4;

Y¹ is:  wherein,

30  is: aryl or heteroaryl,

5 wherein aryl and heteroaryl are optionally substituted with one or more groups
independently selected from the group consisting of:

hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, halo, haloalkyl and haloalkyloxy;

Y^{1a} is: hydrogen,

10 (C₀-C₃)alkyl-aryl,

C(O)-aryl,

heteroaryl,

cycloalkyl,

heterocycloalkyl,

15 aryloxy,

NR⁵(CH₂)_mOR⁵,

aryl-Z-aryl,

aryl-Z-heteroaryl,

aryl-Z-cycloalkyl,

20 aryl-Z-heterocycloalkyl,

heteroaryl-Z-aryl,

heteroaryl-Z-heterocycloalkyl or

heterocycloalkyl-Z-aryl,

25 wherein aryl, cycloalkyl, aryloxy, heteroaryl, and heterocycloalkyl are optionally
substituted with one or more substituents independently selected from the group
consisting of:

halo,

hydroxyl,

nitro,

30 cyano,

C₁-C₆ alkyl,

C₁-C₆ alkoxy optionally substituted with N(R⁵)₂,

haloalkyl,

N(R⁵)₂,

35 N[C(O)R⁵]₂,

-6-

5 $N[S(O)_2R^5]_2$,
 $NR^5S(O)_2R^5$,
 $NR^5C(O)R^5$,
 $NR^5C(O)OR^5$,
 $C(O)N(R^5)_2$,
 10 $C(O)OR^5$ and
 $C(O)R^5$;

 Z is: a bond,
 -oxygen-
 15 $-C(O)NR^5-$
 $-NR^5C(O)-$,
 $-NR^5C(O)O-$,
 $-C(O)-$,
 $-NR^5-$,
 20 $-[O]_p(CH_2)_m-$,
 $-(CH_2)_m[O]_p-$,
 $-NR^5(CH_2)_m-$ or
 $-(CH_2)_mNR^5-$;

25 Y^2 and Y^3 are each independently:

 hydrogen,
 C_1 - C_6 alkyl or
 C_1 - C_6 alkoxy;

30 Y^4 is: $(C_1-C_3)alkyl-NR^5C(O)-(C_0-C_5)alkyl-Y^7$,
 $(C_1-C_3)alkyl-NR^5C(O)-(C_2-C_5)alkenyl-Y^7$,
 $(C_1-C_3)alkyl-NR^5C(O)-(C_2-C_5)alkynyl-Y^7$;
 $(C_1-C_3)alkyl-NR^5C(O)O-(C_0-C_5)alkyl-Y^7$,
 $(C_1-C_3)alkyl-NR^5C(O)NR^5-(C_0-C_5)alkyl-Y^7$,
 35 $(C_1-C_3)alkyl-NR^5C(S)NR^5-(C_0-C_5)alkyl-Y^7$,
 $(C_0-C_3)alkyl-C(O)NR^5-(C_0-C_5)alkyl-Y^7$,

-7-

5 (C₁-C₃)alkyl-OC(O)NY¹⁰Y¹¹,
(C₁-C₃)alkyl-NY¹⁰Y¹¹,
(C₁-C₃)alkyl-O-(C₀-C₅)alkyl-Y⁷,
(C₁-C₃)alkyl-S-(C₀-C₅)alkyl-Y⁷ or
10 CN;

Y⁷ is: hydrogen,

aryl,

heteroaryl,

C₁-C₁₂ alkyl,

15 C₁-C₆ alkoxy,

cycloalkyl,

heterocycloalkyl,

aryloxy,

C(O)-heteroaryl or

20 SR⁶,

wherein alkyl, aryl, aryloxy, alkoxy, heteroaryl, cycloalkyl, and heterocycloalkyl
are optionally substituted with one or more groups independently selected from
R⁷:

25 Y¹⁰ and Y¹¹ are each independently:

hydrogen,

aryl,

heteroaryl,

C₁-C₁₀ alkyl,

30 cycloalkyl,

SO₂(R⁶); or

Y¹⁰ and Y¹¹ together are a 5- to 10-membered heterocycloalkyl ring or
heterocycloalkyl ring fused with aryl, and the heterocycloalkyl ring optionally
containing one or more heteroatoms selected from N, O or S; and wherein,

35

5 aryl, heteroaryl, heterocycloalkyl and alkyl are optionally substituted with one or more substituents independently selected from R^7 ;

R^5 is: hydrogen or C_1 - C_6 alkyl;

10 R^6 is: hydrogen,
 C_1 - C_{10} alkyl,
cycloalkyl,
aryl, or
heteroaryl,

15 wherein alkyl, cycloalkyl, aryl and heteroaryl are optionally substituted with one or more substituents independently selected from R^7 ;

R^7 is: halo,
nitro,
20 oxo,
cyano,
hydroxyl,
benzyl,
phenyl,
25 phenoxy,
heteroaryl,
 $C(O)R^6$,
 C_1 - C_{10} alkyl,
 C_1 - C_6 alkoxy,
30 C_1 - C_6 haloalkyl,
 C_1 - C_6 haloalkyloxy,
 $O(CH_2)_m$ -phenyl,
 $(CH_2)_mOC(O)$ -aryl,
 $C(O)OR^5$,
35 $S(O)_2R^5$,
 $S(O)_2N(R^5)_2$,

5 SR⁵ or
 N(R⁵)₂,

 wherein phenyl and phenoxy are optionally substituted with one or more groups independently selected from halo or trifluoromethyl.

 The compounds of the present invention are useful in the treatment or
10 prevention of diseases or condition relates to hyperglycemia, dyslipidemia, Type II diabetes, Type I diabetes, hypertriglyceridemia, syndrome X, insulin resistance, heart failure, diabetic dyslipidemia, hyperlipidemia, hypercholesteremia, hypertension, obesity, anorexia bulimia, anorexia nervosa, cardiovascular disease and other diseases where insulin resistance is a component.

15 In one embodiment, the present invention also relates to pharmaceutical compositions which comprising at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate, hydrate thereof and a pharmaceutically acceptable carrier. Within the scope of this invention also include a pharmaceutical composition containing additional therapeutic agent as well as at least one compound of
20 the present invention, or a pharmaceutically acceptable salt, solvate, hydrate thereof and a pharmaceutically acceptable carrier.


 In another embodiment, the present invention relates to a method of modulating a peroxisome proliferator activated receptor by contacting the receptor with at least one compound of Formula I, and pharmaceutically acceptable salts, solvates and
25 hydrates thereof.

-10-

5 DETAILED DESCRIPTION OF THE INVENTION

The compounds of the present invention are directed to peroxisome proliferator activated receptor (PPAR) agonists, which are useful for the treatment and/or prevention of disorders modulated by a PPAR.

10 An embodiment of the present invention is a compound of structural formula I:



and pharmaceutically acceptable salts, solvates, hydrates or stereoisomers thereof, wherein:

- 15 n^1 is 2, 3, 4 or 5;
 V is a bond or O;
 X is CH_2 or O;
 p is 0 or 1;
 m is 1-4;

20 Y^1 is: Y^{1a} —(Ar)— where, wherein,



is: aryl or heteroaryl,

wherein aryl and heteroaryl are optionally substituted with one or more groups independently selected from the group consisting of:

25 hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halo, haloalkyl and haloalkyloxy;

Y^{1a} is: hydrogen,

(C_0 - C_3)alkyl-aryl,

$\text{C}(\text{O})$ -aryl,

30 heteroaryl,

-11-

5 cycloalkyl,
heterocycloalkyl,
aryloxy,
NR⁵(CH₂)_mOR⁵,
aryl-Z-aryl,
10 aryl-Z-heteroaryl,
aryl-Z-cycloalkyl,
aryl-Z-heterocycloalkyl,
heteroaryl-Z-aryl,
heteroaryl-Z-heterocycloalkyl or
15 heterocycloalkyl-Z-aryl,
wherein aryl, cycloalkyl, aryloxy, heteroaryl, and heterocycloalkyl are optionally
substituted with one or more substituents independently selected from the group
consisting of:

20 halo,
hydroxyl,
nitro,
cyano,
C₁-C₆ alkyl,
C₁-C₆ alkoxy optionally substituted with N(R⁵)₂,
25 haloalkyl,
N(R⁵)₂,
N[C(O)R⁵]₂,
N[S(O)₂R⁵]₂,
NR⁵S(O)₂R⁵,
30 NR⁵C(O)R⁵,
NR⁵C(O)OR⁵,
C(O)N(R⁵)₂,
C(O)OR⁵ and
C(O)R⁵;

35

-12-

5 Z is: a bond,
 -oxygen-
 -C(O)NR⁵-
 -NR⁵C(O)-,
 -NR⁵C(O)O-,
 10 -C(O)-,
 -NR⁵-,
 -[O]_p(CH₂)_m-,
 -(CH₂)_m[O]_p-,
 -NR⁵(CH₂)_m- or
 15 - (CH₂)_mNR⁵-;

Y² and Y³ are each independently:

hydrogen,
 C₁-C₆ alkyl or
 20 C₁-C₆ alkoxy;

Y⁴ is: (C₁-C₃)alkyl-NR⁵C(O)-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-NR⁵C(O)-(C₂-C₅)alkenyl-Y⁷,
 (C₁-C₃)alkyl-NR⁵C(O)-(C₂-C₅)alkynyl-Y⁷;
 25 (C₁-C₃)alkyl-NR⁵C(O)O-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-NR⁵C(O)NR⁵-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-NR⁵C(S)NR⁵-(C₀-C₅)alkyl-Y⁷,
 (C₀-C₃)alkyl-C(O)NR⁵-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-OC(O)NY¹⁰Y¹¹,
 30 (C₁-C₃)alkyl-NY¹⁰Y¹¹,
 (C₁-C₃)alkyl-O-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-S-(C₀-C₅)alkyl-Y⁷ or
 CN;

35

-13-

5 Y⁷ is: hydrogen,
 aryl,
 heteroaryl,
 C₁-C₁₂ alkyl,
 C₁-C₆ alkoxy,
10 cycloalkyl,
 heterocycloalkyl,
 aryloxy,
 C(O)-heteroaryl or
 SR⁶,
15 wherein alkyl, aryl, aryloxy, alkoxy, heteroaryl, cycloalkyl, and heterocycloalkyl
 are optionally substituted with one or more groups independently selected from
 R⁷:

20 Y¹⁰ and Y¹¹ are each independently:
 hydrogen,
 aryl,
 heteroaryl,
 C₁-C₁₀ alkyl,
 cycloalkyl,
25 SO₂(R⁶); or
 Y¹⁰ and Y¹¹ together are a 5- to 10-membered heterocycloalkyl ring or
 heterocycloalkyl ring fused with aryl, and the heterocycloalkyl ring optionally
 containing one or more heteroatoms selected from N, O or S; and wherein,
 aryl, heteroaryl, heterocycloalkyl and alkyl are optionally substituted with one or
30 more substituents independently selected from R⁷;

R⁵ is: hydrogen or C₁-C₆ alkyl;

-14-


5 R^6 is: hydrogen,
 C_1 - C_{10} alkyl,
 cycloalkyl,
 aryl, or
 heteroaryl,

10 wherein alkyl, cycloalkyl, aryl and heteroaryl are optionally substituted with one
 or more substituents independently selected from R^7 ;

R^7 is: halo,
 nitro,
15 oxo,
 cyano,
 hydroxyl,
 benzyl,
 phenyl,
20 phenoxy,
 heteroaryl,
 $C(O)R^6$,
 C_1 - C_{10} alkyl,
 C_1 - C_6 alkoxy,
25 C_1 - C_6 haloalkyl,
 C_1 - C_6 haloalkyloxy,
 $O(CH_2)_m$ -phenyl,
 $(CH_2)_mOC(O)$ -aryl,
 $C(O)OR^5$,
30 $S(O)_2R^5$,
 $S(O)_2N(R^5)_2$,
 SR^5 or
 $N(R^5)_2$,

 wherein phenyl and phenoxy are optionally substituted with one or more groups
35 independently selected from halo or trifluoromethyl.

- 5 An embodiment of the present invention also include a compound represented by the structural Formula Ia:



Ia

or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

10

Y^1 is an unsubstituted or substituted group selected from the group consisting of: aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aryl- C_1 - C_4 alkyl, heteroaryl- C_1 - C_4 alkyl, cycloalkyl- C_1 - C_4 alkyl, and t-butyl;

Y^2 is selected from the group consisting of:

- 15 H, C_1 - C_{10} alkyl, cycloalkyl, (C_1 - C_{10} alkyl)- Y^5 , O- Y^6 ;

Y^5 is selected from the group consisting of:

an aryl, substituted aryl group, $-COR^4$, $-COOR^4$, $-CONR^6R^7$, $-CSR^4$, and $-C(S)NR^6R^7$;

Y^6 is selected from the group consisting of:

- an aliphatic group, a substituted aliphatic group, an aryl, substituted aryl group,
20 $-COR^4$, $-COOR^4$, $-CONR^6R^7$, $-CSR^4$, and $-C(S)NR^6R^7$;

R^4 , R^6 and R^7 are each independently selected from the group consisting of:

H, an aliphatic group, a substituted aliphatic group, an aryl group and a substituted aryl group;

- 25 Y^3 is selected from the group consisting of:

H, aliphatic, substituted aliphatic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and (C_1 - C_{10} alkyl)- R^8 ; R^8 is selected from the group consisting of aryl, substituted aryl, heteroaryl, and substituted heteroaryl; and

R^5 is selected from the group consisting of:

- 30 H, aliphatic group, a substituted aliphatic group, heteroaryl, substituted heteroaryl, an aryl, a substituted aryl, and (C_1 - C_{10} alkyl)- R^9 ;

5 R^9 is selected from the group consisting of:

aryl, substituted aryl, heteroaryl, and substituted heteroaryl, aminoalkyl, and cycloalkyl;

V is a bond or O;

X is CH_2 or O;

Y^4 is selected from the group consisting of:

10 $-(C_1-C_3)alkyl-O-W-Y^7$, $-C(O)NY^8Y^9$, $-(C_1-C_3)alkyl-NY^{10}Y^{11}$, and

$-(C_1-C_3)alkylN(Y^{13})W-(C_0-C_5)alkyl-Y^{14}$;

W is selected from the group consisting of:

a bond, $-CONY^{12}$, $-C(O)-$, $-OCH_2-$, C_1-C_6 alkyl, $-CO_2-$, $-CHOY^{15}$, $-CSNY^{16}$ and

15 $-SO_2-$;

Y^7 is selected from the group consisting of:

aryl, substituted aryl, heteroaryl, substituted heteroaryl, aliphatic, branched aliphatic and substituted (C_1-C_{10}) alkyl;

Y^8 , Y^9 , Y^{10} , Y^{11} , Y^{12} , Y^{13} , Y^{14} , Y^{15} , and Y^{16} are each independently selected from

20 the group consisting of:


aryl, substituted aryl, heteroaryl, substituted heteroaryl, aliphatic, branched aliphatic and substituted (C_1-C_{10}) alkyl; and n^1 are 2, 3, 4 or 5.

The substituent Y^1 of the compound represented by a formula Ia includes


25 $Y^{1a}-\text{Ar}$ where Y^{1a} and Ar are defined above in formula I, and substituted group includes the substituents recited above and as defined in the embodiment.

5

The compound of present invention as recited above, wherein Y^{1a} is selected from the group consisting of: aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aryloxy,




10 A preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein E is O or S.

- 5 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein q is 1 or 2; and each R^5 is independently hydrogen or methyl.

- 10 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein q is 1 or 2; and each R^5 is independently hydrogen or methyl.

- 5 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein q is 1 or 2; and each R^5 is independently hydrogen or methyl.

- 10 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein q is 0 or 1; and each R^5 is independently hydrogen or methyl.

Another preferred embodiment of the present invention is a compound represented by the following structural formula,




- 15 wherein q is 1 or 2; and each R^5 is independently hydrogen or methyl.

- 5 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein q is 1 or 2; and each R^5 is independently hydrogen or methyl.

- 10 Another preferred embodiment of the present invention is a compound represented by the following structural formula,



wherein q is 1 or 2; and each R^5 is independently hydrogen or methyl.

Another preferred embodiment of the present invention is a compound represented by the following structural formula,




15


wherein q is 1 or 2; and each R^5 , Y^2 and Y^3 are independently hydrogen or methyl.

-21-

- 5 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein Y^{1a} is optionally substituted phenyl, naphthyl,



- 10 and Z is a bond, oxygen, -NH-, -N(CH₃)-, -NHC(O)- or -C(O)NH-.

Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein Y^{1a} is optionally substituted phenyl, naphthyl or  and

- 15 Z is a bond, oxygen, -NH-, -N(CH₃)-, -NHC(O)- or -C(O)NH-.


-22-

- 5 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein Y^{1a} is optionally substituted aryl, heteroaryl, heterocycloalkyl, heteroaryl-Z-heterocycloalkyl or heteroaryl-Z-aryl.

- 10 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein q is 1 or 2; and each R⁵ is independently hydrogen or methyl.

- 15 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




- 5 wherein q is 1 or 2; and each R^5 is independently hydrogen or methyl.

Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein, Y^{1a} is hydrogen, aryl, heteroaryl, or aryloxy; q is 1 or 2; and n^1 is 2, 3, or 4.

- 10 Another preferred embodiment of the present invention is a compound represented by the following structural formula,




wherein, Y^{1a} is hydrogen, aryl, heteroaryl or aryloxy; q is 1 or 2; and n^1 is 2, 3, or 4.

- 5 A preferred compound is a compound represented by the following structural formula,









A preferred compound is a compound represented by the following structural formula,








10






More preferred compounds of the present invention are listed below:







No.	Compound	name
1		3-{2-(Diphenylacetylaminomethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
2		3-{2-[(2-Cyclopropylacetylaminomethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid







No.	Compound	name
3		3-{2-[(3-Methoxybenzoylamino)methyl]-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
4		3-{2-[(Biphenyl-2-carbonyl)amino]methyl}-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
5		3-(4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-[(2,5-dichlorothiophene-3-carbonyl)amino]methyl}phenyl} propionic acid
6		3-{2-(Isopropoxycarbonylaminomethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
7		3-{2-(1,3-Dioxo-1,3-dihydroisoindol-2-ylmethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
8		3-{4-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-[(3-phenylureido)methyl]phenyl} propionic acid







No.	Compound	name
9		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(isopropoxycarbonylaminoethyl)phenyl] propionic acid
10		3-{2-(Isopropoxycarbonylaminoethyl)-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid
11		3-{2-(Benzoylaminoethyl)-4-[3-(biphenyl-4-yloxy)propoxy]phenyl} propionic acid
12		3-(4-[3-(Biphenyl-4-yloxy)propoxy]-2-[[pyridine-2-carbonyl]amino]methyl)phenyl) propionic acid
13		3-{2-Benzylcarbamoyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
14		3-{2-Benzylcarbamoyl-4-[2-(2-biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]phenyl} propionic acid
15		3-{4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-cyanophenyl} propionic acid







No.	Compound	name
16		3-[4-[2-(2-Biphenyl-3-yl-5-methyloxazol-4-yl)ethoxy]-2-(1,3-dioxo-1,3-dihydroisindol-2-ylmethyl)phenyl] propionic acid
17		3-{2-(2-Isopropoxycarbonylaminoethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
18		3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
19		3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[5-methyl-2-(3-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
20		3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[5-methyl-2-(6-phenylpyridin-3-yl)thiazol-4-yl]ethoxy}phenyl) propionic acid








No.	Compound	name
21		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(2-isopropoxycarbonylaminoethyl)phenyl] propionic acid
22		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(isobutoxycarbonylaminoethyl)phenyl] propionic acid
23		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(cyclopentyloxycarbonylaminoethyl)phenyl] propionic acid
24		3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[2-(4-isopropoxyphenyl)-5-methyloxazol-4-yl]ethoxy}phenyl) propionic acid
25		3-(2-Benzylcarbamoyl-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid








No.	Compound	name
26		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-morpholin-4-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
27		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-piperidin-1-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
28		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-pyrimidin-2-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
29		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-pyrazin-2-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
30		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(6-phenoxy pyridin-3-yl)thiazol-4-yl]ethoxy}phenyl) propionic acid
31		3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid

No.	Compound	name
32		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-phenylaminophenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
33	 <p>ClH</p>	3-(4-{2-[5-Methyl-2-(6-phenylpyridin-3-yl)thiazol-4-yl]ethoxy}-2-[(pyridine-2-carbonyl)amino]methyl}phenyl) propionic acid HCl salt
34		3-{4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-[(3-methylbutyrylamino)methyl]phenyl} propionic acid
35		3-{2-(Isopropoxycarbonylaminomethyl)-4-[2-(5-methoxy-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
36		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(6-phenoxy pyridin-3-yl)oxazol-4-yl]ethoxy}phenyl) propionic acid
37		3-{4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-cyclohexylcarbamoxyloxy-methylphenyl} propionic acid

No.	Compound	name
38		3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid
39		3-(2-Cyclohexylcarbamoyloxymethyl-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
40		3-(2-Cyclohexylcarbamoyloxymethyl-4-{2-[5-methyl-2-(4-morpholin-4-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
41		3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[3-(tetrahydropyran-4-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid
42		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(cyclopropylmethoxycarbonylaminomethyl)phenyl] propionic acid
43		3-{2-(Cyclopropylmethoxycarbonylaminomethyl)-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid HCl salt

No.	Compound	name
44		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(cyclobutoxycarbonylaminomethyl)phenyl]propionic acid HCl salt
45		3-{2-(Cyclobutoxycarbonylaminomethyl)-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid HCl salt
46		3-[4-{2-[2-(4-Butyrylamino)phenyl]-5-methyloxazol-4-yl}ethoxy]-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid
47		3-{2-(Isopropoxycarbonylaminomethyl)-4-[2-(5-methyl-2-{4-[(pyridine-2-carbonyl)amino]phenyl}-oxazol-4-yl)-ethoxy]phenyl}-propionic acid
48		3-(4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-[(pyrazine-2-carbonyl)amino]methyl)phenyl) propionic acid
49		3-[4-{2-[2-(3-Cyclohexylcarbamoyl)phenyl]-5-methyloxazol-4-yl}ethoxy]-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid

No.	Compound	name
50		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(2-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
51		3-(2-Cyano-4-{2-[5-methyl-2-(4-phenoxy-phenyl)-oxazol-4-yl]-ethoxy}-phenyl) propionic acid
52		3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(pyridin-2-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid
53		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(4-trifluoromethylphenoxy)methyl] phenyl] propionic acid
54		3-{2-(Isobutoxycarbonylaminomethyl)-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid
55		3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(pyrimidin-2-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid
56		3-[4-[2-(2-Biphenyl-4-yl-5-methoxyoxazol-4-yl)ethoxy]-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid

No.	Compound	name
57		3-(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(pyridine-2-carbonyl)-amino]-methyl}-phenyl)-propionic acid
58		3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(2,4,5-trifluorobenzoylamino)-methyl]-phenyl}-propionic acid
59		3-{2-[(2,4-Difluorobenzoylamino)-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
60		3-(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(thiophene-2-carbonyl)-amino]-methyl}-phenyl)-propionic acid
61		3-(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(thiophene-2-carbonyl)-amino]-methyl}-phenyl)-propionic acid
62		3-{2-(Butyrylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
63		3-{2-[(Cyclobutanecarbonyl)-amino]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid

No.	Compound	name
64		3-{2-(Benzyloxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
65		3-{2-(tert-Butoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
66		3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(2-phenoxy-acetyl-amino)-methyl]-phenyl}-propionic acid
67		3-{2-[(Cyclopentanecarbonyl-amino)-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid

5

Also encompassed by the present invention is a pharmaceutical composition, comprising a pharmaceutically acceptable carrier and at least one compound of the present invention or a pharmaceutically acceptable salt, solvate or hydrate thereof.

Also encompassed by the present invention is a pharmaceutical composition comprising: (1) a compound of formula I according to Claim 1 or a pharmaceutically acceptable salt, solvate, hydrate or stereoisomer thereof; (2) a second therapeutic agent selected from the group consisting of insulin sensitizers, sulfonylureas, biguanides, thiazolidinediones, α -glucosidase inhibitors, insulin secretagogues, insulin, antihyperlipidemic agents, plasma HDL-raising agents, HMG-CoA reductase inhibitors, statins, acyl CoA:cholesterol acyltransferase inhibitors, antiobesity compounds, antihypercholesterolemic agents, fibrates, vitamins and aspirin; and (3) a pharmaceutically acceptable carrier.

5 Also encompassed by the present invention is a method of modulating a peroxisome proliferator activated receptor (PPAR), comprising the step of contacting the receptor with at least one compound of the present invention or a pharmaceutically acceptable salt, solvate or hydrate thereof. The peroxisome proliferator activated receptor is an alpha-receptor or a gamma-receptor.

10 Also encompassed by the present invention is a method for treating or preventing a peroxisome proliferator activated receptor-gamma mediated disease or condition comprising the step of administering a compound of Formula I or Formula Ia.

 Also encompassed by the present invention is a method for lowering blood-glucose comprising the step of administering an effective amount of a compound of
15 formula I or formula Ia.

 Also encompassed by the present invention is a method of treating or preventing disease or condition selected from the group consisting of hyperglycemia, dyslipidemia, Type II diabetes, Type I diabetes, hypertriglyceridemia, syndrome X, insulin resistance, heart failure, diabetic dyslipidemia, hyperlipidemia,
20 hypercholesteremia, hypertension, obesity, anorexia bulimia, anorexia nervosa, cardiovascular disease and other diseases where insulin resistance is a component, comprising the step of administering an effective amount of a compound of formula I or formula Ia.

 Also encompassed by the present invention is a method of treating or
25 preventing diabetes mellitus in a mammal comprising the step of administering to a mammal a therapeutically effective amount of at least one compound of formula I or formula Ia.

 Also encompassed by the present invention is a method of treating or preventing cardiovascular disease in a mammal comprising the step of administering to a
30 mammal a therapeutically effective amount of at least one compound of formula I or formula Ia, or a pharmaceutically acceptable salt, solvate, hydrate or stereoisomer thereof.

-37-

5 Also encompassed by the present invention is a method of treating or preventing syndrome X in a mammal, comprising the step of administering to the mammal a therapeutically effective amount of at least one compound of formula I or formula Ia, or a pharmaceutically acceptable salt, solvate, hydrate or stereoisomer thereof.





10 Also encompassed by the present invention is a method of treating or preventing disease or condition selected from the group consisting of hyperglycemia, dyslipidemia, Type II diabetes, Type I diabetes, hypertriglyceridemia, syndrome X, insulin resistance, heart failure, diabetic dyslipidemia, hyperlipidemia, hypercholesteremia, hypertension, obesity, anorexia bulimia, anorexia nervosa, cardiovascular disease and other diseases where insulin resistance is a component, 15 comprising the step of administering an effective amount of a compound of formula I or Ia, and an effective amount of second therapeutic agent selected from the group consisting of insulin sensitizers, sulfonylureas, biguanides, thiazolidinediones, α -glucosidase inhibitors, insulin secretagogues, insulin, antihyperlipidemic agents, plasma HDL-raising agents, HMG-CoA reductase inhibitors, statins, acyl CoA:cholesterol 20 acyltransferase inhibitors, antiobesity compounds, antihypercholesterolemic agents, fibrates, vitamins and aspirin.

 Also encompassed by the present invention is use of a compound of formula I or formula Ia and pharmaceutically acceptable salt, solvate, hydrate or stereoisomer thereof, for the manufacture of a medicament for the treatment of a 25 condition modulated by a PPAR.


-38-

5 An embodiment of the present invention also includes the compounds having a structural formula II to IX as shown below. These compounds are within the scope of the compound represented by a structural Formula Ia. The substituent Y¹ of



these compound includes $Y^{1a} - \text{Ar} -$, where Y^{1a} and Ar are defined above in formula I, and substituted groups includes substituents recited above and as defined in the
 10 embodiment herein.




-39-



5



wherein Y¹⁸ is (C₁-C₆) straight or branched alkyl; and



10 wherein Y¹⁸ is (C₁-C₆) straight or branched alkyl.

5 The terms used to describe the instant invention have the following meanings.

 As used herein, the term "aliphatic" or "aliphatic group" is a non-aromatic, consisting solely of carbon and hydrogen and may optionally contain one or more units of saturation, e.g., double and/or triple bonds (also refer herein as "alkenyl" and "alkynyl").

10 An aliphatic or aliphatic group may be straight chained, branched (also refer herein as "alkyl") or cyclic (also refer herein as "cycloalkyl"). When straight chained or branched, an aliphatic group typically contains between about 1 and about 10 carbon atoms, more typically between about 1 and about 6 carbon atoms. When cyclic, an aliphatic typically contains between about 3 and about 10 carbon atoms, more typically between about 3 and

15 about 7 carbon atoms. Aliphatics are preferably C₁-C₁₀ straight chained or branched alkyl groups (i.e. completely saturated aliphatic groups), more preferably C₁-C₆ straight chained or branched alkyl groups. Examples include, but are not limited to methyl, ethyl, propyl, n-propyl, iso-propyl, n-butyl, sec-butyl, and tert-butyl. Additional examples include, but are not limited to, cyclopropyl, cyclopentyl, cyclohexyl, cyclopentyl,

20 cyclohexyl and the like.

 The term "alkyl," unless otherwise indicated, refers to those alkyl groups of a designated number of carbon atoms of either a straight or branched saturated configuration. Examples of "alkyl" include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl, pentyl, hexyl, isopentyl and

25 the like. Alkyl as defined above may be optionally substituted with a designated number of substituents as set forth in the embodiment recited above.

 The term "alkenyl" means hydrocarbon chain of a specified number of carbon atoms of either a straight or branched configuration and having at least one carbon-carbon double bond, which may occur at any point along the chain, such as

30 ethenyl, propenyl, butenyl, pentenyl, vinyl, alkyl, 2-butenyl and the like. Alkenyl as defined above may be optionally substituted with designated number of substituents as set forth in the embodiment recited above.

 The term "alkynyl" means hydrocarbon chain of a specified number of carbon atoms of either a straight or branched configuration and having at least one

35 carbon-carbon triple bond, which may occur at any point along the chain. Example of alkynyl is acetylene. Alkynyl as defined above may be optionally substituted with

5 designated number of substituents as set forth in the embodiment recited above.

The term "cycloalkyl" refers to a saturated or partially saturated carbocycle containing one or more rings of from 3 to 12 carbon atoms. Examples of cycloalkyl includes, but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, and the like. Cycloalkyl as defined above also includes a tricycle, such as
10 adamantyl. Cycloalkyl as defined above may be optionally substituted with a designated number of substituents as set forth in the embodiment recited above.

The term "halo" refers to fluoro, chloro, bromo and iodo.

The term "haloalkyl" is a C₁-C₆ alkyl group, which is substituted with one or more halo atoms selected from F, Br, Cl and I. An example of a haloalkyl group is
15 trifluoromethyl.

The term "alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, tert-butoxy, pentoxy, and the like. Alkoxy as defined above may be optionally substituted with a designated number of substituents as set forth in the embodiment
20 recited above.

The term "haloalkyloxy" represents a C₁-C₆ haloalkyl group attached through an oxygen bridge, such as OCF₃. The "haloalkyloxy" as defined above may be optionally substituted with a designated number of substituents as set forth in the embodiment recited above.

25 The term "aryl" includes carbocyclic aromatic ring systems (e.g. phenyl), fused polycyclic aromatic ring systems (e.g. naphthyl and anthracenyl) and aromatic ring systems fused to carbocyclic non-aromatic ring systems (e.g., 1,2,3,4-tetrahydronaphthyl). "Aryl" as defined above may be optionally substituted with a designated number of substituents as set forth in the embodiment recited above.

30 The term "heteroaryl" group, as used herein, is an aromatic ring system having at least one heteroatom such as nitrogen, sulfur or oxygen and includes monocyclic, bicyclic or tricyclic aromatic ring of 5- to 14-carbon atoms containing one or more heteroatoms selected from O, N, or S. The "heteroaryl" as defined above may be optionally substituted with a designated number of substituents as set forth in the
35 embodiment recited above. Examples of heteroaryl are, but are not limited to, furanyl, thienyl (also referred to herein as "thiophenyl") thiazolyl, imidazolyl, isoxazolyl, oxazolyl,

5 pyrazoyl, pyrrolyl, pyrazinyl, pyridyl, pyrimidyl, pyrimidinyl and purinyl, cinnolinyl, benzofuranyl, benzothienyl, benzotriazolyl, benzoxazolyl, quinoline, isoxazolyl, isoquinoline and the like.

The term "heterocycloalkyl" refers to a non-aromatic ring which contains one or more oxygen, nitrogen or sulfur and includes a monocyclic, bicyclic or tricyclic
10 non-aromatic ring of 5 to 14 carbon atoms containing one or more heteroatoms selected from O, N or S. The "heterocycloalkyl" as defined above may be optionally substituted with a designated number of substituents as set forth in the embodiment recited above. Examples of heterocycloalkyl include, but are not limited to, morpholine, piperidine, piperazine, pyrrolidine, and thiomorpholine. The preferred heterocycloalkyl group is
15 morpholine.

An aryl-C₁-C₄-alkyl group, as used herein, is an aryl substituent that is linked to a compound by an alkyl group having from one to four carbon atoms.

A heteroaryl- C₁-C₄-alkyl group, as used herein, is a heteroaryl substituent that is linked to a compound by an alkyl group having from one to four carbon atoms.

20 A cycloalkyl- C₁-C₄-alkyl group, as used herein, is a cycloalkyl substituent that is linked to a compound by an alkyl group having from one to four carbon atoms.

An aminoalkyl group is an alkyl group having from one to six carbon atoms, which is substituted with at least one amine represented by NR¹²R¹² where each R¹² is independently a C₁-C₆ alkyl or both R¹² taken together with the nitrogen to which
25 they are attached form a five or six membered heterocycloalkyl.

Unless otherwise indicated, substituents for alkyl, alkenyl, cycloalkyl, aryl, heteroaryl and heterocycloalkyl also include halo, carboxyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, nitro, cyano, CHO, hydroxyl, C₁-C₆ alkanolic acid and -C(O)NR¹³R¹³ where each R¹³ is independently hydrogen or a C₁-C₆ alkyl.

30 Substituents for thiophen-2,5-diyl and phenylene include H, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl and C₁-C₄ haloalkoxy.

The term "active ingredient" means the compounds generically described by formula I as well as the salts, solvates and prodrugs of such compounds.

The term "pharmaceutically acceptable" means that the carrier, diluents,
35 excipients and salt must be compatible with the other ingredients of the composition, and not deleterious to the recipient thereof. Pharmaceutical compositions of the present

5 invention are prepared by procedures known in the art using well-known and readily available ingredients.

"Preventing" refers to reducing the likelihood that the recipient will incur or develop any of the pathological conditions described herein.

10 "Treating" refers to mediating a disease or condition, and preventing or mitigating its further progression or ameliorate the symptoms associated with the disease or condition.

"Pharmaceutically-effective amount" means that amount of a compound of the present invention, or of its salt, solvate, hydrate or prodrug thereof that will elicit the biological or medical response of a tissue, system or mammal. Such an amount can be
15 administered prophylactically to a patient thought to be susceptible to development of a disease or condition. Such amount when administered prophylactically to a patient can also be effective to prevent or lessen the severity of the mediated condition. Such an amount is intended to include an amount, which is sufficient to modulate a PPAR

20 Conditions mediated by PPAR α or PPAR γ receptors include, for example, diabetes mellitus, cardiovascular disease, Syndrome X, obesity and gastrointestinal disease. Additional conditions associated with the modulation of a PPAR receptor include inflammation related conditions which include, for example, IBD (inflammatory bowel disease), rheumatoid arthritis, psoriasis, Alzheimer's disease, Chrohn's disease and
25 ischemia reprofusion injury (stroke and miocardial infarction).

A "mammal" is an individual animal that is a member of the taxonomic class Mammalia. The class Mammalia includes humans, monkeys, chimpanzees, gorillas, cattle, swine, horses, sheep, dogs, cats, mice, rats and the like.

Administration to a human is most preferred. A human to whom the
30 compounds and compositions of the present invention are administered has a disease or condition in which control blood glucose levels are not adequately controlled without medical intervention, but wherein there is endogenous insulin present in the human's blood. Non-insulin dependent diabetes mellitus (NIDDM) is a chronic disease or condition characterized by the presence of insulin in the blood, even at levels above
35 normal, but resistance or lack of sensitivity to insulin action at the tissues.

5 Those skilled in the art will recognize that stereocenters exist in compound of Formula I and Formula Ia. Accordingly, the present invention includes all possible stereoisomers and geometric isomers of formula I including racemic compounds and the optically active isomers.

10 The compounds of Formula I and Formula Ia contain one or more chiral centers and exist in different optically active forms. When compounds of formula I contain one chiral center, the compounds exist in two enantiomeric forms and the present invention includes both enantiomers and mixtures of enantiomers, such as racemic mixtures. Resolution of the final product, an intermediate or a starting material may be effected by any suitable method known in the art, for example by formation of
15 diastereoisomeric salts which may be separated by crystallization; formation of diastereoisomeric derivatives or complexes which may be separated by crystallization and gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent such as enzymatic esterification; and gas-liquid or liquid chromatography in a chiral environment such as on a chiral support, for example silica
20 with a bound chiral ligand or in the presence of a chiral solvent. See also *Stereochemistry of Carbon Compounds* by E.L. Eliel (Mcgraw Hill, 1962) and *Tables of Resolving Agents* by S. H. Wilen. It will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step is required to liberate the desired enantiomeric form. Alternatively, specific
25 enantiomers may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer into the other by asymmetric transformation. In a more preferred embodiment, the compounds of the present invention are S-enantiomers.

30 When a compound of Formula I or Formula Ia has more than one chiral substituents, it may exist in diastereoisomeric forms. The diastereoisomeric pairs may be separated by methods known to those skilled in the art, for example chromatography or crystallization and the individual enantiomers within each pair may be separated as described above. The present invention includes each diastereoisomer of compounds of formula I and mixtures thereof.

35

5 Certain compounds of formula I or Ia may exist in different stable conformational forms which may be separable. Torsional asymmetry due to restricted rotation about an asymmetric single bond, for example because of steric hindrance or ring strain, may permit separation of different conformers. The present invention includes each conformational isomer of compounds of formula I and Ia and mixtures thereof.

10 Certain compound of formula I or Ia may exist in zwitterionic form, and the present invention includes each zwitterionic form of compounds of formula I or Ia and mixtures thereof.

 Certain compounds of formula I or Ia and their salts may exist in more
15 than one crystal form. Polymorphs of compounds of formula I form part of the present invention and may be prepared by crystallization of a compound of formula I or Ia under different conditions, such as using different solvents or different solvent mixtures for recrystallization; crystallization at different temperatures; and various modes of cooling ranging from very fast to very slow cooling during crystallization. Polymorphs may also
20 be obtained by heating or melting a compound of Formula I or Formula Ia followed by gradual or fast cooling. The presence of polymorphs may be determined by solid probe NMR spectroscopy, IR spectroscopy, differential scanning calorimetry, powder X-ray diffraction or other available techniques.

 Certain compounds of formula I or Ia and their salts may exist in more
25 than one crystal form, and the present invention includes each crystal form and mixtures thereof.

 Certain compounds of formula I or Ia and their salts may also exist in the form of solvates, for example hydrates, and the present invention includes each solvate and mixtures thereof.

30 "Pharmaceutically-acceptable salt" refers to salts of the compounds of formula I or Ia which are substantially non-toxic to mammals. Typical pharmaceutically acceptable salts include those salts prepared by reaction of the compounds of the present invention with a mineral, organic acid: an organic base or inorganic base. Such salts are known as base addition salts, respectively. It should be recognized that the particular
35 counterion forming a part of any salt of the present invention is not of a critical nature so

5 long as the salt as a whole is pharmaceutically acceptable and the counterion does not contribute undesired qualities to the salt as a whole.

By virtue of its acidic moiety, a compound of formula I or Ia forms salts with pharmaceutically acceptable bases. Some examples of base addition salts include metal salts such as aluminum; alkali metal salts such as lithium, sodium or potassium; and
10 alkaline earth metal salts such as calcium, magnesium, ammonium, or substituted ammonium salts. Examples of substituted ammonium salts include, for instance, those with lower alkylamines such as trimethylamine and triethylamine; hydroxyalkylamines such as 2-hydroxyethylamine, bis-(2-hydroxyethyl)-amine or tri-(2-hydroxyethyl)-amine; cycloalkylamines such as bicyclohexylamine or dibenzylpiperidine, N-benzyl- β -
15 phenethylamine, dehydroabietylamine, N,N'-bisdehydro-abietylamine, glucamine, N-piperazine methylglucamine; bases of the pyridine type such as pyridine, collidine, quinine or quinoline; and salts of basic amino acids such as lysine and arginine.

Examples of inorganic bases include, without limitation, sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium
20 bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like.

Compounds of formula I or Ia, which are substituted with a basic group, may exist as salts with pharmaceutically acceptable acids. The present invention includes such salts. Examples of such salts include hydrochlorides, hydrobromides, sulfates, methanesulfonates, nitrates, maleates, acetates, citrates, fumarates, tartrates [e.g. (+)-
25 tartrates, (-)-tartrates or mixtures thereof including racemic mixtures], succinates, benzoates and salts with amino acids such as glutamic acid. These salts may be prepared by methods known to those skilled in the art.

Certain compounds of formula I or Ia and their salts may also exist in the form of solvates, for example hydrates, and the present invention includes each solvate
30 and mixtures thereof.

The compounds of present invention, which bind to and activate the PPARs, lower one or more of glucose, insulin, triglycerides, fatty acids and/or cholesterol, and are therefore useful for the treatment and/or prevention of hyperglycemia, dyslipidemia and in particular Type II diabetes as well as other diseases
35 including syndrome X, Type I diabetes, hypertriglyceridemia, insulin resistance, diabetic dyslipidemia, hyperlipidemia, hypercholesteremia, heart failure, coagulopathy,

5 hypertension, and cardiovascular diseases, especially arteriosclerosis. In addition, these compounds are indicated to be useful for the regulation of appetite and food intake in subjects suffering from disorders such as obesity, anorexia bulimia and anorexia nervosa.

The compounds and compositions of the present invention are also useful to treat acute or transient disorders in insulin sensitivity, which sometimes occurs
10 following a surgery, trauma, myocardial infarction and the like. The compounds and compositions of the present invention are also useful for lowering serum triglyceride levels. Elevated triglyceride level, whether caused by genetic predisposition or by a high fat diet, is a risk factor for the development of heart disease, stroke, and circulatory system disorders and diseases. The physician of ordinary skill will know how to identify
15 humans who can benefit from administration of the compounds and compositions of the present invention.

The present invention further provides a method for the treatment and/or prophylaxis of hyperglycemia in a human or non-human mammal which comprises administering an effective, non-toxic amount of a compound of formula I, or a tautomeric
20 form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof to a hyperglycemic human or non-human mammal in need thereof.

The compounds of the present invention are useful as therapeutic substances in preventing or treating Syndrome X, diabetes mellitus and related endocrine
25 and cardiovascular disorders and diseases in human or non-human animals.

The present invention also relates to the use of a compound of formula I as described above for the manufacture of a medicament for treating a PPAR α or PPAR γ mediated condition, separately or in combination.

A therapeutically effective amount of a compound of formula I can be
30 used for the preparation of a medicament useful for treating Syndrome X, diabetes, treating obesity, lowering tryglyceride levels, raising the plasma level of high density lipoprotein, and for treating, preventing or reducing the risk of developing arteriosclerosis, and for preventing or reducing the risk of having a first or subsequent atherosclerotic disease event in mammals, particularly in humans. In general, a
35 therapeutically effective amount of a compound of formula I of the present invention typically reduces serum glucose levels, more specifically HbA1c, of a patient by about

- 5 0.7% or more; typically reduces serum triglyceride levels of a patient by about 20% or more; and increases serum HDL levels in a patient. Preferably, HDL levels can be increased by about 30% or more.

Additionally, an effective amount of a compound of formula I and a therapeutically effective amount of one or more active agents selected from
10 antihyperlipidemic agent, plasma HDL-raising agents, antihypercholesterolemic agents, fibrates, vitamins, aspirin, insulin secretagogues, insulin and the like can be used together for the preparation of a medicament useful for the above described treatments.

Advantageously, compositions containing the compound of formula I or the salts thereof may be provided in dosage unit form, preferably each dosage unit
15 containing from about 1 to about 500 mg. It is understood that the amount of the compounds or compounds of formula I that will be administered is determined by a physician considering of all the relevant circumstances.

Syndrome X includes pre-diabetic insulin resistance syndrome and the resulting complications thereof, insulin resistance, non-insulin dependent diabetes,
20 dyslipidemia, hyperglycemia obesity, coagulopathy, hypertension and other complications associated with diabetes. The methods and treatments mentioned herein include the above and encompass the treatment and/or prophylaxis of any one of or any combination of the following: pre-diabetic insulin resistance syndrome, the resulting complications thereof, insulin resistance, Type II or non-insulin dependent diabetes,
25 dyslipidemia, hyperglycemia, obesity and the complications associated with diabetes including cardiovascular disease, especially arteriosclerosis.

The compositions are formulated and administered in the same general manner as detailed herein. The compounds of the present invention may be used effectively alone or in combination with one or more additional active agents depending
30 on the desired target therapy. Combination therapy includes administration of a single pharmaceutical dosage composition, which contains a compound of formula I and one or more additional active agents, as well as administration of a compound of formula I and each active agent in its own separate pharmaceutical dosage formulation. For example, a compound of formula I or thereof and an insulin secretagogue such as biguanides,
35 thiazolidinediones, sulfonylureas, insulin or α -glucosidase inhibitors can be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or

5 each agent administered in separate oral dosage formulations. Where separate dosage formulations are used, a compound of formula I and one or more additional active agents can be administered at essentially the same time, i.e., concurrently or at separately staggered times, i.e., sequentially; combination therapy is understood to include all these regimens.

10 An example of combination treatment or prevention of arteriosclerosis may involve administration of a compound of formula I or salts thereof in combination with one or more of second active therapeutic agents: antihyperlipidemic agents; plasma HDL-raising agents; antihypercholesterolemic agents, fibrates, vitamins, aspirin and the like. As noted above, the compounds of formula I can be administered in combination
15 with more than one additional active agent.

Another example of combination therapy can be seen in treating diabetes and related disorders wherein the compounds of formula I or salts thereof can be effectively used in combination with second active therapeutic, such as sulfonylureas, biguanides, thiazolidinediones, α -glucosidase inhibitors, other insulin secretagogues,
20 insulin as well as the active agents discussed above for treating arteriosclerosis.

The examples of second therapeutic agents are insulin sensitizers, PPAR γ agonists, glitazones, troglitazone, pioglitazone, englitazone, MCC-555, BRL 49653, biguanides, metformin, phenformin, insulin, insulin minetics, sulfonylureas, tolbutamide, glipizide, α -glucosidase inhibitors, acarbose, cholesterol lowering agent,
25 HMG-CoA reductase inhibitors, lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin, rivastatin, other statins, sequestrates, cholestyramine, colestipol, dialkylaminoalkyl derivatives of a cross-linked dextran, nicotiny alcohol, nicotinic acid: a nicotinic acid salt, PPAR α agonists, fenofibric acid derivatives, gemfibrozil, clofibrate, fenofibrate, benzaifibrate, inhibitors of cholesterol absorption, beta-sitosterol, acryl
30 CoA:cholesterol acyltransferase inhibitors, melinamide, probucol, PPAR δ agonists, antiobesity compounds, fenfluramine, dexfenfluramine, phentiramine, sulbitramine, orlistat, neuropeptide Y5 inhibitors, β_3 adrenergic receptor agonists, and ileal bile acid transporter inhibitors.

-50-

5 The compounds of the present invention and the pharmaceutically acceptable salts, solvates and hydrates thereof have valuable pharmacological properties and can be used in pharmaceutical compositions containing a therapeutically effective amount of a compound of the present invention, or pharmaceutically acceptable salts, esters or prodrugs thereof, in combination with one or more pharmaceutically acceptable
10 excipients. Excipients are inert substances such as, without limitation carriers, diluents, fillers, flavoring agents, sweeteners, lubricants, solubilizers, suspending agents, wetting agents, binders, disintegrating agents, encapsulating material and other conventional adjuvants. Proper formulation is dependent upon the route of administration chosen. Pharmaceutical compositions typically contain from about 1 to about 99 weight percent of
15 the active ingredient, which is a compound of the present invention.

 Preferably, the pharmaceutical formulation is in unit dosage form. A "unit dosage form" is a physically discrete unit containing a unit dose suitable for administration in human subjects or other mammals. For example, a unit dosage form can be a capsule or tablet, or a number of capsules or tablets. A "unit dose" is a
20 predetermined quantity of the active compound of the present invention, calculated to produce the desired therapeutic effect, in association with one or more pharmaceutically acceptable excipients. The quantity of active ingredient in a unit dose may be varied or adjusted from about 0.1 to about 1000 milligrams or more according to the particular treatment involved.

25 The dosage regimen utilizing the compounds of the present invention is selected by one of ordinary skill in the medical or veterinary arts considering various factors, such as without limitation, the species, age, weight, sex, medical condition of the recipient, the severity of the condition to be treated, the route of administration, the level of metabolic and excretory function of the recipient, the dosage form employed, the
30 particular compound and salt thereof employed, and the like.

 Preferably, the compounds of the present invention are administered in a single daily dose, or the total daily dose may be administered in divided doses of two, three or more times per day. Where delivery is via transdermal forms, administration is continuous.

35

5 Suitable routes of administration of pharmaceutical compositions of the present invention include, for example, oral, eye drop, rectal, transmucosal, topical or intestinal administration; parenteral delivery (bolus or infusion), including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intravenous, intravenous, intraperitoneal, intranasal, or intraocular injections. The compounds of the present invention can also be administered in a targeted drug delivery system, such as in a liposome coated with endothelial cell-specific antibody.

 For oral administration, the compounds of the present invention can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the present invention to be formulated as tablets, pills, powders, sachets, granules, dragees, capsules, liquids, elixirs, tinctures, gels, emulsions, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by combining the active compound with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores.

 For oral administration in the form of a tablet or capsule, the active ingredient may be combined with an oral, non-toxic, pharmaceutically-acceptable carrier, such as, without limitation, lactose, starch, sucrose, glucose, methyl cellulose, calcium carbonate, calcium phosphate, calcium sulfate, sodium carbonate, mannitol, sorbitol, and the like; together with, optionally, disintegrating agents, such as, without limitation, cross-linked polyvinyl pyrrolidone, maize, starch, methyl cellulose, agar, bentonite, xanthan gum, alginic acid: or a salt thereof such as sodium alginate, and the like; and, optionally, binding agents, for example, without limitation, gelatin, acacia, natural sugars, beta-lactose, corn sweeteners, natural and synthetic gums, acacia, tragacanth, sodium alginate, carboxymethyl-cellulose, polyethylene glycol, waxes, and the like; and, optionally, lubricating agents, for example, without limitation, magnesium stearate, sodium stearate, stearic acid: sodium oleate, sodium benzoate, sodium acetate, sodium chloride, talc, and the like. When a dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as a fatty oil.

5 Solid form formulations include powders, tablets and capsules. A solid carrier can be one or more substances, which may also act as flavoring agents, lubricants, solubilisers, suspending agents, binders, tablet disintegrating agents and encapsulating material.

10 In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

15 Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both. A syrup or elixir may contain, in addition to the active ingredient, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor.

20 Sterile liquid formulations include suspensions, emulsions, syrups, and elixirs. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable carrier, such as sterile water, sterile organic solvent, or a mixture of both sterile water and sterile organic solvent.

25 The active ingredient can also be dissolved in a suitable organic solvent, for example, aqueous propylene glycol. Other compositions can be made by dispersing the finely divided active ingredient in aqueous starch or sodium carboxymethyl cellulose solution or in a suitable oil.

30 Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

35 Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active

5 compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added.

All formulations for oral administration should be in dosages suitable for such administration. Particularly suitable compositions for oral administration are unit dosage forms such as tablets and capsules.

10 For parental administration the compounds of the present invention or salts thereof can be combined with sterile aqueous or organic media to form injectable solutions or suspensions. Formulations for injection may be presented in unit dosage form, such as in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or
15 aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that each syringability exists. It must be stable
20 under the conditions of manufacture and storage and must be preserved against any contamination. The carrier can be solvent or dispersion medium containing, for example, water, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiological saline buffer, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), propylene glycol and liquid polyethylene glycol),
25 suitable mixtures thereof, and vegetable oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The injectable solutions prepared in this manner can then be administered intravenously, intraperitoneally, subcutaneously, or intramuscularly, with intramuscular administration being preferred in humans.

30 For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. The active compounds can also be administered intranasally as, for example, liquid drops or spray.

For buccal administration, the compositions may take the form of tablets
35 or lozenges formulated in a conventional manner.

5 For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of a dry powder inhaler, or an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of pressurized
10 aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

 Pharmaceutical compositions of the present invention can be manufactured
15 in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

 In making the compositions of the present invention, the active ingredient will usually be admixed with a carrier, or diluted by a carrier, or enclosed
20 within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, lyophilized solid or paste, semi-solid, or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), or ointment, containing for example up to
25 10% by weight of the active compound. The compounds of the present invention are preferably formulated prior to administration.

 The following pharmaceutical formulations 1 through 8 are illustrative only and are not intended to limit the scope of the invention in any way. "Active Ingredient" refers to a compound of the present invention according to formula I and/or Ia
30 or salts thereof.

5 Formulation 1

Hard gelatin capsules are prepared using the following ingredients:

	Quantity (mg/capsule)
Active Ingredient	250
Starch, dried	200
Magnesium stearate	10
Total	460 mg

Formulation 2

A tablet is prepared using the ingredients below:

	Quantity (mg/tablet)
Active Ingredient	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
Stearic acid	5
Total	665 mg

- 10 The components are blended and compressed to form tablets each weighing 665 mg.

Formulation 3

An aerosol solution is prepared containing the following components:

	Weight
Active Ingredient	0.25
Ethanol	25.75
Propellant 22 (chlorodifluoromethane)	74.00
Total	100.00

- 15 The Active Ingredient is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to 30°C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the propellant. The valve units are then fitted to the container.

-56-

5 Formulation 4

Tablets, each containing 60 mg of Active ingredient, are made as follows:

Active Ingredient	60 mg
Starch	45 mg
Microcrystalline cellulose	35 mg
Polyvinylpyrrolidone (as 10% solution in water)	4 mg
Sodium carboxymethyl starch	4.5 mg
Magnesium stearate	0.5 mg
Talc	1 mg
Total	150 mg

10 The Active Ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The aqueous solution containing polyvinylpyrrolidone is mixed with the resultant powder, and the mixture then is passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

15

Formulation 5

Capsules, each containing 80 mg of Active Ingredient, are made as follows:

Active Ingredient	80 mg
Starch	59 mg
Microcrystalline cellulose	59 mg
Magnesium stearate	2 mg
Total	200 mg

20 The Active Ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

-57-

5 Formulation 6

Suppositories, each containing 225 mg of Active Ingredient, are made as follows:

Active Ingredient	225 mg
Saturated fatty acid glycerides	2,000 mg
Total	2,225 mg

10 The Active Ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2g capacities and allowed to cool.

Formulation 7

15 Suspensions each containing 50 mg of Active Ingredient per 5 ml dose are made as follows:

Active Ingredient	50 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 ml
Benzoic acid solution	0.10 ml
Flavor	q.v.
Color	q.v.
Purified water to total	5 ml

20 The Active Ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with a portion of the water and added, with stirring. Sufficient water is then added to produce the required volume.

5 Formulation 8

An intravenous formulation may be prepared as follows:

Active Ingredient	100 mg
Isotonic saline	1,000 ml

The solution of the above materials generally is administered intravenously to a subject at a rate of 1 ml per minute.

- 10 In yet another embodiment of the present invention, the compound is radiolabelled, such as with carbon-14 or tritiated. Said radiolabelled or tritiated compounds are useful as reference standards for in vitro assays to identify new PPAR α and PPAR γ agonists.

15 Binding and Cotransfection Studies

- The in vitro potency of compounds in modulating PPAR γ , PPAR α and PPAR δ receptors are determined by the procedures detailed below. DNA-dependent binding (ABCD binding) is carried out using Scintillation Proximity Assay (SPA) technology with PPAR receptors. Tritium-labeled PPAR α and PPAR γ agonists are used as radioligands for generating displacement curves and IC₅₀ values with compounds of the present invention. Cotransfection assays are carried out in CV-1 cells. The reporter plasmid contains an acylCoA oxidase (AOX) PPRE and TK promoter upstream of the luciferase reporter cDNA. Appropriate PPARs and RXR α are constitutively expressed using plasmids containing the CMV promoter. Since for PPAR α and PPAR β , interference by endogenous PPAR γ in CV-1 cells is an issue, in order to eliminate such interference, a GAL4 chimeric system is used in which the DNA binding domain of the transfected PPAR is replaced by that of GAL4, and the GAL4 response element is utilized in place of the AOX PPRE. Cotransfection efficacy is determined relative to PPAR α agonist and PPAR γ agonist reference molecules. Efficacies are determined by computer fit to a concentration-response curve, or in some cases at a single high concentration of agonist (10 μ M). A typical range for concentration determination (IC₅₀) is from 1nM to 10 μ M. For binding or cotransfection studies with receptors other than
- 20
- 25
- 30

5 PPARs, similar assays are carried out using appropriate ligands, receptors, reporter constructs and etc. for that particular receptor.

These studies are carried out to evaluate the ability of compounds of the present invention to bind to and/or activate various nuclear transcription factors, particularly huPPAR α ("hu" indicates "human"), huPPAR γ and huPPAR δ . These studies
10 provide in-vitro data concerning efficacy and selectivity of compounds of the present invention. Furthermore, binding and cotransfection data for compounds of the present invention are compared with corresponding data for reference compounds that act on either huPPAR α or huPPAR γ . The typical range of concentration for binding is from 1nM to 10 μ M. The concentration of test compound required to effect 50% maximal
15 activation of PPAR α (IC₅₀ α) and PPAR γ (IC₅₀ γ) is determined.

Evaluation of Triglyceride and Cholesterol Level in HuapoAI Transgenic Mice

Five to six week old male mice, transgenic for human apoAI [C57Bl/6-tgn(apoal)1rub, Jackson Laboratory, Bar Harbor, ME] are housed five per cage
20 (10"x20"x8" with aspen chip bedding) with food (Purina 5001) and water available at all times. After an acclimation period of 2 weeks, animals are individually identified by ear notches, weighed and assigned to groups based on body weight. Beginning the following morning, mice are dosed daily by oral gavage for 7 days using a 20 gauge, 1½" curved disposable feeding needle. Treatments are test compounds (30 mg/kg), a positive control
25 (fenofibrate, 100 mg/kg) or vehicle [1% carboxymethylcellulose (w/v)/ 0.25% Tween80 (w/v); 0.2 ml/mouse]. Prior to termination on day 7, mice are weighed and dosed. Three hours after dosing, animals are anesthetized by inhalation of isoflurane (2-4%) and blood obtained via cardiac puncture (0.7-1.0 ml). Whole blood is transferred to serum separator tubes (Vacutainer SST), chilled on ice and permitted to clot. Serum is obtained after
30 centrifugation at 4°C and frozen until analysis for triglycerides, total cholesterol, compound levels and serum lipoprotein profile by fast protein liquid chromatography (FPLC) coupled to an inline detection system. After sacrifice by cervical dislocation, the liver, heart and epididymal fat pads are excised and weighed.

The animals dosed with vehicle have average triglycerides values of about
35 60 to 80 mg/dl, which are reduced by the positive control fenofibrate (33-58 mg/dl with a mean reduction of 37%). The animals dosed with vehicle have average total serum

-60-

5 cholesterol values of about 140 to 180 mg/dl, which are increased by fenofibrate (about 190 to 280 mg/dl with a mean elevation of 41%). When subject to FPLC analysis, pooled sera from vehicle-treated hu apoA1 transgenic mice have a high-density lipoprotein cholesterol (HDLc) peak area which ranges from 47v-sec to 62v-sec. Fenofibrate increases the amount of HDLc (68-96v-sec with a mean percent increase of 48%). Test
10 compounds evaluated in terms of percent increase in the area under the curve. Representative compounds of the present invention are tested using the above methods or substantially similar methods.

Evaluation of Glucose Levels in db/db Mice

15 Five week old male diabetic (db/db) mice [C57BlKs/j-m +/+ Lepr(db), Jackson Laboratory, Bar Harbor, ME] or lean littermates (db+) are housed 6 per cage (10"x20"x8" with aspen chip bedding) with food (Purina 5015) and water available at all times. After an acclimation period of 2 weeks, animals are individually identified by ear notches, weighed and bled via the tail vein for determination of initial glucose levels.
20 Blood is collected (100 µl) from unfasted animals by wrapping each mouse in a towel, cutting the tip of the tail with a scalpel, and milking blood from the tail into a heparinized capillary tube balanced on the edge of the bench. Sample is discharged into a heparinized microtainer with gel separator (VWR) and retained on ice. Plasma is obtained after centrifugation at 4°C and glucose is measured immediately. Remaining plasma is frozen
25 until the completion of the experiment, and glucose and triglycerides are assayed in all samples. Animals are grouped based on initial glucose levels and body weights. Beginning the following morning, mice are dosed daily by oral gavage for 7 days using a 20 gauge, 1½" curved disposable feeding needle. Treatments are test compounds (30 mg/kg), a positive control agent (30 mg/kg) or vehicle [1% carboxymethylcellulose
30 (w/v)/0.25% Tween80 (w/v); 0.3 ml/mouse]. On day 7, mice are weighed and bled (tail vein) for about 3 hours after dosing. Twenty-four hours after the 7th dose (i.e., day 8), animals are bled again (tail vein). Samples obtained from conscious animals on days 0, 7 and 8 are assayed for glucose. After 24 hour bleed, animals are weighed and dosed for the final time. Three hours after dosing on day 8, animals are anesthetized by inhalation
35 of isoflurane, and blood obtained is via cardiac puncture (0.5-0.7 ml). Whole blood is transferred to serum separator tubes, chilled on ice and permitted to clot. Serum is

5 obtained after centrifugation at 4°C and frozen until analysis for compound levels. After sacrifice by cervical dislocation, the liver, heart and epididymal fat pads are excised and weighed.

The animals dosed with vehicle have average triglycerides values of about 170 to 230 mg/dl, which are reduced by the positive PPAR γ control (about 70 to 120
10 mg/dl with a mean reduction of 50%). Male db/db mice are hyperglycemic (average glucose of about 680 to 730 mg/dl on the 7th day of treatment), while lean animals have average glucose levels between about 190 and 230 mg/dl. Treatment with the positive control agent reduces glucose significantly (about 350 to 550 mg/dl with a mean decrease towards normalization of 56%).

15 Glucose is measured colorimetrically by using commercially purchased reagents (Sigma #315-500). According to the manufacturers, the procedures are modified from published work (McGowan et al. *Clin Chem*, 20:470-5 (1974) and Keston, A. Specific colorimetric enzymatic analytical reagents for glucose. Abstract of papers 129th Meeting ACS, 31C (1956).); and depend on the release of a mole of hydrogen peroxide
20 for each mole of analyte coupled with a color reaction first described by Trinder (Trinder, P. *Ann Clin Biochem*, 6:24 (1969)). The absorbance of the dye produced is linearly related to the analyte in the sample. The assays are further modified for use in a 96 well format. Standards (Sigma #339-11, Sigma #16-11, and Sigma #CC0534 for glucose, triglycerides and total cholesterol, respectively), quality control plasma (Sigma # A2034),
25 and samples (2 or 5 μ l/well) are measured in duplicate using 200 μ l of reagent. An additional aliquot of sample, pipetted to a third well and diluted in 200 μ l water, provided a blank for each specimen. Plates are incubated at room temperature (18, 15, and 10 minutes for glucose, triglycerides and total cholesterol, respectively) on a plate shaker and absorbance read at 500 nm (glucose and total cholesterol) or 540 nm (triglycerides) on a
30 plate reader. Sample absorbance is compared to a standard curve (100-800, 10-500, and 100-400 mg/dl for glucose, triglycerides and total cholesterol, respectively). Values for the quality control sample are consistently within the expected range and the coefficient of variation for samples is below 10%. All samples from an experiment are assayed at the same time to minimize inter-assay variability.

-62-

5 Serum lipoproteins are separated and cholesterol is quantitated with an in-line detection system. Sample is applied to a Superose® 6 HR 10/30-size exclusion column (Amersham Pharmacia Biotech) and eluted with phosphate buffered saline-EDTA at 0.5 ml/min. Cholesterol reagent (Roche Diagnostics Chol/HP 704036) at 0.16 ml/min is mixed with the column effluent through a T-connection, and the mixture is

10 passed through a 15 m x 0.5 mm id knitted tubing reactor immersed in a 37°C water bath. The colored product produced in the presence of cholesterol is monitored in the flow stream at 505 nm, and the analog voltage from the monitor is converted to a digital signal for collection and analysis. The change in voltage corresponding to change in cholesterol concentration is plotted against time, and the area under the curve corresponding to the

15 elution of VLDL, LDL and HDL is calculated (Perkin Elmer Turbochrome software).


The compounds of the present invention can be prepared according to the procedures of the following schemes and examples, which may further illustrate details for the preparation of the compounds of the present invention. The compounds illustrated in the schemes and examples are, however, not to be construed as forming the only genus

20 that is considered as the present invention.

General Reaction Scheme:

The compounds of the present invention, in general, may be prepared according to the Reaction Schemes described below. When describing various aspects

25 of the present compounds, the terms "Tail" and "Head" are used as their concept is illustrated below.




-63-

5 Reaction Scheme 1:

Z_1 = leaving group

Head' = modified headpiece to show OH substitution

- 10 As shown in Reaction Scheme I, the compounds of the present invention, in general, can be divided into Tail and Head regions where a nucleophilic headpiece is coupled with an electrophilic tailpiece. These regions can be further modified as shown in the following reaction schemes.

15 Reaction Scheme 2: Oxazole tailpiece

R_1 is H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halo, haloalkyl or haloalkoxy;
 R_2 is Y^{la} ;

20 R_3 is alkyl (methyl) or tolyl.


-64-

5 As shown in Reaction Scheme 2, an intermediate oxazole tailpiece can be prepared by a condensation of dionemonoxime (1) with aldehyde (2) such as bromobenzaldehyde in the presence of acid such as hydrochloric acid or acetic acid to give an oxazole n-oxide compound (3). The oxazole n-oxide is then treated with phosphorous oxychloride in an organic solvent to form chloromethyl substituted-oxazole

10 (4). Compound (4) is further treated with a cyanide to form cyanomethyl oxazole compound (5). The cyano group of compound (5) is converted to a carboxylic acid group by treatment with an alkali metal hydroxide such as NaOH to form carboxymethyl substituted oxazole (6), which is further treated with a carboxylic acid reducing agent, such as borane or lithium aluminum hydride (LAH) to form compound (7). Compound

15 (7) can be converted to oxazolyl sulfonyl ester (8) in the presence of a base by treatment with a sulfonyl halide or sulfonyl anhydride (R_3SO_2Cl or $(R_3SO_2)_2O$), such as tosyl anhydride, mesyl anhydride, tosyl chloride or mesyl chloride.

Reaction Scheme 3: Oxazole tailpiece



20


Alternatively, an intermediate of oxazole tailpiece can be prepared as shown in Reaction Scheme 3. Acid chloride (9) is reacted with L-aspartic acid dimethyl

25 ester (10) to give amide compound (11), which undergoes cyclization to form an oxazole

-65-

- 5 ring (12) by treatment with a dehydrating agent such as P_2O_5 . The ester compound (12) is reduced by treating with LAH to give alcohol (13), which is then converted to oxazolyl sulfonyl ester (8a) as described above in Reaction Scheme 2.


10 Reaction Scheme 4: Oxazole tailpiece




- 15 Another route to an intermediate of oxazole tailpiece is shown in Reaction Scheme 4. Acid chloride (9) and L-aspartic acid monomethyl ester (10) are reacted to give amide compound (11), which is further reacted to give ketone (14). The ketone compound undergoes a cyclization in the presence of dehydrating agent such as $POCl_3$ or H_2SO_4 /acetic anhydride to form oxazole ring (15). Compound (15) undergoes reduction to give alcohol (16), which is then converted to oxazolyl sulfonyl ester (8) as described above in Reaction Scheme 2.

20

-66-

5 Reaction Scheme 5: Oxazole tailpiece


Another route to an intermediate of the oxazole tailpiece is shown in Reaction Scheme 5. The oxazole compound (17) can undergo a coupling reaction in the presence of palladium catalyst with an aryl boronic acid, aryl alcohol or aryl amine followed by deprotection to yield the corresponding compound (18). Compound (18) is then converted to oxazolyl sulfonyl ester (19) as described above in Reaction Scheme 2.

15 Reaction Scheme 6: Thiazole tailpiece

5 As shown in Reaction Scheme 6, an intermediate thiazole tailpiece can be prepared by the condensation of compound (20) with bromo alkyl ester (21) in the presence of 1,4-dioxane followed by cyclization to give thiazole compound (22). The thiazole (22) then undergoes an ester reduction to give alcohol (13), which is further converted to thiazole sulfonyl ester (8) as described above in Reaction Scheme 2.


10

Reaction Scheme 7: Pyrazole tailpiece




15 As shown in Reaction Scheme 7, an intermediate pyrazole tailpiece can be prepared by the condensation of arylaldehyde (25) with compound (26) in the presence of base followed by cyclization to give pyrazole compound (27). Compound (27) is treated with ethylene carbonate in the presence of base such as NaH to give alkylated compound (28), which is then converted to pyrazole sulfonyl ester (29) as described above in Reaction Scheme 2.

20

5 Reaction Scheme 8: Preparation of aryether bromide

Reaction Scheme 8 shows the preparation of aryether bromide by a nucleophilic substitution reaction of an aryl alcohol with a dibromide in the presence of a base.


10 Reaction Scheme 9: Preparation of Aminomethyldihydrocinnamate

$\text{Z}_2 = \text{COR}, \text{CONHR}, \text{CSNHR}, \text{COOR}, \text{SO}_2\text{R};$
where R is selected from Y^7

15 Reaction Scheme 9 shows the synthetic route to prepare aminomethyldihydrocinnamate headpiece. 3-Hydroxybenzaldehyde (29) is reacted with bromine to give 2-bromo-5-hydroxybenzaldehyde (30), which is then coupled to *t*-butylacrylate to give the compound (31). Compound (31) is treated with NH_2OH to give oxime (32), which undergoes hydrogenation to give aminomethyldihydrocinnamate (33).


- 5 Various functional groups (Z_2) can be introduced at the aminomethyl portion of compound (33) to afford compound (34).

Reaction Scheme 10: N-alkyl headpiece




- As shown in Reaction Scheme 10, aminomethyl portion of the headpiece compound (34) can be further modified. Phenol (34) is protected with a protecting group such as a benzyl group to give compound (35). Compound (35) undergoes a nucleophilic substitution reaction with an alkyl halide (RX) under basic conditions followed by a
 15 deprotection to give phenol compound (36).

Reaction Scheme 11: Transesterification



- Reaction Scheme 11 illustrates a transesterification reaction of the compound as shown above. The tert-butyl ester can be cleaved under acidic conditions such as TFA. The corresponding carboxylic acid group can be re-esterified with an appropriate alcohol, such as methanol in the presence of an acid such as H_2SO_4 .
- 25


5 **Reaction Scheme 12: Tail-Head coupling with aminomethyldihydrocinnamates**



As shown in Reaction Scheme 12, headpiece and tailpiece can be modified to introduce various functional groups. In Path A, the protecting group (TBS) is first attached to the phenol of compound (37) and then radical bromination followed by alkylation with potassium phthalimide yields compound (38). Compound (38) undergoes a Heck coupling reaction with tert-butyl acrylate in the presence of palladium catalyst and then deprotection of silyl group followed by hydrogenation to give compound (39). Compound (39) is coupled with a tailpiece and removal of phthaloyl group yields

- 5 compound (40). After neutralization, the amino function of (40) can be modified to give (41). Ester hydrolysis under acidic conditions gives (42). Alternatively, as shown in Path B, compound (42) can be obtained by coupling headpiece (43) with a tailpiece to give compound (41) followed by ester hydrolysis.


10 Reaction Scheme 13: Synthesis of meta-aminomethyldihydrocinnamates



- Reaction Scheme 13 shows a synthesis for meta-substituted aminomethyldihydrocinnamates. A protecting group such as benzyl is attached to the phenol of compound (44) to give protected compound (45). Compound (45) undergoes glycol formation followed by cleavage and oxidation reactions to give carboxylic acid compound (46), which is then converted to amide compound (47). Compound (47) undergoes a rearrangement and subsequent amine protection with Boc anhydride followed by deprotection of benzyl ether to afford phenol (48). Compound (48) is
- 15

- 5 coupled with a tailpiece followed by deprotection to give the intermediate, which then
undergoes further modification at the aminomethyl portion of the compound followed by
hydrolysis to give compound (49).

Reaction Scheme 14: Synthesis of Aminoethyldihydrocinnamates




10

Reaction Scheme 14 shows the synthetic route to prepare aminoethyldihydrocinnamate headpiece. A protecting group such as benzyl is attached to aryl alcohol of compound (50) to give protected compound (51), which is then converted to nitro-olefin compound (52). Compound (52) undergoes reduction of the nitro-olefin by

- 5 treatment with LAH followed by introduction of an amine protecting group such as Boc to give compound (53). Compound (53) undergoes a coupling reaction with methyl acrylate in the presence of a palladium catalyst followed by deprotection of phenol group to give the phenol compound (54). Compound (54) is then coupled with a tailpiece followed by deprotection of Boc group to give the intermediate compound, which then
- 10 can undergo further modification at the aminoethyl portion of the compound followed by hydrolysis to give compound (55). Alternatively, compound (57) is deprotected followed by modification at the aminoethyl portion of the compound to give compound (56). Compound (56) can be coupled with a tailpiece followed by ester hydrolysis to afford compound (55).

15


-74-

5 Reaction Scheme 15: Synthesis of Aminomethylfibrates

Reaction Scheme 15 shows a synthetic route to prepare aminomethylfibrate compounds. The reaction is carried out by following a substantially similar synthetic route as described in Reaction Scheme 13.

5 **Reaction Scheme 16: Synthesis of Carboxamidodihydrocinnamates**

Path A




10 Reaction Scheme 16 shows the synthetic routes to
 carboxamidodihydrocinnamates by Path A or Path B. Path A allows for rapid amide
 variation and Path B for tailpiece variation. In path A, the protecting group such as
 benzyl is attached to the phenol of compound (58) to give protected compound (59).
 Compound (59) is oxidized to give carboxylic acid compound (60), which then undergoes
 15 esterification followed by deprotection of the phenol to give compound (61). Compound
 (61) is coupled with a tailpiece followed by deprotection of carboxylic acid group to give

- 5 compound (62), which then undergoes amide formation followed by ester hydrolysis to afford compound (63). Alternatively as shown in path B, compound (64) can undergo an amide formation followed by reduction and deprotection of phenol group to give compound (65). Compound (65) then can be coupled with a tailpiece followed by ester hydrolysis to yield compound (63).

10


Reaction Scheme 17: Synthesis of Cyanodihydrocinnamates




- 15 Cyanodihydrocinnamates can be prepared as shown in Reaction Scheme 17. Compound (66) is brominated with a brominating agent such as NBS to give compound (67), which is then undergoes a Heck coupling reaction with ethyl acrylate in the presence of palladium catalyst followed by hydrogenation to give compound (68). Compound (68) is coupled with a tailpiece followed by ester hydrolysis to yield
- 20 carboxylic acid compound (69).

5 Reaction Scheme 18: Synthesis of alkoxydihydrocinnamates

Route (a)




Route (b)



- 5 Alkoxydihydrocinnamates can be prepared as shown in Reaction Scheme 18. In route (a), compound (31) is coupled with a tailpiece to give compound (70), which then undergoes sequential reduction of double bond and aldehyde to give alcohol compound (71). Compound (71) undergoes the condensation reaction with an isocyanate or thioisocyanate, alkyl halide or aryl halide followed by ester hydrolysis to give
- 10 compound (72). Alternatively, compound (71) can be converted to the corresponding halide (73) using a carbon tetrahalide and triphenylphosphine. A nucleophilic substitution reaction followed by ester hydrolysis affords compound (74).


- In route (b), the protecting group such as TBDPSi is attached to the aryl alcohol of compound (75) to give compound (76), which then undergoes a similar
- 15 reaction sequence as described in route (a) to give phenol (77). Alternatively, compound (76) can undergo the condensation reaction with Z_3-X followed by deprotection to afford phenol (78). Phenol compounds (77) and (78) can be taken separately to final compounds by reaction with a tailpiece followed by acid hydrolysis.

20 Reaction Scheme 19: Synthesis of Alkoxyalkylfibrates



- 5 Reaction Scheme 19 illustrates a synthetic route to prepare alkoxyalkylfibrate compounds. Compound (79) is reacted with ethyl 2-bromoisobutyrate in the presence of base to give compound (80). The phenol of compound (80) is deprotected followed by aldehyde reduction in the presence of a reducing agent such as NaBH_4 to give compound (81). Phenol (81) then can undergo reaction with $\text{Z}_3\text{-X}$
- 10 followed by hydrolysis to give alkoxyalkylfibrate compound (82). Alternatively, phenol compound (81) can be converted to the corresponding halide (83) using a carbon tetrahalide and triphenylphosphine. Treatment of compound (83) with ArZ_4H followed by ester hydrolysis affords compound (84).


15 Reaction Scheme 20: Modification to the tailpiece



$\text{Ar-Y}^7\text{-Ar}$; Ar-Ar ; $\text{Ar-O-Ar} = \text{Y}^{1a}$
 $\text{R} = \text{H, or alkyl}$

5 Reaction Scheme 20 illustrates the synthetic routes to compounds with modified tailpieces. Oxazole compound (85) can undergo a coupling reaction with an arylboronic, aryl alcohol, aryl amine or secondary amine in the presence of palladium catalyst to give modified tailpiece compound (86). Alternatively, compound (85) undergoes a coupling reaction with pinacol diborane in the presence of palladium catalyst
10 to give compound (87), which then is further coupled with arylhalide to give biaryl compound (88). Alternatively, the compound (87) can be oxidized to give phenol (89), which is then coupled with arylhalide in the presence of palladium catalyst to give aryl-aryloxy compound (90) as shown in route (a). Alternatively, compound (89) can undergo a nucleophilic reaction with an alkyl halide or alcohol as shown in route (b) to form ether
15 compound (91).

Reaction Scheme 21: Modification to the tailpiece




$\text{Ar-C(O)NRR'} = \text{Y}^{1a}$

20 $\text{NR'R} = \text{substituents defined in Y}^{1a}$


Reaction Scheme 21 shows a synthetic route to modify the tailpiece of the present compounds. In route (a), the oxazole tailpiece compound (92) undergoes carbonylation to give carboxylic acid compound (93), which is converted to the acid
25 chloride followed by reaction with an amine to form amide compound (94). Similarly in route (b), modified compound (94) is achieved by palladium catalyzed carbonylation to form an intermediate hydroxypyridine ester followed by a reaction with amine to form amide compound.

-81-


5 Reaction Scheme 22: Modification to the tailpiece

NH-Z^2 = substituents defined in Y^{1a}

Reaction Scheme 22 shows a synthetic route to modify the tailpiece of the present compounds. Compound oxazolyl sulfonate (95) is nitrated to give compound (96), which then undergoes a coupling reaction with a headpiece to give compound (97). The nitro group of the phenyl ring is reduced to give aniline compound, which is modified to give compound (99).

5 Reaction Scheme 23: Modification to the headpiece N

Reaction Scheme 23 shows a synthetic route to modify the headpiece of the present compounds. The headpiece compound (100) can be modified by coupling with arylhalide (Ar-X) in the presence of palladium catalyst followed by ester hydrolysis to give the compound (101). Alternatively, compound (100) can be reacted with Z₂-X in the presence of base to give the modified aminomethyl compound (102). In route (a), compound (102) can undergo nucleophilic substitution at the amine followed by ester hydrolysis to give acid compound (105). Alternatively, by route (b) with Z₂ as a trifluoro acetyl group, alkylation under basic conditions followed by aqueous base hydrolysis gives secondary amine compound (104). Subsequent functionalization of the amine with Z₂-X followed by acid hydrolysis affords compound (105).

5 Reaction Scheme 24: Oxazole tailpiece

$\text{R}_2 = \text{Y}^{\text{Ia}}$ such as aryl, aryl-Z-aryl or heteroaryl-Z-aryl, or alkyl
 $\text{R} = \text{alkyl}$

10

An alternative synthetic route to oxazole tailpiece is shown in Reaction Scheme 24. Carboxylic acid (106) is condensed with 2-bromo-3-oxopentanoate (preferably methyl ester) (107) to give ketoester (108). The latter is converted to an intermediate enamine (109) by treatment with anhydrous ammonium acetate. Subsequent cyclization of compound (109) in acetic acid in the presence of anhydrous ammonium acetate gives compound (110). The use of anhydrous ammonium acetate obtained by azeotropic evaporation with ethanol eliminates water in the reaction, which causes decarboxylation of compound (110). Additionally, some of the water liberated in the reaction is removed at the enamine stage. These modifications along with a simplified isolation procedure lead to higher yields of oxazole (110).

20

In the Schemes, Preparations and Examples below, various reagent symbols and abbreviations have the following meanings:

25	BINAP	2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl
	Boc	t-butoxycarbonyl
	CBZ	benzyloxycarbonyl
	DCM	dichloromethane
	DEAD	diethyl azodicarboxylate
	DI	deionized

5	DIAD	diisopropyl azodicarboxylate
	DIPEA	diisopropylethylamine
	DMAP	4-dimethylamino pyridine
	DMF	N,N-dimethylformamide
	DMSO	dimethylsulfoxide
10	eq. (equiv)	equivalent(s)
	EDC	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide HCl
	ESI-MS	electron spray ion-mass spectroscopy
	Et	ethyl
	EtOAc	ethyl acetate
15	Fmoc	9-Fluorenylmethyl carbamate
	HATU	O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
	HOAc	acetic acid
	HOAT:	1-hydroxy-7-azabenzotriazole
20	HOBT	1-hydroxybenzotriazole hydrate
	HPLC	high performance liquid chromatography
	HRMS	high resolution mass
	h	hour(s)
	LRMS	low resolution mass
25	LAH	lithium aluminum hydride
	Me	methyl
	Ms	methanesulfonyl
	NBS	N-bromosuccinimide
	Pd ₂ (dba) ₃	tris(dibenzylideneacetone) dipalladium(0)
30	Ph	phenyl
	Phe	phenylalanine
	Pr	propyl
	r.t.	room temperature
	TBAF	tetrabutylammonium fluoride
35	TBS	tertbutyldimethylsilyl
	TFA	trifluoroacetic acid

5	TEA	triethylamine
	THF	tetrahydrofuran
	TLC	thin-layer chromatography

Standard Procedure

10 The following standard procedures (A) to (E) are used for preparing the compounds of the present invention as illustrated in Examples below.

Standard Procedure (A): Mitsunobu Coupling of Aryl alcohol Headpieces with Alcohol Tailpieces

15 A mixture of 2-(5-methyl-2-naphthalen-2-yl-oxazol-4-yl)-ethanol (112 mg, 0.442 mmol, 1 equiv), 3-[4-hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid *tert*-butyl ester (150 mg, 0.445 mmol, 1.00 equiv), and triphenylphosphine (116 mg, 0.442 mmol, 1.00 equiv) in toluene (10 mL) at room temperature was treated with diisopropyl azodicarboxylate (90 μ L, 92 mg, 0.46 mmol, 1.0 equiv) over a period of
20 about 3 minutes. The mixture was stirred for about 23 hours and concentrated. The crude material was purified by silica gel chromatography.

Standard Procedure (B): Ester Hydrolysis under Acidic Conditions

25 A solution of *tert*-butyl ester-containing compound (0.5 mmol) in CH_2Cl_2 (6 mL) was treated with 90 % trifluoroacetic acid/water (3 mL) and stirred at ambient temperature for about 3 hours and then concentrated to give the corresponding carboxylic acid compound. The crude material was purified by silica gel chromatography if necessary.

Standard Procedure (C): Ester Hydrolysis under Acidic Conditions

30 A mixture of *tert*-butyl ester-containing compound (0.2 mmol) in 4 M HCl in 1,4-dioxane (5 mL) was stirred at ambient temperature for about 16 hours and concentrated to give the corresponding carboxylic acid compound. The crude material was purified by silica gel chromatography if necessary.

35

-86-

5 Standard Procedure (D): Ester Hydrolysis under Basic Conditions:


A solution of ester (0.25 mmol) in MeOH (3 mL) and THF (1.5 mL) was treated with 2N NaOH (1 mL) and heated at 55°C for about 2 hours. The mixture was cooled and concentrated. The residue was partitioned between CH₂Cl₂ (10 mL), brine (5 mL) and 5N HCl (1 mL). The organic layer was dried (Na₂SO₄) and concentrated. The
10 crude material was purified by silica gel chromatography if necessary.

PREPARATION OF INTERMEDIATES

Tailpiece Preparation - Oxazoles

Preparation 1

15 Toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)ethyl ester

Step A: 4,5-Dimethyl-2-(4-bromophenyl)-oxazole oxide

A solution of 2,3-butanedione monooxime (50 g, 0.49 mol) and 4-bromo-
20 benzaldehyde (101 g, 0.54 mol) in acetic acid (500 mL) was cooled to 0°C and then
gaseous HCl was bubbled through the solution for 35 min while the reaction was stirred
in an ice bath. Diethyl ether (500 mL) was added to the reaction to precipitate the
product, and the resultant slurry was stirred 45 min at 0°C before being filtered. The
solids were rinsed with Et₂O (50 mL), taken up in water (1 L), and conc. NH₄OH (60 mL)
25 was added to the slurry. This mixture was extracted with CHCl₃. The organic layer was
dried (MgSO₄) and concentrated to give 97.4 g (74%) of 4,5-dimethyl-2-(4-
bromophenyl)-oxazole oxide as a white solid. This compound should be used directly
within 24–48 h. ¹H NMR (500 MHz, CDCl₃) 8.34 (d, *J* = 9.0 Hz, 2H), 7.61 (d, *J* = 9.0
Hz, 2H), 2.35 (s, 3H), 2.20 (s, 3H); ¹³C (125 MHz, CDCl₃) 142.1, 131.9, 129.5, 126.3,
30 124.1, 122.2, 11.1, 6.2; IR (KBr) 1685, 1529, 1418, 1377, 1233, 1165 cm⁻¹; UV (EtOH)
_{max} 307 nm (24371); HRMS (TOF) *m/z* calculated for C₁₁H₁₁⁷⁹BrNO₂: 267.997, found
267.9951.

5 Step B: 2-(4-Bromophenyl)-4-(chloromethyl)-5-methyloxazole

A solution of 4,5-dimethyl-2-(4-bromophenyl)-oxazole oxide (96.6 g, 0.36 mol) in CHCl_3 (0.90 L) was treated dropwise with phosphorous oxychloride (61.1 g, 0.40 mol) allowing the reaction to exotherm and then was stirred at reflux for 30 min. The reaction was cooled to room temperature and washed with water (2 x 1 L). The combined aqueous washes were back extracted with CH_2Cl_2 (2 x 400 mL). The organic layers were dried (MgSO_4) and concentrated to give crude product that was recrystallized from hot hexanes (300 mL), decanting the hot supernatant away from a dark oily material. The remaining dark oil was agitated in additional hot hexanes (200 mL), and the combined supernatants were cooled to 0°C . The product was isolated by filtration as a lime-green powder (74.2 g, 72%): $R_f = 0.39$ in 20% ethyl acetate/hexanes. ^1H NMR (500 MHz, CDCl_3) 7.88–7.86 (m, 2H), 7.59–7.56 (m, 2H), 4.54 (s, 2H), 2.42 (s, 3H); ^{13}C (125 MHz, CDCl_3) 159.2, 146.9, 133.2, 132.0, 127.6, 126.1, 124.7, 37.1, 11.5; IR (KBr) 2970, 1633, 1599, 1481, 1401, 1258, 1117, 1008 cm^{-1} ; UV (EtOH) max 281 nm (21349); HRMS (FAB) m/z calculated for $\text{C}_{11}\text{H}_{10}^{79}\text{BrClNO}$: 285.9634, found 285.9641; Anal. Calculated for $\text{C}_{11}\text{H}_9\text{ClBrNO}$: C, 46.11; H, 3.17; N, 4.89; Cl, 12.37; Br, 27.88. Found C, 46.28; H 3.07; N, 4.81; Cl, 12.36; Br, 27.88.

Step C: 2-(4-Bromophenyl)-5-methyl-4-oxazoleacetic acid


To a solution of 2-(4-bromophenyl)-4-(chloromethyl)-5-methyloxazole (64.8 g, 0.23 mol) in DMF (400 mL) was added powdered potassium cyanide (22.1 g, 0.34 mol) and potassium iodide (28.6 g, 0.17 mol), and the resultant mixture was heated to 85°C for 3.5 h. The reaction mixture was cooled to room temperature. Potassium carbonate (5 g) was dissolved in water (800 mL) and added dropwise to the reaction to precipitate 2-(4-bromophenyl)-4-(cyanomethyl)-5-methyloxazole (stir vigorously 15 min following addition) which was isolated by filtration and washed with water (2 x 400 mL). The crude 2-(4-bromophenyl)-4-(cyanomethyl)-5-methyloxazole was used in the next step without purification. ^1H NMR (300 MHz, CDCl_3) 7.85 (m, 2H), 7.58 (m, 2H), 3.64 (s, 3H), 2.43 (s, 3H).

5 The crude 2-(4-bromophenyl)-4-(cyanomethyl)-5-methyloxazole (assume 0.22 mol) was combined with 2-methoxyethanol (630 mL) and 85% solid KOH (74.6 g, 1.33 mol) in water (360 mL) was added to the reaction. The mixture was heated to reflux for 3 h, cooled, quenched with 2 M HCl (500 mL), and extracted with CH₂Cl₂. The organic layer was dried (MgSO₄) and concentrated, using toluene to remove residual 2-methoxyethanol azeotropically. The crude product (57.3 g) was recrystallized from toluene (450 mL) to give 39.8 g (60%) of 2-(4-bromophenyl)-5-methyl-4-oxazoleacetic acid as an off-white powder. R_f = 0.23 in 10% MeOH/CH₂Cl₂; ¹H NMR (500 MHz, CDCl₃) 9.00 (br s, 1H), 7.85–7.83 (m, 2H), 7.58–7.56 (m, 2H), 3.62 (s, 3H), 2.36 (s, 3H); ¹³C (125 MHz, CDCl₃) 173.8, 159.0, 146.2, 132.0, 129.1, 127.6, 125.9, 124.7, 31.5, 10.2; IR (CHCl₃) 2923, 1699, 1641, 1481, 1428, 1306, 1234, 1010, 829, 727 cm⁻¹; UV (EtOH)_{max} 288 nm (19626).

10

15


Step D: 2-(4-Bromophenyl)-5-methyl-4-oxazoleethanol



20 A solution of 2-(4-bromophenyl)-5-methyl-4-oxazoleacetic acid (39.1 g, 0.13 mol) in dry THF (175 mL) was treated dropwise with borane/THF complex (227 mL of a 1.0 M solution in THF, 1.3 mol) over 2 h at about 35°C. After stirring 2 h at room temperature under N₂, the reaction was quenched with slow addition of methanol (60 mL) and stirred overnight at room temperature. The reaction was diluted with 1 N NaOH (50 mL) and extracted with CH₂Cl₂ (2 x 200 mL). The organic layer was washed with H₂O (3 x 100 mL), dried (MgSO₄), and concentrated. The crude product (38.7 g) was recrystallized from toluene (200 mL, wash solid with cold hexanes) to give 26.9 g (72%) of 2-(4-bromophenyl)-5-methyl-4-oxazoleethanol as a white powder. R_f = 0.37 in 10% MeOH/CH₂Cl₂. ¹H NMR (500 MHz, CDCl₃) 7.84–7.82 (m, 2H), 7.57–7.55 (m, 2H), 3.91 (q, J = 5.5 Hz, 2H), 3.14 (t, J = 6 Hz, OH), 2.72 (t, J = 5.5 Hz, 2H), 2.33 (s, 3H); ¹³C (125 MHz, CDCl₃) 158.7, 144.5, 134.2, 131.9, 127.4, 126.4, 124.3, 61.8, 28.1, 10.1; IR (KBr) 3293, 2948, 1642, 15985, 1480, 1472, 1401, 1053, 1003, 836, 734 cm⁻¹; UV (EtOH)_{max} 290 nm (20860); Anal. Calculated for C₁₂H₁₂BrNO₂: C, 51.09; H, 4.29; N, 4.96; Br, 28.32. Found C, 51.31; H 4.06; N, 4.90; Br, 28.19.

25

30

5 Step E: 2-(Biphenyl-4-yl-5-methyl-oxazol-4-yl)ethanol

2-(4-Bromophenyl)-5-methyl-4-oxazoleethanol (10.0 g, 35.0 mmol) and phenylboronic acid (4.5 g, 38.0 mmol) were dissolved in n-propanol (120 mL) before adding triphenylphosphine (165.2 mg, 0.63 mmol), palladium acetate (46 mg, 2.1 mmol), and Na_2CO_3 (4.5 g, 42 mmol dissolved in 30 mL distilled H_2O). The solution was heated at reflux and stirred for 1.5 h. After cooling to ambient temperature, the mixture was concentrated and then partitioned between CH_2Cl_2 (100 mL) and 1N NaOH (100 mL). The aqueous phase was extracted with CH_2Cl_2 (2 x 50 mL). The combined organic phases were dried (MgSO_4) and concentrated under reduced pressure to provide 2-(4-biphenyl)-5-methyl-4-oxazoleethanol (9.5 g, 97% yield) as a white solid which was used directly without further purification. ^1H NMR (500 MHz, CDCl_3) 8.01 (d, 2H), 7.77-7.50 (m, 4H), 7.46 (m, 2H), 7.38 (m, 1H), 3.91 (q, $J = 5.5$ Hz, 2H), 3.18 (t, $J = 6$ Hz, OH), 2.72 (t, $J = 5.5$ Hz, 2H), 2.33 (s, 3H).

20 Step F: Toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)ethyl ester


To a solution of 2-(biphenyl-4-yl-5-methyl-oxazol-4-yl)ethanol (15.8 g, 56.6 mmol) in CH_2Cl_2 (250 mL) at room temperature under N_2 was added pyridine (14.7 g, 185 mmol, 15.0 mL), DMAP (2.03 g, 16.6 mmol), and then tosyl anhydride (24.57 g, 75.2 mmol) portion wise. The reaction exothermed to 32°C and was stirred 30 min before additional tosyl anhydride (2.3 g) was added. The mixture was diluted with CH_2Cl_2 (100 mL) and stirred vigorously with 1N HCl (150 mL) for 15 min. The organic phase was dried (MgSO_4) and filtered through a pad of silica gel (100 mL, packed with CH_2Cl_2). The silica gel was eluted with ethyl acetate (100 mL), and the solution was concentrated to give toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)ethyl ester as a white solid (23.3 g, 95%). $R_f = 0.51$ in 60% ethyl acetate/hexanes. ^1H NMR (400 MHz, CDCl_3) 7.97 (d, 2H), 7.70 (d, 2H), 7.66 (t, 2H), 7.65 (d, 2H), 7.51 (t, 1H), 7.42 (d, 2H), 7.24 (d, 2H), 4.37 (t, 2H), 2.88 (t, 2H), 2.37 (s, 3H), 2.26 (s, 3H).

-90-

5

Preparation 2


Toluene-4-sulfonic acid 2-(4-Bromophenyl-5-methyl-oxazol-4-yl)ethyl ester



The title compound was prepared from 2-(4-bromophenyl)-5-methyl-4-oxazoleethanol according to Procedure 1, Step F: MS (ESI) m/z 436.0 ($M+H$)⁺.

10


The following intermediate compounds are prepared by a substantially similar manner as described in Preparations 1 and 2.

2-(3-Bromophenyl)-5-methyl-4-oxazoleethanol:

15


¹H NMR (300 MHz, d₆-DMSO) δ 7.99 (s, 1H), 7.88 (d, J = 7.7 Hz, 1H), 7.64 (d, J = 7.7 Hz, 1H), 7.44 (t, J = 7.7 Hz, 1H), 4.61 (t, J = 5.5 Hz, OH), 3.63 (q, J = 5.5 Hz, 2H), 2.60 (t, J = 6.6 Hz, 2H), 2.32 (s, 3H);

20

Toluene-4-sulfonic acid 2-(3-bromophenyl-5-methyl-oxazol-4-yl)ethyl ester


¹H NMR (400 MHz, CDCl₃) δ 8.01 (t, J = 1.6 Hz, 1H) 7.80 (dt, J = 7.6 Hz, J = 1.6 Hz, 1H), 7.66 (d, J = 8.4 Hz, 2H), 7.53 (dt, J = 7.6 Hz, J = 1.6 Hz, 1H), 7.30 (t, J = 7.6 Hz, 1H), 7.19 (d, J = 8.4 Hz, 2H), 4.30 (t, J = 6.4 Hz, 2H), 2.82 (t, J = 6.4 Hz, 2H), 2.31 (s, 3H), 2.24 (s, 3H); MS (ESI) m/z 436.0 ($M+H$)⁺.

25

5 Toluene-4-sulfonic acid 2-(2-biphenyl-3-yl-5-methyl-oxazol-4-yl)ethyl ester


¹H NMR (300 MHz, CDCl₃) δ 7.99 (s, 1H), 7.80 (d, J = 8.0 Hz, 1H), 7.64 (d, 8.0 Hz, 1H), 7.51 (d, J = 9.0 Hz, 1H), 7.30 (t, J = 8.0 Hz, 1H), 7.20 (d, J = 9.0 Hz, 1H), 4.30 (t, J = 7.0 Hz, 2H), 2.80 (t, J = 7.0 Hz, 2H), 2.30 (s, 3H), 2.23 (s, 3H).

10

2-(5-Methyl-2-thiophen-2-yl-4-oxazoleethanol


¹H NMR (500 MHz, CDCl₃): δ 7.54 (m, 1H), 7.33 (m, 1H), 7.03 (m, 1H), 3.87 (t, J = 5.8 Hz, 2H), 3.5 (s, 1H), 2.67 (t, J = 5.8 Hz, 2H), 2.25 (s, 3 H)


15

Toluene-4-sulfonic acid 2-(5-methyl-2-thiophen-2-yloxazol-4-yl)ethyl ester


¹H NMR (400 MHz, CDCl₃): δ 7.67 (d, J = 8.3 Hz, 2H), 7.51 (dd, J = 3.8, 1.4 Hz, 1H), 7.37 (dd, J = 4.9, 1.2 Hz, 1H), 7.21 (d, J = 7.9 Hz, 2H), 7.08 (dd, J = 4.8, 3.5 Hz, 1H), 4.28 (t, J = 6.3 Hz, 2H), 2.80 (t, J = 6.3 Hz, 2H), 2.28 (s, 3H), 2.26 (s, 3 H); mp 107-109°C.

20


-92-

5 2-[2-(4-Benzoyloxy-phenyl)-5-methyl-oxazol-4-yl]-ethanol

¹H NMR (500 MHz, CDCl₃) δ 7.91 (d, 2H, J = 8.60 Hz), 7.45-7.34 (m, 5H), 7.02 (d, 2H, J = 8.60 Hz), 5.11 (s, 2H), 3.91 (t, 2H, J = 5.7 Hz), 2.71 (t, 2H, J = 5.7 Hz), 2.31 (s, 3H); MS (ES⁺) Calculated for C₁₉H₂₀NO₃: Found m/e 310 (M + 1, 100%)

10

Toluene-4-sulfonic acid 2-[2-(4-benzyloxy-phenyl)-5-methyl-oxazol-4-yl]-ethyl
ester





¹H NMR (500 MHz, CDCl₃) δ 7.80-7.78 (m, 2H), 7.67-7.65 (m, 2H), 7.45-7.34 (m, 5H), 7.25-7.17 (m, 2H), 7.02-6.99 (m, 2H), 5.12 (s, 2H), 4.29 (t, 2H, J = 6.45 Hz), 2.80 (t, 2H, J = 6.45 Hz), 2.27 (s, 3H), 2.22 (s, 3H); HRMS (ES⁺) m/z exact mass calculated for C₂₆H₂₆NO₅S 464.1532, found 464.1531; Anal. Calculated for C₂₆H₂₅NO₅S: C, 67.37; H, 5.44; N, 3.02. Found C, 66.59; H 5.33; N, 3.06.

20

5

Preparation 3

Toluene-4-sulfonic acid 2-(5-methyl-2-phenethyl-oxazol-4-yl)-ethyl ester

Step A: 2-(3-Phenyl-propionylamino)-succinic acid 4-methyl ester

10


Methyl L-aspartate (15.0 g, 0.082 mol), DI water (245 mL), acetone (20 mL), and Na_2CO_3 (30.8 g, 0.286 mol) were combined and cooled the solution to 5°C.

The compound 3-phenyl-propionyl chloride (13.3 mL, 0.089 mol) was added dropwise via addition funnel over 10 min. The reaction was allowed to warm to ambient temperature and stir for 2 h. Conc. HCl (50 mL) was added to the thick slurry until the

15

pH was ≤ 4.0 . The reaction mixture was extracted with CH_2Cl_2 (3 x 100 mL). The combined organic layers were washed with water, dried (MgSO_4), filtered, and concentrated. The clear, colorless oil was used without further purification. ^1H NMR (400 MHz, CDCl_3) δ 7.92 (br s, 1H), 7.28-7.17 (m, 5H), 6.57 (d, $J = 7.6$ Hz, 1H), 4.87 (m, 1H), 3.67 (s, 3H), 2.96 (t, $J = 7.6$ Hz, 2H), 2.89 (A of ABX, $J_{AB} = 17.6$ Hz, $J_{AX} = 4.8$ Hz, 1H), 2.88 (B of ABX, $J_{BA} = 17.6$ Hz, $J_{BX} = 4.0$ Hz, 1H), 2.69 (t, $J = 7.6$ Hz, 2H); MS (EI+) 280 (M+H), 302 (M+H+Na).

20

Step B: 4-Oxo-3-(3-phenyl-propionylamino)-pentanoic acid methyl ester


25

2-(3-Phenyl-propionylamino)-succinic acid 4-methyl ester (10 g, 36 mmol), pyridine (50 mL) and acetic anhydride (45 mL) were combined in a 500 mL flask. The reaction mixture was heated at 90°C for 2 h and then cooled to ambient temperature. After concentrating the reaction mixture under reduced pressure, DI water was added

- 5 (100 mL). The reaction mixture was partitioned between water and CH_2Cl_2 (200 mL). The organic phase was washed with 1N HCl (50 mL), dried (MgSO_4), filtered, and concentrated. The material was used without further purification. ^1H NMR (400 MHz, CDCl_3) δ 7.31-7.20 (m, 5H), 6.79 (br d, $J = 7.6$ Hz, 1H), 4.72 (X of ABX, 1H), 3.65 (s, 3H), 3.01-2.93 (m, 3H), 2.71-2.62 (m, 3H), 2.11 (s, 3H); MS (EI) 278.1 (M+H).


10

Step C: (5-Methyl-2-phenethyl-oxazol-4-yl)-acetic acid methyl ester



- In a 100 mL flask, 4-oxo-3-(3-phenyl-propionylamino)-pentanoic acid methyl ester (10 g, 36 mmol) and acetic anhydride (28 mL) were combined. Following addition of concentrated H_2SO_4 (1 mL), the solution was heated to 90°C for 30 min and cooled to ambient temperature. The reaction was slowly diluted with DI water (30 mL, potential exotherm). The reaction mixture was partitioned between CH_2Cl_2 (150 mL) and water (150 mL). The organic phase was washed with DI water, 10% NaHCO_3 (aq), brine (150 mL), and then was dried (MgSO_4) and concentrated to a brown oil. The residue was purified by column chromatography (600 mL SiO_2 , 35% EtOAc/hexanes) to provide the desired product (3.25 g) as a pale yellow oil. ^1H NMR (400 MHz, CDCl_3) δ 7.33-7.20 (m, 5H), 3.72 (s, 3H), 3.47 (s, 2H), 3.08-2.96 (m, 4H), 2.24 (s, 3H); MS (EI+) 260 (M+H).


- 25 Step D: (5-Methyl-2-phenethyl-oxazol-4-yl)-acetic acid



- (5-Methyl-2-phenethyl-oxazol-4-yl)-acetic acid methyl ester (8.75 g, 33.8 mmol), in MeOH (120 mL) was treated with 5N NaOH (40 mL), and then the solution was warmed to 40°C . After 40 min, the reaction mixture was concentrated. The residue was suspended in water (75 mL) and acidified to pH=1 with 5N HCl. The mixture was extracted with EtOAc (2x), dried (MgSO_4), and concentrated to provide 5.25 g (63%) of

5 the product as an off-white solid. ^1H NMR (400 MHz, CDCl_3) δ 7.33-7.20 (m, 5H), 3.52 (s, 2H), 3.06-3.03 (m, 4H), 2.24 (s, 3H).

Step E: 2-(5-Methyl-2-phenethyl-oxazol-4-yl)-ethanol



10 BH_3 -THF complex (49 mL of a 1.0 M solution in THF) was added dropwise via addition funnel over 50 min to a solution of (5-methyl-2-phenethyl-oxazol-4-yl)-acetic acid (5.05 g, 20.6 mmol) in THF (35 mL). The reaction mixture was stirred at ambient temperature for 3 h, and then quenched with MeOH (12 mL). After heating at 50 °C for 2 h, the reaction mixture was cooled to ambient temperature, and then
15 partitioned between CH_2Cl_2 and 1N NaOH. The organic phase was washed with brine (1x 50 mL), dried over MgSO_4 and concentrated to obtain a residue, which was purified by column chromatography (500 mL SiO_2 , 35% EtOAc/hexanes) to provide 3.99 g (84%) of the desired product as a clear, colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 7.33-7.20 (m, 5H), 3.84 (q, J = 5.6 Hz, 2H), 3.06-2.67 (m, 4H), 2.62 (t, J = 5.6 Hz, 2H), 2.22 (s, 3H); MS (EI+) 232.19 (M+H); 254.15 (M+H+Na).
20


Step F: Toluene-4-sulfonic acid 2-(5-methyl-2-phenethyl-oxazol-4-yl)-ethyl ester

A solution of 2-(5-methyl-2-phenethyl-oxazol-4-yl)-ethanol (1.2 g, 5.19 mmol) in CH_2Cl_2 at 0°C was treated with pyridine (1.64 g, 20.7 mmol, 1.68 mL), DMAP (190 mg, 1.56 mmol), and tosyl anhydride (2.2 g, 6.75 mmol). The reaction was warmed
25 to ambient temperature and, after 90 min, the solution was filtered through a pad of silica gel (rinsed with CH_2Cl_2). The product was used without further purification. ^1H NMR (400 MHz, CDCl_3) δ 7.73 (d, J = 8.4 Hz, 2H), 7.31-7.17 (m, 7H), 4.21 (t, J = 6.8 Hz, 2H), 3.01-2.88 (m, 4H), 2.75 (t, J = 6.8 Hz, 2H), 2.43 (s, 3H), 2.19 (s, 3H).
30

-96-


- 5 The following intermediate compounds are prepared by a substantially similar manner as described in Preparations 3.

2-(2-Cyclohexyl-5-methyl-oxazol-4-yl)-ethanol:




- 10 ^1H NMR (400 MHz, CDCl_3) δ 3.73 (t, $J = 6.8$ Hz, 2H), 2.58 (tt, $J = 11.6, 3.6$ Hz, 1H), 2.54 (t, $J = 6.8$ Hz, 2H), 2.13 (s, 3H), 1.93-1.89 (m, 2H), 1.74 (dt, $J = 12.8, 3.6$ Hz, 2H), 1.67-1.62 (m, 1H), 1.41 (qd, $J = 12.0, 3.2$ Hz, 1H), 1.33-1.17 (m, 4H); MS (EI+) 210.1 (M+H).

- 15 Toluene-4-sulfonic acid 2-(2-cyclohexyl-5-methyl-oxazol-4-yl)-ethyl ester



- ^1H NMR (400 MHz, CDCl_3) δ 7.67 (d, $J = 8.4$ Hz, 2H), 7.27 (d, $J = 8.4$ Hz, 2H), 4.16 (t, $J = 6.8$ Hz, 2H), 2.70 (t, $J = 6.8$ Hz, 2H), 2.56 (tt, $J = 11.6, 3.6$ Hz, 1H), 2.39 (s, 3H), 2.13 (s, 3H), 1.93-1.89 (m, 2H), 1.74 (dt, $J = 12.8, 3.6$ Hz, 2H), 1.67-1.62 (m, 1H), 1.41 (qd, $J = 12.0, 3.2$ Hz, 1H), 1.33-1.17 (m, 4H); MS (EI+) 364.1 (M+H) $^+$.
- 20

2-[5-Methyl-2-(1-methylcyclohexyl)oxazol-4-yl]ethanol




MS (EI+) 224.1 (M+H) $^+$.

25


-97-

5 Toluene-4-sulfonic acid 2-[5-methyl-2-(1-methylcyclohexyl)oxazol-4-yl]ethyl
ester



¹H NMR (400 MHz, CDCl₃) δ 7.73 (d, J = 8.4 Hz, 2H), 7.30 (t, J = 8.4 Hz, 2H), 4.12 (t, J = 6.6 Hz, 2H), 2.76 (t, J = 6.6 Hz, 2H), 2.42 (s, 3H), 2.17 (s, 3H), 2.61-2.02 (m, 2H),
 10 1.56-1.30 (m, 8H), 1.19 (s, 3H); MS (EI) 378.2 (M + H)⁺.


2-[5-Methyl-2-(tetrahydro-pyran-4-yl)-oxazol-4-yl]-ethanol



MS (EI) 212.2 (M + H)⁺.

15


Toluene-4-sulfonic acid 2-[5-methyl-2-(tetrahydro-pyran-4-yl)-oxazol-4-yl]-ethyl
ester



MS (EI) 366.2 (M + H)⁺.

20

2-(2-Benzyl-5-methyl-oxazol-4-yl)-ethanol





MS (EI) 218.0 (M + H)⁺.

25

-98-


5

Toluene-4-sulfonic acid 2-(2-benzyl-5-methyl-oxazol-4-yl)-ethyl esterMS (EI) 372.1 (M + H)⁺.2-(2-Benzo[b]thiophen-2-yl-5-methyl-oxazol-4-yl)-ethanol

10


¹H NMR (CDCl₃) δ 7.81 (m, 3 H), 7.38 (m, 2 H), 3.94 (m, 2 H), 3.07 (br s, 1 H), 2.73 (t, 2 H, *J* = 6 Hz), 2.34 (s, 3 H); ¹³C NMR (CDCl₃) δ 155.9, 145.0, 140.5, 139.8, 134.5, 129.9, 125.9, 125.1, 124.7, 123.7, 122.7, 61.9, 28.5, 10.4; MS (EI) 260.1 (M + H)⁺.

15





Toluene-4-sulfonic acid 2-(2-benzo[b]thiophen-2-yl-5-methyl-oxazol-4-yl)-ethyl ester

20



¹H NMR (CDCl₃) δ 7.85 (m, 1 H), 7.84 (m, 1 H), 7.75 (s, 1 H), 7.67 (d, 2 H, *J* = 8 Hz), 7.39 (m, 2 H), 7.21 (m, 2 H), 4.31 (t, 2 H, *J* = 2 Hz), 2.83 (t, 2 H, *J* = 6 Hz), 2.32 (s, 3 H), 2.19 (s, 3 H).

2-(5-Methyl-2-naphthalen-2-yl-oxazol-4-yl)-ethanolHRMS Calcd for C₁₆H₁₆NO₂: *m/z* 254.1181. Found: 254.1167.

-99-

5 2-[5-Methyl-2-(4-trifluoromethyl-phenyl)-oxazol-4-yl]-ethanolMS (EI) 272 (M + H)⁺.10 Toluene-4-sulfonic acid 2-[5-methyl-2-(4-trifluoromethyl-phenyl)-oxazol-4-yl]-ethyl esterMS (EI) 426 (M + H)⁺; mp 110.2°C.15 2-[2-(4-Butoxy-phenyl)-5-methyl-oxazol-4-yl]-ethanolMS (EI) 276 (M + H)⁺.20 Toluene-4-sulfonic acid 2-[2-(4-butoxy-phenyl)-5-methyl-oxazol-4-yl]-ethyl esterMS (EI) 430 (M + H)⁺; mp 84.9°C.


-100-

5 2-(2-Bromophenyl-5-methyl-oxazol-4-yl)-ethanolMS (EI) 282.1 (M + H)⁺.Toluene-4-sulfonic acid 2-(2-bromophenyl-5-methyl-oxazol-4-yl)ethyl ester


10

MS (EI) 438.1 (M + H)⁺.Preparation 4


Toluene-4-sulfonic acid 2-[2-(6-chloro-pyridin-3-yl)-5-methyl-oxazol-4-yl]-ethyl ester



15

Step A: 3-[2-(6-Chloro-pyridin-3-yl)-5-methyl-oxazole-4-yl]-acetic acid methyl ester

According to Preparation 3, Steps A to C, 6-chloronicotinic acid was converted into the title compound. MS (ESI) m/z 267 (M+H)⁺.

5 Step B: 3-[2-(6-Chloro-pyridin-3-yl)-5-methyl-oxazole-4-yl]-ethanol

A solution of 3-[2-(6-chloro-pyridin-3-yl)-5-methyl-oxazole-4-yl]-acetic acid methyl ester (500 mg, 1.88 mmol) in THF (20 mL) at 0°C was treated LAH (90 mg, 2.3 mmol). The reaction mixture was stirred for 1 h and was quenched with water (0.1 mL), 15% NaOH (0.1 mL), and water (0.3 mL). The mixture was filtered through Celite and concentrated to give the title alcohol which as used in the next step without further purification. MS (ESI) m/z 239 ($M+H$)⁺.


15 Step C: Toluene-4-sulfonic acid 2-[2-(6-chloro-pyridin-3-yl)-5-methyl-oxazol-4-yl]-ethyl ester

A solution of crude 3-[2-(6-chloro-pyridin-3-yl)-5-methyl-oxazole-4-yl]-ethanol (1.88 mmol max) in CH₂Cl₂ (10 mL) was treated with *para*-toluenesulfonyl chloride (0.4 g, 2.3 mmol), DMAP (40 mg), and triethylamine (0.4 mL, 2.82 mmol). The reaction mixture was stirred at ambient temperature overnight and was diluted with CH₂Cl₂ (20 mL). The mixture was washed with water, and the organic layer was dried (MgSO₄), filtered, and concentrated. The crude product was purified by silica gel chromatography (hexanes/EtOAc 10/1 to 2/1) to afford the title compound (295 mg, 40% over two steps). MS (ESI) m/z 393 ($M+H$)⁺.


5

Preparation 5

Toluene-4-sulfonic acid 2-{5-methyl-2-[4-(methyl-phenyl-amino)-phenyl]-oxazol-4-yl}-ethyl ester



Step A: 4-(2-Benzyloxy-ethyl)-2-(4-bromo-phenyl)-5-methyl-oxazole




10

A solution of 2-[2-(4-bromo-phenyl)-5-methyl-oxazol-4-yl]-ethanol (3.17 g, 11.2 mmol) in DMF (25 mL) was treated with NaH (0.67 g, 60% oil dispersion) at 0°C and stirred for 5 min. Benzyl bromide (2.90 g, 16.9 mmol) was added, and the resulting mixture was stirred at room temperature for 3 h. The reaction was quenched with water, and the mixture was extracted with EtOAc (2 x 150 mL). The combined organics were dried (Na₂SO₄), concentrated, and purified by silica gel chromatography column (10% EtOAc/hexanes) to yield the title compound as an oil (2.50 g, 60%).

15


Step B: {4-[4-(2-Benzyloxy-ethyl)-5-methyl-oxazol-2-yl]-phenyl}-methyl-phenyl-amine



20

A solution of 4-(2-benzyloxy-ethyl)-2-(4-bromo-phenyl)-5-methyl-oxazole (200 mg, 0.538 mmol) in toluene (5.0 mL) in a seal tube under nitrogen gas flow was treated with Pd(OAc)₂ (50 mg), 2-(di-*t*-butylphosphino)biphenyl (20 mg), *N*-methyl aniline (115 mg, 1.08 mmol), and sodium *t*-butoxide (104 mg, 1.08 mmol). The tube was sealed and heated at 105 °C for 14 h. The mixture was cooled and purified directly by silica gel column chromatography (30-50% EtOAc/hexanes) to yield the title compound (195 mg, 91%). MS (ESI) *m/z* 399.3 (M+H)⁺.

25

5 Step C: 2-{5-Methyl-2-[4-(methyl-phenyl-amino)-phenyl]-oxazol-4-yl}-ethanol


A solution of {4-[4-(2-benzyloxy-ethyl)-5-methyl-oxazol-2-yl]-phenyl}-methyl-phenyl-amine (195 mg, 0.490 mmol) in THF (2 mL) and EtOH (10 mL) was treated a slurry of Pd/C (200 mg) in EtOH (2 mL). The resulting mixture was treated with hydrogen under balloon pressure for 14 h and filtered through a pad of Celite. The filtrate was concentrated, and crude product was purified by silica gel chromatography column (50% EtOAc/hexanes) to yield the title compound (91 mg, 60%).

15 Step D: Toluene-4-sulfonic acid 2-{5-methyl-2-[4-(methyl-phenyl-amino)-phenyl]-oxazol-4-yl}-ethyl ester

A solution of 2-{5-methyl-2-[4-(methyl-phenyl-amino)-phenyl]-oxazol-4-yl}-ethanol (91 mg, 0.30 mmol) in CH₂Cl₂ (4.0 mL) was treated with *para*-toluenesulfonyl chloride (68 mg, 0.36 mmol), triethyl amine (0.20 mL) and a few crystals of DMAP. The resulting mixture was stirred at room temperature for 14 h and was quenched with water (0.2 mL). The mixture was purified directly by silica gel column chromatography (40% EtOAc/hexanes) to yield the title compound (120 mg, 83%). MS (ESI) *m/z* 463.1 (M+H)⁺.


- 5 The following intermediate compounds are prepared by a substantially similar manner as described in Preparation 5.

Toluene-4-sulfonic acid 2-[5-methyl-2-(4-phenylamino-phenyl)-oxazol-4-yl]-ethyl ester



- 10 MS (ESI) m/z 449.1 ($M+H$)⁺.


Toluene-4-sulfonic acid 2-[5-methyl-2-(4-morpholin-4-yl-phenyl)-oxazol-4-yl]-ethyl ester




- 15 MS (ESI) m/z 443.1 ($M+H$)⁺.

Preparation 6

Toluene-4-sulfonic acid 2-[5-methyl-2-(4-phenoxy-phenyl)-oxazol-4-yl]-ethyl ester



- 20 Step A: 4-(2-Benzyloxy-ethyl)-5-methyl-2-(4-phenoxy-phenyl)-oxazole



- A mixture of 4-(2-benzyloxy-ethyl)-2-(4-bromo-phenyl)-5-methyl-oxazole (0.025 mol, 9.2 g), phenol (0.03 mol, 2.8 g), K_3PO_4 (0.05 mol, 10.6 g), 2-(di-*tert*-butylphosphino)biphenyl (1.8 mmol, 0.54 g) and $Pd(OAc)_2$ (1.2 mmol, 0.28 g) in toluene (350 mL) was degassed with nitrogen and heated at 100°C for 18 h. Additional $Pd(OAc)_2$ (0.5 g) and phosphine ligand (1.0 g) were added, and the mixture was heated 5 h at
- 25

-105-

- 5 100°C. The reaction was concentrated and purified directly by silica gel chromatography (4/1 hexanes/ethyl acetate) to give the title compound (7.6 g).


Step B: Toluene-4-sulfonic acid 2-[5-methyl-2-(4-phenoxy-phenyl)-oxazol-4-yl]-ethyl ester

- 10 According to Preparation 5, Steps C to D, 4-(2-benzyloxy-ethyl)-5-methyl-2-(4-phenoxy-phenyl)-oxazole was converted into the title compound. ¹HNMR (400 MHz, CDCl₃) δ 7.81 (d, 2H, J=9.1 Hz), 7.67 (d, 2H, J=8.2 Hz), 7.37 (t, 2H, J=8.2 Hz), 7.15 (m, 3H), 7.12 (m, 4H), 4.39 (t, 2H, J=6.4 Hz), 2.81 (t, 2H, 6.4 Hz), 2.30 (s, 3H), 2.25 (s, 3H).

15

Preparation 7

4-Methyl-3-nitro-benzenesulfonic acid 2-[5-methyl-2-(4-nitro-phenyl)-oxazol-4-yl]-ethyl ester





- 20 A mixture of potassium nitrate (3.0 g, 30 mmol, 2.7 equiv) and sulfuric acid (10 mL, 18 g, 94 mmol, 17 equiv) was cooled to 0°C. Toluene-4-sulfonic acid 2-(5-methyl-2-phenyl-oxazol-4-yl)-ethyl ester (4.00 g, 11.2 mmol, 1 equiv) was added and the ice bath was removed. The reaction mixture was heated with a heat gun until the tosylate dissolved. After 30 min, the solution was poured into H₂O (100 mL) and extracted with
- 25 EtOAc (100 mL). The organic layer was dried (Na₂SO₄) and concentrated (75°C) to an orange oil (4.41 g). The crude product was purified by silica gel flash chromatography (30-50% EtOAc/hexanes) to give the title compound as a yellow solid (3.64 g, 73%). MS (ESI) m/z 447 (M+H)⁺.

30

5

Preparation 8


Toluene-4-sulfonic acid 2-(5-methoxy-2-phenyl-oxazol-4-yl)-ethyl ester

Step A: 2-Benzoylamino-succinic acid dimethyl ester

10

A mixture of benzoyl chloride (3.20 mL, 27.7 mmol), L-aspartic acid dimethyl ester (5.0 g, 25.2 mmol) and triethyl amine (5.3 mL, 38 mmol) in CH₂Cl₂ (50 mL) was stirred at ambient temperature for 2 h and diluted with water. The organic layer was dried (MgSO₄), filtered, and concentrated. The residue was purified by silica gel chromatography (hexanes/EtOAc 1/1) to afford a white solid (5.3 g, 79%). MS (ESI) m/z 266 (M+H)⁺.


15

Step B: 3-(5-Methoxy-2-phenyl-oxazol-4-yl)-acetic acid methyl ester

A mixture of 2-benzoylamino-succinic acid dimethyl ester (5.3 g, 20 mmol) in 1,2-dichloroethane (15 mL) was treated with P₂O₅ (5.3 g, 30 mmol) and Celite (3.2 g) and was heated at 85°C for 2 h. The solvent was decanted and concentrated. The residue was dissolved in CH₂Cl₂ and washed with saturated aqueous NaHCO₃. The organic layer was dried (MgSO₄), filtered, and concentrated. The residue was purified by silica gel chromatography (hexanes/EtOAc 10/1 to 3/1) to afford the title compound (2.9 g, 59%). MS (ESI) m/z 247 (M+H)⁺.

25

-107-

5 Step C: 3-(5-Methoxy-2-phenyl-oxazol-4-yl)-ethanol

A suspension of LAH (0.56 g, 14.1 mmol) in THF (100 mL) at -78°C was treated dropwise with a solution of 3-(5-methoxy-2-phenyl-oxazol-4-yl)-acetic acid methyl ester (2.9 g, 11.7 mmol) in THF (100 mL). After the addition was completed, the reaction mixture was warmed up to ambient temperature, cooled to -20°C, and quenched with H₂O (0.8 mL), 15% NaOH (0.8 mL), and H₂O (2.4 mL). The mixture was filtered through Celite and concentrated to the title compound as an oil. MS (ESI) m/z 220.3 (M+H)⁺.


15 Step D: Toluene-4-sulfonic acid 2-(5-methoxy-2-phenyl-oxazol-4-yl)-ethyl ester

A solution of crude 3-(5-methoxy-2-phenyl-oxazol-4-yl)-ethanol (11.7 mmol max) in CH₂Cl₂ (100 mL) was treated with *para*-toluenesulfonyl chloride (2.7 g, 14.0 mmol), DMAP (100 mg), and triethylamine (2.5 mL, 17.6 mmol). The reaction mixture was stirred at ambient temperature for 16h and was washed with water. The organic layer was dried (MgSO₄), filtered, and concentrated. The residue was purified by silica gel chromatography (hexanes/EtOAc, 10/1 to 1/1) to afford the title compound (2.0 g, 46% over two steps). MS (ESI) m/z 374 (M+H)⁺.

-108-

5 The following intermediate compounds are prepared by a substantially similar manner as described in Preparation 8.


2-(2-Biphenyl-4-yl-5-methoxy-oxazol-4-yl)-ethanol



MS (ESI) m/z 296.0 ($M+H$)⁺.

10

Toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methoxy-oxazol-4-yl)-ethyl ester




MS (ESI) m/z 450.1 ($M+H$)⁺.


15

Preparation 9

Toluene-4-sulphonic acid 2-[5-methyl-2-(6-phenyl-pyridin-3-yl)thiazol-4-yl]ethyl ester



Step A: 2-Phenyl-5-cyanopyridine




20

5-Cyano-2-chloropyridine (5.0 g, 36.1 mmol), phenylboronic acid (6.6 g, 54 mmol), tetrakis(triphenylphosphine) palladium (0) (0.5 g), and aqueous Na₂CO₃ (7.6 g), in toluene (100 mL) were heated at 90°C for 16 h. The mixture was diluted with

-109-

- 5 EtOAc and washed with H₂O. The organic layer was dried (MgSO₄), filtered, and concentrated. The residue was purified by silica gel chromatography (hexanes/EtOAc 2/1) to afford the title compound (6.1 g, 94%). MS (ESI) m/z 181(M+H)⁺.


Step B:



10


- A mixture of 2-phenyl-5-cyanopyridine (6.0 g, 33 mmol) and thioacetamide (4.0 g, 53 mmol) in 4N HCl in 1,4-dioxane (50 mL) was heated at 98°C for 20 h. The reaction mixture was cooled and poured into aqueous saturated NaHCO₃. The precipitate was collected, washed with water, and dried under vacuum (60°C) to afford the title compound as a yellow solid (7.0 g, 99%).
- 15

Step C: [5-Methyl-2-(6-phenyl-pyridin-3-yl)-thiazol-4-yl]-acetic acid methyl ester



- A mixture of 6-phenyl-thionicotinamide (7.0 g) and 4-bromo-3-oxo-pentanoic acid methyl ester (9.15 g, 35 mmol) in 1,4-dioxane (30 mL) was heated at reflux for 4 h. The reaction mixture was cooled, poured into aqueous saturated NaHCO₃, and extracted with CH₂Cl₂. The organic layer was dried (MgSO₄), filtered, and concentrated. The residue was purified by silica gel chromatography (hexanes/EtOAc, 2/1) to afford the title compound (6.0 g, 56%). MS (ESI) m/z 325 (M+H)⁺.
- 20
- 25

-110-


5 Step D: [5-Methyl-2-(6-phenyl-pyridin-3-yl)-thiazol-4-yl]-ethanol

A solution of [5-methyl-2-(6-phenyl-pyridin-3-yl)-thiazol-4-yl]-acetic acid methyl ester (6.0 g, 18.5 mmol) in THF (500 mL) was added dropwise to a suspension of LAH (0.90 g, 22.2 mmol) in THF (300 mL) at -78°C. After the addition was completed, the reaction mixture was allowed to warm to ambient temperature, cooled to -20 °C, and quenched sequentially with H₂O (1.1 mL), 15% NaOH (1.1 mL), and H₂O (3.3 mL). The mixture was filtered through Celite, and the filtrate was concentrated to give the title compound as an oil that was used directly in the next step.

15 Step E: Toluene-4-sulponic acid 2-[5-methyl-2-(6-phenyl-pyridin-3-yl)thiazol-4-yl]ethyl ester

A mixture of [5-methyl-2-(6-phenyl-pyridin-3-yl)-thiazol-4-yl]-ethanol (18.5 mmol max), *para*-toluenesulfonyl chloride (3.89 g, 20.5 mmol), DMAP (500 mg), and triethylamine (4.0 mL, 28.0 mmol) in CH₂Cl₂ (300 mL) was stirred at ambient temperature for 2.5 h. The reaction mixture was diluted with water, and the organic layer was separated, dried (MgSO₄), filtered, and concentrated. The residue was purified by silica gel chromatography (hexanes/EtOAc, 10/1 to 1/1) to afford the title compound as a solid (2.0 g, 46% over two steps). MS (ESI) *m/z* 451 (M+H)⁺.


25 The following intermediate compounds are prepared by a substantially similar manner as described in Preparation 9.

2-[5-Methyl-2-(5-phenyl-pyridin-3-yl)-thiazol-4-yl]-ethanol

MS (ESI) *m/z* 297 (M+H)⁺.


-111-

5 Toluene-4-sulfonic acid 2-[5-methyl-2-(5-phenyl-pyridin-3-yl)thiazol-4-yl]ethyl
ester



MS (ESI) m/z 451 ($M+H$)⁺.


10 Toluene-4-sulfonic acid 2-[5-methyl-2-(6-phenoxy-pyridin-3-yl)-thiazol-4-yl]-ethyl ester



¹HNMR 400 MHz (CDCl₃) δ 8.52 (2, 1H), 8.03 (d, 1H, J=6.9 Hz), 7.63 (d, 2H, J=8.6 Hz), 7.42 (t, 2H, J=7.7 Hz), 7.2 (m, 5H), 6.91 (1H, d, J=7.7 Hz), 4.37 (t, 2H, J=6.3 Hz), 3.02 (t, 2H, J=6.3 Hz), 2.39 (s, 3H), 2.28 (s, 3H).

15


Toluene-4-sulfonic acid 2-[5-methyl-2-(6-morpholin-4-yl-pyridin-3-yl)-thiazol-4-yl]-
ethyl ester



20 ¹HNMR 400 MHz (CDCl₃) δ 8.51 (s, 1H), 7.82 (d, 1H, J=8.5 Hz), 7.63 (d, 2H, J=7.8 Hz), 7.19 (d, 2H, J=7.8 Hz), 6.61 (d, 1H, J=8.5 Hz), 4.37 (t, 2H, J=6.5 Hz), 3.82 (t, 2H, J=5.2 Hz), 3.58 (t, 2H, J=5.2 Hz), 2.99 (t, 2H, J=6.5 Hz), 2.35 (s, 3H), 2.27 (s, 3H).


-112-

- 5 4-{5-Methyl-4-[2-(toluene-4-sulfonyloxy)-ethyl]-thiazol-2-yl}-piperazine-1-carboxylic acid tert-butyl ester




MS (ESI) m/z 482 (M+H)⁺.

- 10 Toluene-4-sulfonic acid 2-[5-methyl-2-(4-methyl-piperazin-1-yl)-thiazol-4-yl]-ethyl ester



MS (ESI) m/z 396.1 (M+H)⁺.


- Toluene-4-sulfonic acid 2-[5-methyl-2-(4-phenyl-piperazin-1-yl)-thiazol-4-yl]-ethyl ester



15





MS (ESI) m/z 458.1 (M+H)⁺.

- 2-(5-Methyl-2-phenyl-thiazol-4-yl)-ethanol






- 20 MS (ESI) m/z 220 (M+H)⁺.

-113-

5 Toluene-4-sulfonic acid 2-(5-methyl-2-phenyl-thiazol-4-yl)-ethyl ester:MS (ESI) m/z 374 (M+H)⁺.10 2-(2-Biphenyl-4-yl-5-methyl-thiazol-4-yl)-ethanolMS (ESI) m/z 296 (M+H)⁺.15 Toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methyl-thiazol-4-yl)-ethyl esterMS (ESI) m/z 450 (M+H)⁺.20 Toluene-4-sulfonic acid 2-(5-methyl-2-pyridin-2-ylthiazol-4-yl) ethyl esterMS (ESI) m/z 375.1 (M+H)⁺.

-114-



5 Toluene-4-sulfonic acid 2-(5-methyl-2-pyridin-3-ylthiazol-4-yl) ethyl esterMS (ESI) m/z 375.1 (M+H)⁺.10 Toluene-4-sulfonic acid 2-(5-methyl-2-pyridin-4-ylthiazol-4-yl) ethyl esterMS (ESI) m/z 375 (M+H)⁺, MS (ESI) m/z 383.1 (M+H)⁺.15 Toluene-4-sulfonic acid 2-[2-(2-methoxyethylamino)-5-methylthiazol-4-yl] ethyl esterMS (ESI) m/z 371 (M+H)⁺.

-115-

Pyrazoles

Preparation 10


Toluene-4-sulfonic acid 2-(5-methyl-3-phenyl-pyrazol-1-yl)-ethyl ester

Step A: 5-Methyl-3-phenyl-1H-pyrazole

10

Hydrazine hydrate (9.0 mL, 99 mmol, 35 wt.% in H₂O; 0.64 equiv) was added to a solution of benzoylacetone (25.00 g, 154.1 mmol, 1 equiv) in ethanol (250 mL). After stirring 14 h, more hydrazine hydrate (8.0 mL, 88 mmol, 0.57 equiv) was added. After 2 h, the reaction solution was concentrated (95°C) to give the title compound as a white solid (24.31 g, 99.7%). HRMS Calculated for C₁₀H₁₁N₂: m/z 159.0922. Found: 159.0917

15

Step B: 2-(5-Methyl-3-phenyl-pyrazol-1-yl)-ethanol

20

Sodium hydride (2.5 g, 1.5 g NaH, 62 mmol, 1.1 equiv) was added over a period of 3 min to a solution of 5-methyl-3-phenyl-1H-pyrazole (9.00 g, 56.9 mmol, 1 equiv) in DMF (90 mL) cooled to 0 °C in an ice bath. After stirring 15 min, ethylene carbonate (7.6 mL, 10 g, 110 mmol, 2.0 equiv) was added. The bath was removed, and the reaction mixture was stirred for 15 h. The mixture was treated with 4 M aq K₂CO₃ (90 mL), heated at reflux for 5 h, and diluted with H₂O (200 mL). After allowing the hot mixture to cool for 15 min, more H₂O (100 mL) and then hexanes (100 mL) were added. The mixture was shaken vigorously and then allowed to separate. Crystals formed and stayed with the top organic layer. The aqueous layer was separated, and the crystals were collected by vacuum filtration and washed with hexanes (2 x 50 mL). The crystals were dissolved in Et₂O/EtOAc (1:1; 200 mL), and the solution was dried (Na₂SO₄), filtered,

25

30

-116-


- 5 and concentrated (75 °C) to give the title compound as an off-white crystalline solid (6.86 g, 59.6%). HRMS Calculated for $C_{12}H_{15}N_2O$: m/z 203.1184. Found: 203.1168

Step C: Toluene-4-sulfonic acid 2-(5-methyl-3-phenyl-pyrazol-1-yl)-ethyl ester


- 10 According to Preparation 9, Step E, 2-(5-methyl-3-phenyl-pyrazol-1-yl)-ethanol was converted into the title compound. MS (ESI) m/z 357 (M+H)⁺.

Preparation 11

2-(3-Biphenyl-4-yl-5-methyl-pyrazol-1-yl)-ethanol



- 15 Step A: 5-Biphenyl-4-yl-3-methyl-1H-pyrazole



- To a stirred mixture of NaH (1.98 g, .049 mol, 60% oil dispersion) in dry THF (30 mL) was added a suspension of diethoxyphosphorylacetone tosyl hydrazone (8.97 g, .024 mol; N Almirante *Syn. Lett.* 1999, 302.) in a mixture of THF (35 mL) and DMF (5.0 mL) dropwise over 15 min. The yellow suspension was stirred at 0-5°C for 30 min and was treated with a 4-biphenyl carboxaldehyde (3.10 g, .0169 mol) in dry THF (30 mL) at 0-5°C over 15 min. The orange solution was heated and stirred at reflux for 4 h and stirred at ambient temperature overnight. The mixture was poured into 5% aq. NaH_2PO_4 (350 mL) and extracted with EtOAc (2 x 200mL). The organic layers were combined, washed with brine, dried ($MgSO_4$), filtered, and concentrated to a yellow semi-solid. This material was triturated with hot EtOAc (20 mL) and filtered. The solid was washed with EtOAc (2 x 10 mL) and dried under high vacuum to give the title compound (2.61 g, 47%): HRMS Calculated for $C_{16}H_{15}N_2$: m/z 235.1235. Found: 235.1230.


- 30 Step B: 2-(3-Biphenyl-4-yl-5-methyl-pyrazol-1-yl)-ethanol

The title compound was prepared from 5-biphenyl-4-yl-3-methyl-1H-pyrazole according to the Preparation 10, Step B. HRMS Calculated for $C_{18}H_{19}N_2O$: m/z 279.1497. Found: 279.1496.

-117-

5 The following intermediate compounds are prepared by a substantially similar manner as described in Preparations 10 and 11.


2-[3-(4-Bromo-phenyl)-5-methyl-pyrazol-1-yl]-ethanol



10


HRMS Calculated for $C_{12}H_{14}BrN_2O$: m/z 281.0289. Found: 281.0288.

3-Methyl-5-naphthalen-2-yl-1H-pyrazole



15 HRMS Calculated for $C_{14}H_{12}N_2$: m/z 208.1001. Found: 208.0981.


2-(5-Methyl-3-naphthalen-2-yl-pyrazol-1-yl)-ethanol



HRMS Calculated for $C_{16}H_{17}N_2O$: m/z 253.1341. Found: 253.1339.

20


3-Methyl-5-naphthalen-1-yl-1H-pyrazole



Anal Calculated for $C_{14}H_{12}N_2$: C, 80.74; H, 5.81; N, 13.45. Found: C, 80.93; H, 5.70; N, 13.42; mp 115-117°C.

25


-118-

5 2-(5-Methyl-3-naphthalen-1-yl-pyrazol-1-yl)-ethanol.

Arylether bromide Preparations

Preparation 12


10 2-(2-Bromo-ethoxy)-6-methoxynaphthalene



To a solution of 6-methoxynaphthalen-2-ol (1.07 g, 6.14 mmol) in DMF (4 mL) were added cesium carbonate (3.11 g, 9.55 mmol) and dibromoethane (2.5 mL, 29 mmol). The mixture was stirred and heated at 55°C for 48 h. The reaction mixture was cooled, filtered, diluted with EtOAc, and washed with brine (2 x 30 mL). The organic layer was dried (Na₂SO₄) and concentrated. The crude product was purified using radial chromatography (2:98 to 25:75 EtOAc:Hex) to give the title compound as a white solid (0.52 g, 30 %): ¹H NMR (400 MHz, CDCl₃) δ 3.61 (t, J= 6.1 Hz, 2H), 3.82 (s, 3H), 4.30 (t, J= 6.4 Hz, 2H), 7.01-7.08 (m, 4H), 7.56 (dd, J= 12.0, 9.0 Hz, 2H).

20

The following intermediate compounds are prepared by a substantially similar manner as described in Preparation 12.


6-(2-Bromoethoxy)-3-phenylbenzofuran

25

The above compound is prepared from 3-phenylbenzofuran-6-ol (see *Bull. Soc. Chim. Fr.*, 942 (1962)). ¹H NMR (400 MHz, CDCl₃) δ 3.60 (t, J= 6.4 Hz, 2H), 4.28 (t, J= 6.4

- 5 Hz, 2H), 6.88 (dd, $J = 8.8, 2.4$ Hz, 1H), 7.00 (d, $J = 2.4$ Hz, 1H), 7.26-7.30 (m, 1H), 7.36-7.44 (m, 2H), 7.52-7.56 (m, 2H), 7.63 (d, $J = 9.8$ Hz, 2H).

4-(2-Bromoethoxy)-1-phenoxybenzene



- 10 ^1H NMR (400 MHz, CDCl_3) δ 3.55 (t, $J = 6.4$ Hz, 2H), 4.19 (t, $J = 6.1$ Hz, 2H), 6.80-6.90 (m, 6H), 6.96 (t, $J = 7.3$ Hz, 1H), 7.17-7.24 (m, 2H).

4-(3-Bromoethoxy)biphenyl: MS (ESI) m/z 295 ($\text{M} + \text{NH}_3$) $^+$.

- 3-(2-Bromoethoxy)biphenyl: ^1H NMR (400 MHz, CDCl_3) δ 3.58 (t, $J = 6.4$ Hz, 2H), 4.27 (t, $J = 6.4$ Hz, 2H), 6.81 (dd, $J = 8.3, 2.4$ Hz, 1H), 6.90 (d, $J = 8.8$ Hz, 1H), 7.13 (dd, $J = 7.8, 1.0$ Hz, 1H), 7.26-7.33 (m, 2H), 7.34-7.37 (m, 2H), 7.43-7.50 (m, 2H).


6-(4-Bromopropoxy)-3-phenylbenzofuran: ^1H NMR (400 MHz, CDCl_3) δ 2.26-2.33 (m, 2H), 3.57 (t, $J = 6.4$ Hz, 2H), 4.11 (t, $J = 5.9$ Hz, 2H), 6.89 (dd, $J = 8.6, 2.2$ Hz, 1H), 7.01 (d, $J = 2.0$ Hz, 1H), 7.29 (t, $J = 7.6$ Hz, 1H), 7.40 (t, $J = 7.6$ Hz, 2H), 7.56 (d, $J = 6.8$ Hz, 2H), 7.57-7.65 (m, 2H).

- 20 2-(4-Bromopropoxy)-6-methoxynaphthalene: ^1H NMR (400 MHz, CDCl_3) δ 2.29 (t, $J = 6.1$ Hz, 2H), 3.56 (t, $J = 6.4$ Hz, 2H), 3.81 (4.11 (t, $J = 5.9$ Hz, 2H), 7.01-7.14 (m, 4H), 7.52-7.57 (m, 2H).

- 3-(4-Bromopropoxy)biphenyl: ^1H NMR (400 MHz, CDCl_3) δ 2.27 (t, $J = 6.1$ Hz, 2H), 3.55 (t, $J = 6.6$ Hz, 2H), 4.09 (t, $J = 5.9$ Hz, 2H), 6.81 (dd, $J = 2.9, 1.0$ Hz, 1H), 7.05 (t, $J = 2.0$ Hz, 1H), 7.07 (t, $J = 2.0$ Hz, 1H), 7.22-7.24 (m, 2H), 7.26-7.37 (m, 2H), 7.43-7.52 (m, 3H).


-120-

5


4-(3-Bromopropoxy)-biphenyl:

$^1\text{H-NMR}$ (200.15 MHz, CDCl_3): δ 7.57-7.29 (m, 7H), 6.98 (dd, 2H, $J=6.72, 1.88$), 4.15 (t, 2H, $J=5.92$), 3.62 (t, 2H, $J=6.44$), 2.34 (qn, 2H, $J=5.92$).

10 4-(3-Bromopropoxy)-1-phenoxybenzene: $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.3(2H, m), 7.1(2H, m), 7.0(2H, m), 6.9(2H, m), 4.1(2H, m); 3.6(2H, m); 2.3(2H, m).

2-(4-Bromobutoxy)-6-methoxynaphthalene


15 $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 1.88-1.97 (m, 2H), 1.99-2.10 (m, 2H), 3.41-3.48 (m, 2H), 3.81 (s, 3H), 4.00 (t, $J= 5.9$ Hz, 2H), 7.00-7.05 (m, 3H), 7.13-7.19 (m, 1H), 7.54 (t, $J= 8.1$ Hz, 2H).

6-(4-Bromobutoxy)-3-phenylbenzofuran

20 $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 1.90-1.98 (m, 2H), 2.01-2.04 (m, 2H), 3.41-3.44 (m, 2H), 3.94-3.99 (m, 2H), 6.83-6.91 (m, 1H), 6.96-6.97 (m, 1H), 7.27-7.29 (m, 1H), 7.36-7.43 (m, 2H), 7.53-7.62 (m, 4H).


-121-

5

4-(4-Bromobutoxy)biphenyl


^1H NMR (400 MHz, CDCl_3) δ 2.03 (t, J = 6.8 Hz, 2H), 3.42 (t, J = 6.6 Hz, 2H), 3.96 (t, J = 6.5 Hz, 2H), 6.87 (d, J = 7.8 Hz, 2H), 7.17-7.23 (m, 3H), 7.32 (t, J = 7.6 Hz, 1H), 7.42-7.47 (m, 4H).

10

3-(4-Bromobutoxy)biphenyl

^1H NMR (400 MHz, CDCl_3) δ 1.86-1.95 (m, 2H), 1.96-2.05 (m, 2H), 2.42 (t, J = 6.6 Hz, 2H), 3.98 (t, J = 6.1 Hz, 2H), 6.79 (dd, J = 7.3, 2.0 Hz, 1H), 7.03-7.11 (m, 1H), 7.26 (t, J = 7.8 Hz, 2H), 7.35 (t, J = 7.8 Hz, 2H), 7.42-7.51 (m, 2H).

15

4-(4-Bromobutoxy)-1-phenoxybenzene

^1H NMR (400 MHz, CDCl_3) δ 1.83-2.03 (m, 2H), 1.96-2.03 (m, 2H), 3.41 (t, J = 6.6 Hz, 2H), 3.90 (t, J = 6.5 Hz, 2H), 6.76-6.78 (m, 2H), 6.79-6.90 (m, 4H), 6.94-6.97 (m, 1H), 7.19-7.23 (m, 2H).

20



-122-

5

Headpiece Preparations

Preparation 13


3-(2-Aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester

Step A: 2-Bromo-5-hydroxybenzaldehyde

10

To a suspension of 3-hydroxybenzaldehyde (1 kg, 8.2 mol) in CH_2Cl_2 (10 L) was added a solution of bromine (1.3 kg, 8.2 mol) in CH_2Cl_2 (620 mL) over 1 h. After stirring for about 22 h, the solids were isolated by filtration, washed with CH_2Cl_2 (4 L), and dried under vacuum to afford of 2-bromo-5-hydroxybenzaldehyde as a tan solid (1.05 kg, 63%).

15

Step B: 3-(2-Formyl-4-hydroxy-phenyl)-acrylic acid tert-butyl ester

To a solution of 2-bromo-5-hydroxybenzaldehyde (200 g, 1.0 mole) in propionitrile (6 L) under N_2 was added tri-*o*-tolyl phosphine (75.4 g, 0.25 mol) and diisopropylethylamine (350 mL, 260 g, 2.0 mol). The solution was degassed and purged with N_2 three times. To this solution was added tert-butylacrylate (440 mL, 385 g, 3.0 mol) and palladium acetate trimer (27.8 g, 0.12 mol). The mixture was degassed three times and heated to 80°C over 30 min. After 30 min, the reaction was cooled to ambient temperature and filtered. The solids were washed with acetonitrile. The filtrate was transferred to a separatory funnel with acetonitrile (4 L total) and was extracted with hexanes (4 x 5 L). The propionitrile/acetonitrile layer was concentrated by rotary evaporation. The residue was dissolved in toluene (4 L) and washed successively with 1N HCl (1 L), water (1 L), and brine (1 L). The organic layer was dried (Na_2SO_4), filtered, and concentrated to afford crude product as a wet brown solid (293 g). The crude

20

25


30

-123-

- 5 product was dissolved in CH_2Cl_2 (600 mL) with stirring. To this solution was added hexanes (1.8 L) over ~2 h, and the resulting slurry was stirred at ambient temperature for 1 h. The slurry was cooled in an ice water bath and stirred for 1 h. The product was isolated by filtration, washed with hexanes, and dried under vacuum to afford the product as a tan solid (199.5 g, 80.6%).

10

Step C: 3-[4-Hydroxy-2-(hydroxyimino-methyl)-phenyl]-acrylic acid tert-butyl ester



- A slurry of hydroxylamine hydrochloride (8.33 g, 0.12 mol), pyridine (9.49 g, 0.12 mol), and ethanol (100 mL) was stirred at ambient temperature for 30 min.
- 15 3-(2-Formyl-4-hydroxy-phenyl)-acrylic acid tert-butyl ester (14.90 g, 0.06 mol) was added followed by an ethanol (49 mL) rinse. The resulting brown solution was stirred at ambient temperature for 16 h and concentrated. The residue was suspended in water (150 mL) and extracted with ethyl acetate (3 x 100 mL). The combined organic layers were washed with saturated NaCl solution (2 x 100 mL), dried (Na_2SO_4), and filtered through a
- 20 plug of silica gel to remove trace polar impurities. The filtrate was concentrated to give crude product (16.04 g). A 13.56 g sample of the crude material was crystallized from heptane (70 mL) and ethyl acetate (80 mL) to the title compound (7.41 g, 55%), mp 146-149°C. ^1H NMR (CD_3OD): δ 1.53 (s, 9H), 4.83 (s, 2H), 6.15-6.23 (d, 1H), 6.82-6.87 (dd, 1H), 7.10-7.12 (d, 1H), 7.52-7.55 (d, 1H), 7.98-8.03 (d, 1H), 8.37 (s, 1H). MS (ES^+) m/z 286.1 ($[\text{M}+\text{Na}]^+$). MS (ES^-) m/z 262.2 ($[\text{M}-\text{H}]^-$). Anal. Calculated for $\text{C}_{14}\text{H}_{17}\text{NO}_4$: C: 63.8658; H: 6.5081; N: 5.3198. Found: C: 63.71; H: 6.38; N: 5.51.
- 25

Step D: 3-(2-Aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester


- A solution of 3-[4-hydroxy-2-(hydroxyimino-methyl)-phenyl]-acrylic acid
- 30 tert-butyl ester (0.50 g, 1.9 mmol) in ethanol (50 mL) was treated with 10% palladium on carbon (0.25 g). The resulting suspension was treated with hydrogen using a Parr shaker apparatus at ambient temperature and 50 psi for 16 h. The reaction mixture was filtered through Celite, rinsed with ethanol (20 mL), and concentrated. The residue was dissolved in ethyl acetate (50 mL) and washed with 10% K_2CO_3 solution (2 x 25 mL) and 1N

-124-

- 5 NaOH (2 x 25 mL). The organic layer was dried (Na_2SO_4), filtered, and concentrated. The residue was crystallized from isopropyl acetate to give the title compound (0.14 g, 40%), mp 154-157°C. ^1H NMR (CD_3OD): δ 1.37 (s, 9H), 2.44-2.50 (t, 2H), 2.80-2.85 (t, 2H), 3.75 (s, 2H), 4.83 (s, 3H), 6.57-6.63 (dd, 1H), 6.75-6.77 (d, 1H), 6.95-7.00 (d, 1H). MS (ES^+) m/z 252.1 ($[\text{M}+\text{H}]^+$). MS (ES^-) m/z 250.2 ($[\text{M}-\text{H}]^-$). Anal. Calculated for $\text{C}_{14}\text{H}_{21}\text{NO}_3$: C: 66.9069; H: 8.4222; N: 5.5731. Found: C: 66.57; H: 8.29; N: 5.60.

Preparation 14

3-(2-Aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester Ethandioate
(2: 1)



15


- Alternative to the crystallization of 3-(2-Aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester from isopropyl acetate. The crude tert-butyl-3-[2-(aminomethyl)-4-hydroxyphenyl]propanoate (0.50 g, 0.002 mol) was dissolved in refluxing ethyl acetate (10 mL). A solution of oxalic acid (0.18 g, 0.002 mol) dissolved in ethyl acetate (5 mL) was added and produced a precipitate immediately. The resulting slurry was cooled to 0°C and filtered. The isolated product was slurred in ethanol (5 mL), cooled to 0°C, and filtered to give 3-(2-aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester ethandioate (2:1) (0.41 g, 69%). mp 188-190°C. ^1H NMR ($\text{DMSO}-d_6$): δ 1.36 (s, 18H), 2.39-2.45 (t, 4H), 2.47-2.52 (m, 4H), 2.70-2.76 (t, 4H), 3.83 (s, 4H), 6.61-6.65 (dd, 2H), 6.77-6.82 (d, 2H), 6.97-7.01 (d, 2H). MS (ES^+) m/z 252.1 ($[\text{M}+\text{H}]^+$). MS (ES^-) m/z 250.2 ($[\text{M}-\text{H}]^-$). Anal. Calculated for $\text{C}_{30}\text{H}_{44}\text{N}_2\text{O}_{10}$: C: 60.80; H: 7.48; N: 4.73. Found: C: 60.79; H: 7.51; N: 4.81.

25

5

Preparation 15

3-[2-(tert-Butoxycarbonylamino-methyl)-4-hydroxy-phenyl]-propionic acid methyl ester




10 A solution of 3-(2-aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester (1.00 g, 3.98 mmol) in CH_2Cl_2 (10 mL) was treated with water (12 drops) then TFA (10 mL). The solution was stirred at ambient temperature for 6.5 h and concentrated. The residue was diluted with 2,2-dimethoxypropane (40 mL), conc. HCl (4 mL), and then MeOH (10 mL). The solution was stirred for 18 h and concentrated to 3-(2-aminomethyl-4-hydroxy-phenyl)-propionic acid: HCl salt as a tan solid. MS (ES^-) m/z 208 [M-1].

15 The solid was diluted with THF (20 mL) and saturated aq. NaHCO_3 solution (12 mL) and was treated with di-*tert*-butyl dicarbonate (1.09 g, 5.00 mmol). The mixture was stirred for 3 h and concentrated, and the residue was partitioned between EtOAc (75 mL) and 1N HCl (30 mL). The organic layer was dried (Na_2SO_4), and concentrated to give the title compound (1.16 g, 94 %). MS (ES^-) m/z 308 [M-1].

20

Preparation 16

3-[4-hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester



25


A slurry of 3-(2-aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester (75.4 g, 0.3 mol) in CH_2Cl_2 (900 mL) at 1°C was treated with triethylamine (60.7 g,

-126-

- 5 0.6 mol). Isopropyl chloroformate (300 mL, 0.3 mol, 1M in toluene) was added while maintaining the temperature less than 12°C. The resulting solution was stirred 16 h at ambient temperature. After 16 h, additional isopropyl chloroformate (15 mL, 0.015 mol, 1.0M in toluene) was added, and the reaction was stirred for 1 h. The reaction mixture was washed with 1N HCl (2 x 200 mL) and saturated NaHCO₃ solution (2 x 200 mL).
- 10 The organic layer was dried (Na₂SO₄) and concentrated. The residue was purified by filtration through Merck silica gel 62 (750 grams, CH₂Cl₂/MeOH 100/0 to 96/4) to give the title compound (95.48 g, 94.3%). ¹H NMR (300 MHz, CDCl₃): δ 1.20-1.24 (d, 6H), 1.40 (s, 9H), 2.46-2.52 (t, 2H), 2.81-2.86 (t, 2H), 4.29-4.32 (d, 2H), 4.86-4.97 (m, 1H), 5.19-5.28 (m, 1H), 6.67-6.72 (dd, 2H), 6.76 (s, 1H), 6.97-7.00 (d, 1H). MS (ES⁺) m/z
- 15 336.1 [M-H]⁺.


The following intermediate compounds are prepared by a substantially similar manner as described in Preparation 16.

- 20 3-[4-Hydroxy-2-(isobutoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester



MS (ES⁺) m/z 350 [M-H]⁺.


- 25 3-[2-(Cyclopropylmethoxycarbonylamino-methyl)-4-hydroxy-phenyl]-propionic acid tert-butyl ester



HRMS Calculated for C₁₉H₂₆NO₅: m/z 348.1811. Found: 348.1817.


-127-

5 3-[2-(Cyclobutoxycarbonylamino-methyl)-4-hydroxy-phenyl]-propionic acid *tert*-butyl ester



HRMS Calculated for $C_{19}H_{26}NO_5$: m/z 348.1811. Found: 348.1817.


10 3-[2-(Cyclopentyloxycarbonylamino-methyl)-4-hydroxy-phenyl]-propionic acid *tert*-butyl ester



MS (ES^+) m/z 362 $[M-H]^+$.

15 Preparation 17

3-[4-Hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid methyl ester



A solution of 3-[4-hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid *tert*-butyl ester (3.0 mmol, 1.0g) in CH_2Cl_2 (10 mL) was treated with 90% CF_3CO_2H/H_2O (20 mL) at ambient temperature. The mixture was stirred for 3 h and concentrated. The residue was dissolved in MeOH (20 mL), treated with conc. H_2SO_4 (1


-128-

- 5 mL), and heated at reflux for 14 h. The mixture was cooled and concentrated. The residue was dissolved in water/ethyl acetate and neutralized with K_2CO_3 . The organic layer was dried (Na_2SO_4), and concentrated to give the title compound (0.76 g, 86 %). HRMS Calculated for $C_{15}H_{22}NO_5$: m/z 296.1498. Found: 296.1504.

10

Preparation 18

3-[4-Hydroxy-2-(3-isopropyl-ureidomethyl)-phenyl]-propionic acid tert-butyl ester




- A slurry of 3-(2-aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester (2.24 g, 8.91 mmol) in CH_2Cl_2 (30 mL) was treated with isopropyl isocyanate (1.1 mL, 11 mmol). The reaction mixture was diluted with water, the CH_2Cl_2 was removed under reduced pressure, and the aqueous layer was extracted with EtOAc. The organic layer was dried ($MgSO_4$) and concentrated to the title compound as a white solid (2.6 g, 87%). MS (ES^+) m/z 337 $[M+H]^+$.


-129-

5 **Preparation 19**

3-(2-{[(2,5-dichloro-thiophene-3-carbonyl)-amino]-methyl}-4-hydroxy-phenyl)-
propionic acid tert-butyl ester



Step A: 2,5-Dichloro-thiophene-3-carboxylic acid



10

A mixture of the 1-(2,5-dichloro-thiophen-3-yl)-ethanone (10 g, 51.26 mmol) and 9.5% NaOCl (150 mL, 230 mmol, 4.5 eq., commercial bleach) was treated with 5N NaOH (1 mL, 5 mmol, 0.1 eq.). The mixture was stirred vigorously and heated to 55 °C. The internal temperature was monitored closely and heat was removed to control the exotherm. After 6 h at 61°C, starting material was completely consumed. The mixture was cooled to 0°C and carefully quenched with 20 % aq. NaHSO₃ solution(20 mL). At 0°C, 6M HCl (12 mL) was added to adjust the pH to 1.5. The mixture was extracted with EtOAc (300 mL and 3 x 50 mL). The combined organic layers were washed with brine (200 mL), dried (Na₂SO₄), and concentrated to a white solid (8.8 g).

15

-130-


- 5 Step B: 3-(2-[[2,5-dichloro-thiophene-3-carbonyl]-amino]-methyl)-4-hydroxy-phenyl)-propionic acid tert-butyl ester

A solution of the 2,5-dichloro-thiophene-3-carboxylic acid (12.9g, 65.5 mmol) and 4-methylmorpholine (7.17 mL, 65.2 mmol) in dry THF (400 mL) was cooled to -15°C. Isobutyl chloroformate (8.46 mL, 65.2 mmol) was added. The mixture was stirred 3 min and triethylamine (9.1 mL, 65 mmol) was added. A solution of 3-(2-aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester (16.4 g, 65.3 mmol) in DMF (130 mL) pre-cooled to -15°C was added via cannula over 15 min. After stirring 1 h, TLC indicated complete reaction. The reaction mixture was allowed to warm to ambient temperature. Solids were removed by filtration and washed with THF (100 mL). The filtrate was diluted with Et₂O (500 mL) and washed with water (250 mL) then brine (150 mL). The organic layer was dried (Na₂SO₄) and concentrated. The crude brown oil was purified by silica gel chromatography (hexanes/EtOAc 2/1) and recrystallization (toluene) to afford the title compound as a white crystalline solid (22.3 g, 79.6%). MS (ES⁺) m/z 430.1 [M+H]⁺.

20

Preparation 20

3-(4-Hydroxy-2-[[pyrazine-2-carbonyl]-amino]-methyl)-phenyl)-propionic acid tert-butyl ester



- 25 To a solution of pyrazine-2-carboxylic acid (0.570 g, 4.60 mmol) in CH₂Cl₂ (50 mL) was added EDC (0.960 g, 5.01 mmol) and HOBT (0.620 g, 4.60 mmol) at room temperature. The mixture was stirred for 10 min, and 3-(2-aminomethyl-4-hydroxy-phenyl)-propionic acid tert-butyl ester (1.05 g, 4.18 mmol) was added. The reaction mixture was stirred for 18 h at room temperature, treated with water (50 mL), and extracted with CH₂Cl₂ (2 x 50 mL). The combined organic layers were dried (Na₂SO₄), concentrated, and purified by silica gel chromatography (EtOAc/hexanes, 1:1
- 30


-131-

5 to 1:0) to obtain the title compound (1.05 g, 70%). ^1H NMR (400 MHz, CDCl_3): δ 1.40 (s, 9H), 2.52 (t, 2 H, $J = 7.6$ Hz), 2.91 (t, 2H, $J = 7.6$ Hz), 4.66 (d, 2H, $J = 6.4$ Hz), 6.73 (dd, 1H, $J = 2.9, 7.8$ Hz), 6.83 (d, 1H, $J = 2.4$ Hz), 7.07 (d, 1H, $J = 8.3$ Hz), 8.17 (br s, 1H), 8.49 (dd, 1H, $J = 1.5, 2.4$ Hz), 8.73 (d, 1H, $J = 2.4$ Hz), 9.41 (d, 1H, $J = 1.5$ Hz). MS (ES) m/z 356.1 ($[\text{M}-\text{H}]^-$).

10

The following intermediate compounds are prepared by a substantially similar manner as described in Preparation 20.


15 3-(4-Hydroxy-2-{[(pyridine-2-carbonyl)-amino]-methyl}-phenyl)-propionic acid tert-butyl ester



^1H -NMR(CDCl_3): 1.40 (s, 9H), 2.51 (t, $J=7.6$ Hz, 2H), 2.91 (t, $J=7.6$ Hz, 2H), 4.64 (d, $J=5.9$ Hz, 2H), 6.72 (d, $J=7.8$ Hz, 1H), 6.85 (s, 1H), 7.06 (d, $J=8.2$ Hz, 1H), 7.41 (t, $J=7.6$ Hz, 1H), 7.84 (t, $J=7.0$ Hz, 1H), 8.20 (d, $J=7.4$ Hz, 1H), 8.38 (br s, 1H), 8.50 (d, $J=3.5$ Hz, 1 H); MS (ES) m/z 348 ($\text{M}+\text{H}$).

-132-

5

Preparation 213-[2-(benzoylamino-methyl)-4-hydroxy-phenyl]propionic acid *tert*-butyl ester


A slurry of 3-(2-aminomethyl-4-hydroxy-phenyl)-propionic acid *tert*-butyl ester (0.36 g, 1.43 mmol) in CH_2Cl_2 (10 mL) was cooled in an ice-water bath and treated with triethyl amine (0.40 mL, 0.28 mmol) then dropwise with benzoyl chloride (0.18 mL, 1.55 mmol). The resulting solution was stirred 30 min, and the cooling bath was removed. After 30 min, the solution was concentrated, and the residue was partitioned between EtOAc (50 mL) and 1N HCl (10 mL). The organic layer was washed with brine (10 mL), dried (Na_2SO_4), and concentrated to a solid. The crude mixture was purified by radial chromatography (hexanes:EtOAc 3:1 to 2:1) to give the title compound as a white solid (325 mg, 64%): ^1H NMR (400 MHz, CDCl_3) δ 1.26 (s, 9H), 2.54 (t, J = 7.1 Hz, 2H), 2.81 (t, J = 6.8 Hz, 2H), 4.55 (d, J = 2H), 6.68 (dd, J = 8.3, 2.4 Hz, 1H), 6.86 (d, J = 2.9 Hz, 1H), 6.96 (d, J = 8.3 Hz, 1H), 7.25-7.29 (m, 2H), 7.34-7.37 (m, 1H), 7.49 (br s, 1H), 7.16-7.79 (m, 2H); MS (ES) m/z 356.2 (M+H).

20

-133-

5 The following intermediate compound is prepared by a substantially similar manner as described in Preparation 21.

3-{2-[(Cyclobutanecarbonyl-amino)-methyl]-4-hydroxy-phenyl}-propionic acid tert-butyl ester




10

MS (ESI) m/z 334.2 ($M+H$)⁺.


Preparation 22

3-[2-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-hydroxy-phenyl]-propionic acid tert-butyl ester

15



Step A: (4-Bromo-3-methyl-phenoxy)-tert-butyl-dimethyl-silane




20 A 12 L flask was charged with 4-bromo-3-methyl phenol (428 g, 2.29 mol), CH₂Cl₂ (7.5 L), triethylamine (480 mL, 3.45 mol), and tert-butyldimethylsilyl chloride (324 g, 2.15 mol). To the solution was added 4-dimethylaminopyridine (15.0 g, 0.123 mol). The reaction mixture was stirred at ambient temperature overnight. The reaction was washed with saturated ammonium chloride (2.2 L) and then DI water (0.9

-134-

- 5 L). The organic layer was dried (Na_2SO_4), filtered, and concentrated to crude product (699 g). This material was purified by silica gel chromatography (heptane) to give the title compound (637 g, 98.5%).

Step B: (4-Bromo-3-bromomethyl-phenoxy)-tert-butyl-dimethyl-silane




10

- (4-Bromo-3-methyl-phenoxy)-tert-butyl-dimethyl-silane (255 g, 0.846 mol), dichloroethane (2.5 L), N-bromosuccinimide (165 g, 0.927 mol) and 2,2'-azobisisobutyronitrile (19.0 g, 0.116 mol) were combined in a 5 L flask. The mixture was degassed by evacuating and purging with N_2 (5x). The reaction mixture was heated to 47°C, and the heat was shut off. An exotherm to 76°C occurred. GC analysis showed 6.5% unreacted starting material. The heat was applied again, and the reaction was held at reflux (83°C) for 15 min. After cooling to 8°C, heptane (1.0 L) was added. The resulting slurry was stirred at 4 °C for 30 min and filtered. The filtrate was evaporated to dryness. The residue was treated with heptane (1 L), placed in the freezer overnight, and filtered. The filtrate was concentrated to the title compound (326 g, 101%).

20

Step C: 2-[2-Bromo-5-(tert-butyl-dimethyl-silanyloxy)-benzyl]-isoindole-1,3-dione




- A 12 L flask was charged with (4-bromo-3-bromomethyl-phenoxy)-tert-butyl-dimethyl-silane (568 g 1.49 mol), DMF (3.1 L), and potassium phthalimide (316 g 1.71 mol). An exotherm to 34 °C occurred. After 40 min, the reaction mixture was cooled to 18°C. Ether (6.2 L) and DI water (4.9 L) were added, and the layers were separated. The organic layer was washed with saturated NaCl solution (2 L), dried (Na_2SO_4), filtered, and concentrated. The residue was recrystallized from heptane (1.5 L) to give the title compound (454 g, 68%).

30

-135-

- 5 Step D: 3-[2-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-hydroxy-phenyl]-acrylic acid tert-butyl ester



- A 12 L flask was charged with 2-[2-bromo-5-(tert-butyl-dimethyl-silanyloxy)-benzyl]-isoindole-1,3-dione (461 g, 1.03 mol), propionitrile (7 L), tri-*ortho*-tolyl phosphine (76.0 g, 0.250 mol) and diisopropyl ethyl amine (365 mL, 2.10 mol). The reaction mixture was degassed/purged with N₂ (3x), and tert-butyl acrylate (465 mL, 3.17 mol) was added. After degassing/purging one time, palladium (II) acetate (28.0 g, 0.125 mol) was added. The stirred mixture was degassed/purged with N₂ three times and heated to 95°C for 20 h. The mixture was filtered through a hyflo cake, washed with
- 10 acetonitrile, and concentrated to a brown oil (841 g). The residue was dissolved in THF (3.5 L), and tetrabutylammonium fluoride (TBAF, 650 mL, 0.65 mol, 1M in THF) was added. After 1 h, additional TBAF (95 mL) was added. The mixture was rotated on the rotary evaporator for 10 min and was concentrated to crude product (987 g). This material was purified by silica gel chromatography (toluene/ethyl acetate, 100/0 to 75/25)
- 15 to give the title compound (340 g, 86.8%).
- 20

Step E: 3-[2-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-hydroxy-phenyl]-propionic acid tert-butyl ester


- A 1 gallon autoclave was charged with 3-[2-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-hydroxy-phenyl]-acrylic acid tert-butyl ester (196 g, 0.517 mol), ethyl acetate (2.6 L) and 5 % palladium on carbon (75 g). The autoclave was kept at 25°C under 60 psi of hydrogen for 21 h. The temperature of the reaction was increased to 40 °C, and the pressure was increased to 75 psi for 5 h. The mixture was filtered through a pad of hyflo and concentrated to the title compound (186 g, 94.4 %): MS (ESI) m/z
- 25
- 30 380.2 (M-H)⁻.

-136-

5


Preparation 23

3-{4-Hydroxy-2-[(isopropoxycarbonyl-methyl-amino)-methyl]-phenyl}-propionic acid
tert-butyl ester



10

Step A: 3-[5-Benzyloxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid
tert-butyl ester



15

To a solution of 3-[5-hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (14.8 mmol) in DMF (100 mL) was added Cs_2CO_3 (29.6 mmol) and benzyl bromide (16.3 mmol). The reaction mixture was stirred overnight at ambient temperature and diluted with EtOAc (200 mL). The mixture was washed with brine (100 mL) and water (100 mL). The organic layer was dried (Na_2SO_4) and concentrated to a yellow oil (6.8 g). MS [ES] m/z 428 ($\text{M}+\text{H}$)⁺.

20

Step B: 3-{5-Hydroxy-2-[(isopropoxycarbonyl-methyl-amino)-methyl]-phenyl}-propionic acid tert-butyl ester

25

To a solution of 3-[5-benzyloxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (6.3 g, 0.015 mol) in DMF (100 mL) was added NaH (29.4 mmol, 60% oil dispersion). The reaction mixture was stirred at 0°C for 15 min, treated with methyl iodide (4.18 g, 29.4 mmol), and stirred overnight under N_2 at ambient temperature. The mixture was concentrated and triturated with hexanes (200 mL). The residue was diluted with EtOAc (200 mL), and the solution was washed with

-137-

5 water (200 mL) and brine (200 mL), dried (Na_2SO_4), and concentrated to a yellow oil (5.2 g). The material was dissolved in EtOAc (100 mL), and 5% Pd/C (0.725 g) was added. The reaction mixture was shaken under a hydrogen atmosphere in a Parr apparatus at 60psi overnight at 40°C. The mixture was filtered and concentrated to give the title compound (2.1 g). MS [ES] m/z 352 ($\text{M}+\text{H}$)⁺.


10

The following intermediate compound is prepared by a substantially similar manner as described in Preparation 23.

3-(5-Hydroxy-2-{{methyl-(pyridine-2-carbonyl)-amino}-methyl}-phenyl)-propionic acid

15

tert-butyl ester




MS [ES] m/z 371 ($\text{M}+\text{H}$)⁺.

20 The following General Procedures are used to prepare the compounds of present invention as illustrated below.

General Procedures I and II

General Procedure I

General Procedure for the Parallel Synthesis of Carboxamides from Carboxylic Acids



25

3-{2-Aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester: acetic acid salt (704 mg, 1.4 mmol) was


-138-

5 dissolved in CH_2Cl_2 (25 mL) and washed with saturated NaHCO_3 solution (15 mL). The organic layer was dried (NaSO_4), filtered, and concentrated to 3-{2-aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester (636 mg, 1.4 mmol). This amine (0.1 mmol, 0.1 M in CH_2Cl_2) was added to a carboxylic acid (1.4 eq) in a 1 dram vial. N-Hydroxybenzotriazole hydrate (1.4eq, 0.17 M in $\text{CH}_2\text{Cl}_2/\text{DMF}$ 10/1; HOBt) and EDC (1.4 eq, 0.18 M CH_2Cl_2) were added, and the vials were capped and shaken for 18 h. DMF (0.5 mL) and triethylamine (0.5 mL) were added and the vials were shaken for 72 h. Brine (1 mL) was added, and the mixtures were transferred to a ChemElute cartridge and eluted with CH_2Cl_2 . The solvent was removed under a stream of N_2 . The residue was treated with a mixture of $\text{CH}_2\text{Cl}_2/\text{TFA}/\text{water}$ 60/35/5 (1 mL), 15 agitated briefly, and allowed to stand at ambient temperature for 2 h. The solvent was removed under a stream of N_2 . The residue was treated with 10% $\text{CCl}_4/\text{CH}_2\text{Cl}_2$ (1 mL) and concentrated under a stream of N_2 (2x). The crude products were dried under vacuum and purified by mass-directed HPLC.

20

General Procedure II

General Procedure for the Parallel Synthesis of Carboxamides, Carbamates, Sulfonamides, Ureas, and Thioureas from Carboxylic Acid Chlorides, Chloroformates, Sulfonyl Chlorides, Isocyanates, and Isothiocyanates




25

3-{2-Aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester: acetic acid salt (720 mg, 1.4 mmol) was dissolved in CH_2Cl_2 (25 mL) and washed with saturated NaHCO_3 solution (15 mL). The organic layer was dried (NaSO_4), filtered, and concentrated to 3-{2-aminomethyl-4-[2-(5-


-139-

- 5 methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester (586 mg, 92%). This amine (0.1 mmol, 0.1 M in CH_2Cl_2) and carboxylic acid chloride, chloroformate, sulfonyl chloride, isocyanate, or thioisocyanate (1 eq), and triethylamine (0.2 mL) were placed in a 1 dram vial. The vials were capped and shaken for 18 h. DMF (0.5 mL) was added to incomplete reaction mixtures, and the vials were shaken for 2 h.
- 10 Brine (1 mL) was added, and the mixtures were transferred to a ChemElute cartridge and eluted with CH_2Cl_2 . The solvent was removed under a stream of N_2 . The residue was treated with a mixture of CH_2Cl_2 /TFA/water 60/35/5 (1 mL), agitated briefly, and allowed to stand at ambient temperature for 2 h. The solvent was removed under a stream of N_2 . The residue was treated with 10% $\text{CCl}_4/\text{CH}_2\text{Cl}_2$ (1 mL) and concentrated under a stream
- 15 of N_2 (2x). The crude products were dried under vacuum and purified by mass-directed HPLC.

Example 1

20


Step A: 3-{2-(1,3-Dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester



5 A 2 L 3-neck flask was charged with 3-[2-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-4-hydroxy-phenyl]-propionic acid tert-butyl ester (67.9 g, 0.178mol),
toluene-4-sulfonic acid 2-(5-methyl-2-phenyl-oxazol-4-yl)-ethyl ester (76.6 g, 0.214 mol)
and DMF (680 mL). Cesium carbonate (75.4 g, 0.231 mol) was added, and the reaction
mixture was heated at 55 °C for 18 h. After cooling, ethyl acetate (890 mL) and DI water
10 (1200 mL) were added, the mixture was agitated, and the layers were separated. The
aqueous layer was back-extracted with ethyl acetate (740 mL). The organic layers were
combined and washed with 1N NaOH (375 mL) then saturated NaCl solution (2 x 375
mL). The organic solution was dried (Na₂SO₄), filtered, and concentrated to an oil (107
g). The crude oil was dissolved in toluene (50 mL), and heptane was added (~100 mL)
15 until cloudiness remained even with agitation. The mixture was warmed to 50 °C on a
rotary evaporator to yield a solution. Seed crystals were added, and rotation was
continued at ambient temperature overnight. The product slurry was placed in the freezer
overnight and filtered. The title compound was dried in a vacuum oven at 35 °C (71.4 g,
70.8 %).

20

Step B: 3-{2-Aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester: acetic acid salt






A 3 L 3-neck flask was charged with 3-{2-(1,3-dioxo-1,3-dihydro-
25 isoindol-2-ylmethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic
acid tert-butyl ester (71.0 g, 0.125 mol) and isopropanol (IPA, 1.35 L). After warming to
40 °C and adding IPA (250 mL), a solution was still not obtained. After cooling to <30
°C, sodium borohydride (25.6 g, 0.677 mol) was added carefully in portions. Tap water
was added to the bath, and the reaction was allowed to stir overnight. A spatula was used
30 to break up the solids in the thick slurry, and IPA (200 mL) was added. Glacial acetic
acid (130 mL, 2.27 mol) was added dropwise over 2 h. The reaction mixture was heated
at reflux for 10 h, allowed to cool over the weekend, and concentrated to crude product
(237 g). CH₂Cl₂ (100 mL) was added to give a thick gel which was poured onto a column







-141-







- 5 of silica gel 60 equilibrated with CH_2Cl_2 . Methanol was used to help dissolve the crude material. Eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (1/1) gave the product mixture (182 g). Ethyl acetate (200 mL) was added, and a precipitate formed. The mixture was heated to 40 °C, and the solution was cooled to 0 °C. The pure product was filtered, washed with heptane/EtOAc (1:1, 2 x 100 mL), and dried to yield a white fluffy solid (35.4 g, 57%).
- 10 The filtrate was concentrated (116 g) and purified again by silica gel chromatography ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 100/0 to 80/20) to give additional product as an amber oil (13.9 g, 22%).

Examples 2- 230







- Examples 2-230 are prepared by following a substantially similar
- 15 procedure as described in Example 1 and General Procedures I and II.

No	Compounds	MS (ES+)	General Procedure
2		515.5 FIA	II
3		513.5 FIA	I
4		514.4 FIA	II

No	Compounds	MS (ES+)	General Procedur e
5		535.2 FIA	II
6		501.3 FIA	II
7		557.1 FIA	II
8		505.3	II
9		491.3	I
10		493.3	I







No	Compounds	MS (ES+)	General Procedur e
11		451.2	I
12		451.2	II
13		463.3	II
14		505.3	II
15		521.6	II
16		531.6	II







No	Compounds	MS (ES+)	General Procedur e
17		575.7	II
18		621.6	II
19		559.8	II
20		543.7	II
21		555.4	II
22		525.2	II







No	Compounds	MS (ES+)	General Procedur
23		529.3	II
24		515.3	II
25		510.6	II
26		543.7	II
27		539.3	II
28		569.4	II

No	Compounds	MS (ES+)	General Procedures
29		571.3	II
30		513.4	II
31		619.4	II
32		539.3	II
33		543.4	II
34		571.3	II







-147-

No	Compounds	MS (ES+)	General Procedur e
35		589.2	II
36		577.3	II
37		529.7	I
38		505.3	I
39		525.2	I
40		549.3	I







No	Compounds	MS (ES+)	General Procedur
41		491.3	I
42		521.3	I
43		489.3	I
44		485.3	I
45		505.3	I
46		505.3	I

No	Compounds	MS (ES+)	General Procedur e
47		505.3	I
48		491.3	I
49		571.2	I
50		597.4	I
51		505.3	I
52		499.3	I







-150-







No	Compounds	MS (ES+)	General Procedur e
53		533.3	I
54		547.3	I
55		547.3	I
56		575	I
57		575	I
58		591	I

-151-







No	Compounds	MS (ES+)	General Procedur e
59		521	I
60		591	I
61		477.2	I
62		465	I
63		491	I
64		519	I







-152-







No	Compounds	MS (ES+)	General Procedur e
65		541	I
66		493	I
67		541	I
68		554	I
69		507	I
70		527	I







No	Compounds	MS (ES+)	General Procedur
71		463	I
72		521	I
73		554	I
74		520	I
75		545	I
76		503	I







No	Compounds	MS (ES+)	General Procedures
77		554	I
78		503	I
79		553	I
80		545	I
81		499	I
82		515.2	I







No	Compounds	MS (ES+)	General Procedur e
83		545.3	I
84		554	I
85		515	I
86		554	I
87		554	I
88		598	I







No	Compounds	MS (ES+)	General Procedur e
89		499	I
90		555	I
91		529	I
92		465	I
93		533	I
94		465	I







No	Compounds	MS (ES+)	General Procedur
95		538	I
96		499	I
97		479	I
98		521	I
99		538	I
100		521	I

No	Compounds	MS (ES+)	General Procedur e
101		553	I
102		538	I
103		513	I
104		513	I
105		503	I
106		527	I







No	Compounds	MS (ES+)	General Procedur
107		513	I
108		513	I
109		513	I
110		521	I
111		513	I
112		535	I







No	Compounds	MS (ES+)	General Procedur
113		545	I
114		545	I
115		527	I
116		555	I
117		541	I
118		538	I







No	Compounds	MS (ES+)	General Procedur
119		564	I
120		577	I
121		569	I
122		564	I
123		575	I
124		587	I

No	Compounds	MS (ES+)	General Procedur e
125		587	I
126		575	I
127		491	I
128		564	I
129		538	I
130		560	I







No	Compounds	MS (ES+)	General Procedur e
131		575	I
132		621	I
133		527	I
134		569	I
135		571	I
136		571	I







No	Compounds	MS (ES+)	General Procedur e
137		597	I
138		569	I
139		621	I
140		571	I
141		594	I
142		476	I







No	Compounds	MS (ES+)	General Procedur e
143		617	I
144		500	I
145		469	I
146		561	I
147		509	I
148		521	I

No	Compounds	MS (ES+)	General Procedur e
149		577	I
150		589	I
151		561	I
152		595	I
153		575	I
154		599	I







-167-







No	Compounds	MS (ES+)	General Procedur
155		566	I
156		599	I
157		543	I
158		650	I
159		619	I
160		615	I

No	Compounds	MS (ES+)	General Procedur
161		437.2	I
162		571.2	I
163		see text	I
164		493.2	II
165		483.2	II
166		551.2	II







No	Compounds	MS (ES+)	General Procedur
167		467.2	II
168		481.2	II
169		501.2	II
170		439.2	II
171		481.3	II
172		453.2	II







-170-

No	Compounds	MS (ES+)	General Procedur e
173		565.4	II
174		535.2	II
175		475.2	I
176		536.2	I
177		524.2	I
178		487.2	I

No	Compounds	MS (ES+)	General Procedur e
179		479.2	I
180		486.2	I
181		486.2	I
182	 HCl	474.2	I
183	 HCl	486.2	I
184		512	I







No	Compounds	MS (ES+)	General Procedur e
185		512.2	I
186		568.1	I
187		556.1	I
188		537.1	I
189		574.1	I
190		559.1	I







No	Compounds	MS (ES+)	General Procedur e
191		487.3	II
192		473.0	II
193		487.3	II
194		521.3	II
195		527.2	II
196		466.3	II







No	Compounds	MS (ES+)	General Procedur e
197		528.3	II
198		534.3	II
199		480.3	II
200		535.2	II
201		611.1	II
202		539.2	II

No	Compounds	MS (ES+)	General Procedures
203		539.1	II
204		557.1	II
205		557.1	II
206		500.2	II
207		550.2	II
208		592.2	II





-176-

No	Compounds	MS (ES+)	General Procedur e
209		516.1	II
210		575.1	II
211		539.2	II
212		529	II
213		467.3	II
214		495.3	II

No	Compounds	MS (ES+)	General Procedures
215		761.2 (FIA)	II
216		661.1 (FIA)	II
217		459.4 (FIA)	II
218		682	II
219		423	I
220		569	I

No	Compounds	MS (ES+)	General Procedur e
221		552	II
222		480	II
223		751	II
224		829	II
225		589	II
226		697	II

-179-


No	Compounds	MS (ES+)	General Procedure
227		733	II
228		733	II
229		733	II
230		494.1	II

-180-


5

Example 231

3-(4-[2-(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-[[2,5-dichloro-thiophene-3-carbonyl)-amino]-methyl}-phenyl)-propionic acid



10 Step A: 3-(4-[2-(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-[[2,5-dichloro-thiophene-3-carbonyl)-amino]-methyl}-phenyl)-propionic acid tert-butyl ester





A mixture of 3-(2-[[2,5-dichloro-thiophene-3-carbonyl)-amino]-methyl}-4-hydroxy-phenyl)-propionic acid tert-butyl ester (280 mg, 0.651 mmol) and toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethyl ester (498 mg, 1.15 mmol),
 15 were reacted according to Standard procedure A to give, after radial chromatography (hexanes/ EtOAc 98:2 to 90:10), the title compound as a white solid (367 mg, 82%). ¹H NMR (400 MHz, CDCl₃): δ 1.36 (s, 9 H), 2.37 (s, 3 H), 2.55 (t, 2H, J = 7.3 Hz), 2.87 (t, 2H, J = 7.3 Hz), 2.96 (t, 2 H, J = 6.6 Hz), 4.22 (t, 2H, J = 6.8 Hz), 4.58 (d, 2H, J = 5.4 Hz), 6.80 (dd, 1H, J = 8.3 Hz, J = 2.4 Hz), 6.88 (d, 1H, J = 2.9 Hz), 6.97 (t, 1H, J = 5.9 Hz), 7.10 (d, 1 H, J = 8.8 Hz), 7.13 (s, 1 H), 7.36 (t, 1H, J = 7.3 Hz), 7.45 (t, 2H, J = 7.8 Hz), 7.63 (d, 2 H, J = 8.3 Hz), 7.65 (d, 2H, J = 6.3 Hz), 8.02 (d, 2H, J = 7.8 Hz). MS (ES) m/e 691.2 [M+1].
 20







- 5 Step B: 3-(4-[2-(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-[[2,5-dichloro-thiophene-3-carbonyl)-amino]-methyl}-phenyl)-propionic acid

A solution of the 3-(4-[2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-[[2,5-dichloro-thiophene-3-carbonyl)-amino]-methyl}-phenyl)-propionic acid tert-butyl ester (367 mg, 0.531 mmol) in CH₂Cl₂ (20 mL) was treated with anisole (2.0 mL) then TFA (6.0 mL). The mixture was stirred at room temperature for 2 h and concentrated. The residue was co-evaporated with CCl₄ three times, dried under vacuum, triturated with ether, and filtered to obtain the title compound as a white solid (301 mg, 73%). ¹H NMR (400 MHz, DMSO-d₆): δ 2.25 (s, 3 H), 2.38 (t, 2H, J = 8.3 Hz), 2.71 (t, 2H, J = 7.3 Hz), 2.82 (t, 2H, J = 6.3 Hz), 4.07 (t, 2H, J = 6.6 Hz), 4.28 (d, 2H, J = 4.9 Hz), 6.68 (d, 1H, J = 9.3 Hz), 6.77 (s, 1H), 7.00 (d, 1H, J = 8.3 Hz), 7.24 (s, 1H), 7.29 (t, 1 H, J = 5.9 Hz), 7.39 (t, 2H, J = 7.8 Hz), 7.63 (t, 2H, J = 7.8 Hz), 7.71 (d, 2 H, J = 8.3 Hz), 7.88 (d, 2H, J = 8.3 Hz), 8.69 (t, 2H, J = 5.9 Hz). MS (ES) m/e 635.1 [M+1].







Examples 232-318







20 Examples 232-318 are prepared by following a substantially similar procedure as described in Example 231 and Standard coupling procedure A and Standard hydrolysis procedures C to E.







No.	Compounds	MS (ES+)
232		481.3
233		500.3







No.	Compounds	MS (ES+)
234		514.3
235		495.3
236		475.1
237		494.3
238		542.2
239		523.3

-183-







No.	Compounds	MS (ES+)
240		480.3
241		499.2
242	 ClH	503 ClH
243	 CH	484 CH
244		524.4
245		586.3







No.	Compounds	MS (ES+)
246		567.3
247		505.3
248		559.2
249		483.2
250		487.3
251		578.3

No.	Compounds	MS (ES+)
252		502.2
253		503.3
254		587.2
255		587.2
256		545/547
257		545/547







No.	Compounds	MS (ES+)
258		535
259		560
260		579
261	 <p>ClH</p>	484
262		539
263	 <p>ClH</p>	484







No.	Compounds	MS (ES+)
264		569
265		560
266		576.1
267		558.4
268		557.2
269		542







No.	Compounds	MS (ES+)
270		481.3
271		543.2
272		561.3
273		561.3
274		491.3
275		491.2







No.	Compounds	MS (ES+)
276		539.3
278		539.3
279		469.3
280		469.3
281		488.1
282		473.3

No.	Compounds	MS (ES+)
283		473.2
284		562.2
285		562.3
286		492.2
287		492.2
288		511.2

No.	Compounds	MS (ES+)
289		635.2
290		565.2
291		565
292		511
293		517
294		517

No.	Compounds	MS (ES+)
295		536
296		531
297		572.2
298		541
299		483
300		502

No.	Compounds	MS (ES+)
301		551.2770 (HRMS)
302		504.2171 (HRMS)
303		550.2
304		554
305		557.2
306		490.2

No.	Compounds	MS (ES+)
307		555.2426 (HRMS)
308		571.2436 (HRMS)
309		585.2595 (HRMS)
310		551.2753 (HRMS)
311		504.2144 (HRMS)
312		563.2

-195-

No.	Compounds	MS (ES+)
313		579.2
314		572.2
315		495.2
316		512.2
317		506.2
318		¹ H NMR

-196-


5 ¹H NMR data for Example 318:

(400 MHz, CDCl₃): d 6.98 (1H, d, J=7.9 Hz), 6.71 (1H, s), 6.68 (1H, d, J=7.9 Hz), 4.97 (1H, m), 4.86 (2H, m), 4.31 (2H, bs), 4.08 (2H, t, J=6.5 Hz), 3.45 (4H, m), 3.27 (4H, m), 2.86 (4H, m), 2.55 (2H, bs), 2.18 (3H, s), 1.42 (9H, s), 1.16 (6H, d, J=7.2 Hz).

10


Example 319

{4-[2-(5-Methyl-2-(4-phenyl)phenyl-oxazol-4-yl)-ethoxy]-2-(isopropylcarbamate)-methyl}-propionic acid



Step A: {4-[2-(5-Methyl-2-(4-phenyl)phenyl-oxazol-4-yl)-ethoxy]-2-

15 (isopropylcarbamate)methyl}-propionic acid tert-butyl ester



3-[4-hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (258 mg, 0.763 mmol) and toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethyl ester (654 mg, 1.51 mmol) were reacted according to

20 Standard procedure A. Purification by radial chromatography (CH₂Cl₂/EtOAc 98/2 to 90/10) gave the title compound in quantitative yield. ¹H NMR (400 MHz, CDCl₃):

δ 1.14 (d, 6H, J = 5.9 Hz), 1.31 (s, 9 H), 2.30 (s, 3 H), 2.39 (t, 2H, J = 7.6 Hz), 2.89 (t, 2H, J = 6.6 Hz), 4.14 (t, 2H, J = 6.6 Hz), 4.25 (d, 2H, J = 5.4 Hz), 4.85 (heptet, 1H, J = 5.8 Hz), 6.67 (dd, 1H, J = 8.3 Hz, J = 2.4 Hz), 6.75 (d, 1H, J = 2.4 Hz), 6.99 (d, 1H, J = 8.3 Hz), 7.26-7.30 (m, 1 H), 7.37 (t, 2H, J = 7.6 Hz), 7.53-7.58 (m, 4 H), 7.95 (d, 2H, J = 8.3 Hz). MS (ES) m/e 599.4 [M+1].

25


-197-

- 5 Step B: {4-[2-(5-Methyl-2-(4-phenyl)phenyl-oxazol-4-yl)-ethoxy]-2-(isopropylcarbamate)methyl}-propionic acid


A solution of {4-[2-(5-methyl-2-(4-phenyl)phenyl-oxazol-4-yl)-ethoxy]-2-(isopropylcarbamate)methyl}-propionic acid (414 mg, 0.763 mmol) in CH₂Cl₂ (10 mL) was treated with anisole (1.0 mL) then trifluoroacetic acid (TFA, 3.0 mL). The solution
10 was stirred at room temperature for 2 h and concentrated. The residue was co-evaporated with CCl₄ twice, dried under vacuum, triturated with ether, and filtered to obtain the title compound as a white solid (287 mg, 69%). ¹H NMR (400 MHz, CDCl₃) δ 1.11 (d, 6H, J = 5.9 Hz), 2.28 (s, 3 H), 2.46 (t, 2H, J = 7.6 Hz), 2.79 (t, 2H, J = 7.6 Hz), 2.86 (t, 2H, J = 6.6 Hz), 4.11 (t, 2H, J = 6.8 Hz), 4.24 (d, 2H, J = 5.4 Hz), 4.82 (heptet, 1H, J = 6.1 Hz),
15 5.05 (bs, 1 H), 6.66 (dd, 1H, J = 2.4, 8.3 Hz), 6.73 (d, 1H, J = 2.4 Hz), 6.99 (d, 1H, J = 8.3 Hz), 7.24-7.28 (m, 1 H), 7.35 (t, 2H, J = 7.6 Hz), 7.51-7.56 (m, 4 H), 7.98 (d, 2H, J = 8.3 Hz). MS (ES) m/e 543.1 [M+1].

Example 320

- 20 3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-2-morpholin-4-yl-thiazol-4-yl)ethoxy]-phenyl}-propionic acid



Step A: 3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-2-morpholin-4-yl-thiazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester



5 3-[4-hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic
acid tert-butyl ester (272 mg, 0.806 mmol) and toluene-4-sulfonic acid 2-(5-methyl-2-
morpholin-4-yl-thiazol-4-yl)-ethyl ester (522 mg, 1.36 mmol) were reacted according to
Standard procedure A. Purification using radial chromatography (CH₂Cl₂/EtOAc 98:2 to
75:25) gave the title compound as a white solid (387 mg, 98%). ¹H NMR (400 MHz,
10 CDCl₃) δ 1.14 (d, 6H, J = 5.9 Hz), 1.32 (s, 9 H), 2.16 (s, 3 H), 2.39 (t, 2H, J = 7.6 Hz),
2.77 (t, 2H, J = 7.6 Hz), 2.84 (t, 2H, J = 7.1 Hz), 3.28 (t, 4H, J = 4.9 Hz), 3.70 (t, 4H, J =
4.9 Hz), 4.08 (t, 2H, J = 7.1 Hz), 4.25 (d, 2H, J = 5.4 Hz), 4.83 (heptet, 1H, J = 4.9 Hz),
6.66 (dd, 1H, J = 2.4, 8.3 Hz), 6.72 (d, 1H, J = 2.4 Hz), 6.98 (d, 1H, J = 8.3 Hz). MS (ES)
m/e 548.3 [M+1].

15

Step B: 3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-2-morpholin-4-yl-
thiazol-4-yl)ethoxy]-phenyl}propionic acid

A solution of 3-{2-(isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-2-
morpholin-4-yl-thiazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester (387 mg,
20 0.706 mmol) in CH₂Cl₂ (10 mL) was treated with anisole (1 mL) then TFA (3 mL). The
solution was stirred at room temperature for 2 h and concentrated. The residue was co-
evaporated with CCl₄ twice, dried under vacuum, triturated with ether, and filtered to
obtain the title compound as a white solid (236 mg, 60%), isolated as the trifluoroacetic
acid salt: ¹H NMR (400 MHz, CDCl₃) δ 1.12 (d, 6H, J = 6.4 Hz), 2.16 (s, 3H), 2.47 (t,
25 2H, J = 7.6 Hz), 2.79 (t, 2H, J = 7.6 Hz), 2.87 (t, 2H, J = 6.4 Hz), 3.35 (t, 4H, J = 4.9 Hz),
3.70 (t, 4H, J = 4.9 Hz), 4.07 (t, 2H, J = 6.6 Hz), 4.23 (d, 2H, J = 5.4 Hz), 4.82 (heptet,
1H, J = 6.1 Hz), 5.07 (bs, 1H), 6.63 (dd, 1H, J = 2.7, 8.6 Hz), 6.69 (d, 1H, J = 2.9 Hz),
6.98 (d, 1H, J = 8.3 Hz). MS (ES) m/e 492.1 [M+1].


30

-199-

5


Example 321

3-(2-(2-Isopropoxycarbonyl-ethyl)-4-{2-[5-methyl-2-(6-phenoxy-pyridin-3-yl)-oxazol-4-yl]ethoxy}-phenyl)-propionic acid



10

Step A: 3-(2-(2-Isopropoxycarbonyl-ethyl)-4-{2-[5-methyl-2-(6-chloro-pyridin-3-yl)-oxazol-4-yl]ethoxy}-phenyl)-propionic acid t-butyl ester



15

Toluene-4-sulfonic acid 2-[2-(6-chloro-pyridin-3-yl)-5-methyl-oxazol-4-yl]-ethyl ester tosylate (290 mg, 0.74 mmol) and 3-[4-hydroxy-2-(2-isopropoxycarbonyl-ethyl)-phenyl]-propionic acid t-butyl ester (274 mg, 0.81 mmol) were converted to the title compound (183 mg, 44 %) according to Standard procedure A. MS (ESI) m/z 559 ($M+H$)⁺.

20

Step B: 3-(2-(2-Isopropoxycarbonyl-ethyl)-4-{2-[5-methyl-2-(6-phenoxy-pyridin-3-yl)-oxazol-4-yl]ethoxy}-phenyl)-propionic acid

25


To a suspension of NaH (20 mg, 0.48 mmol) in DMF (10 mL) at ambient temperature was added phenol (45 mg, 0.48 mmol). The mixture was stirred at 60°C for 20 min, and 3-(2-(2-isopropoxycarbonyl-ethyl)-4-{2-[5-methyl-2-(6-chloro-pyridin-3-yl)-oxazol-4-yl]ethoxy}-phenyl)-propionic acid t-butyl ester (180 mg, 0.32 mmol) was added. The resulting mixture was stirred at 60°C overnight, cooled, diluted with EtOAc, and washed with water/brine (4x). The organic layer was dried (MgSO₄), filtered and

-200-

5 concentrated. The residue was purified by silica gel chromatography (hexanes/EtOAc 3/1 to 1/1) to afford the intermediate ester as an oil (60 mg, 30%). The title compound was generated using Standard hydrolysis procedure C. MS (ESI) m/z 560 ($M+H$)⁺. ¹H NMR (CDCl₃, 250 MHz) δ 1.25 (d, $J=0.5$ Hz, 6H), 2.42 (s, 3H), 2.65 (m, 2H), 2.96 (m, 4H), 4.38 (m, 2H), 4.18 (m, 2H), 4.38 (m, 1H), 5.00 (m, 2H), 6.82 (m, 3H), 6.96 (m, 1H), 7.12
10 (m, 1H), 7.20 (m, 2H), 7.45 (m, 2H), 8.28 (m, 1H), 8.80 (m, 1H).

Example 322

3-[4-[2-(3-Biphenyl-4-yl-5-methyl-pyrazol-1-yl)-ethoxy]-2-(isopropoxycarbonyl-amino-methyl)-phenyl]-propionic acid



15

The title compound was prepared from 3-[4-hydroxy-2-(isopropoxycarbonyl-amino-methyl)-phenyl]-propionic acid tert-butyl ester (Prep 16) and 2-(3-biphenyl-4-yl-5-methyl-pyrazol-1-yl)-ethanol (Prep 11) using Standard coupling procedure B followed by Standard hydrolysis procedure C. MS [ES] m/e 542 ($M+1$)⁺.


20

-201-

5 The following Examples 323 to 333 are prepared by following a substantially similar procedure as described in Example 322.

Examples 323

10 3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-3-phenyl-pyrazol-1-yl)-ethoxy]-phenyl}-propionic acid




HRMS Calculated for $C_{26}H_{32}N_3O_5$: m/z 466.2342. Found: 466.2331.

15

Examples 324

3-(4-[2-(3-Biphenyl-4-yl-5-methyl-pyrazol-1-yl)-ethoxy]-2-[[pyridine-2-carbonyl]-amino]-methyl)-phenyl)-propionic acid trifluoroacetate



HRMS Calculated for $C_{28}H_{30}N_4O_4$: m/z 486.2267. Found: 486.2233.


20

-202-

5

Examples 325

3-[4-[2-(5-Biphenyl-4-yl-3-methyl-pyrazol-1-yl)-ethoxy]-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




HRMS Calculated for $C_{32}H_{36}N_3O_5$: m/z 542.2655. Found: 542.2678

10

Examples 326


3-[4-{2-[3-(4-Bromo-phenyl)-5-methyl-pyrazol-1-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester



15 HRMS Calculated for $C_{30}H_{39}BrN_3O_5$: m/z 600.2073. Found: 600.2064;

Examples 327

3-[4-{2-[3-(4-Bromo-phenyl)-5-methyl-pyrazol-1-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



20


HRMS Calculated for $C_{26}H_{31}BrN_3O_5$: m/z 544.1447. Found: 544.1456.

-203-

5

Examples 328

3-(4-[2-(5-Methyl-3-naphthalen-2-yl-pyrazol-1-yl)-ethoxy]-2-[[pyridine-2-carbonyl]-amino]-methyl}-phenyl)-propionic acid hydrochloride




The above compound is prepared by using Standard hydrolysis procedure D. HRMS

10 Calculated for $C_{32}H_{31}N_4O_4$: m/z 535.2345. Found: 535.2343

Examples 329

3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-3-naphthalen-2-yl-pyrazol-1-yl)-ethoxy]-phenyl}-propionic acid




15

HRMS Calculated for $C_{30}H_{34}N_3O_5$: m/z 516.2498. Found: 516.2485.

Examples 330

3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-3-naphthalen-1-yl-pyrazol-1-yl)-ethoxy]-phenyl}-propionic acid

20




HRMS Calculated for $C_{30}H_{34}N_3O_5$: m/z 516.2498. Found: 516.2529.

5

Examples 331

3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-2-naphthalen-2-yl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid




HRMS Calculated for $C_{30}H_{33}N_2O_6$: m/z 517.2339. Found: 517.2328.

10

Examples 332


3-{4-[2-(3-Biphenyl-4-yl-5-methyl-pyrazol-1-yl)-ethoxy]-2-[(isopropoxy-carbonyl-methyl-amino)-methyl]-phenyl}-propionic acid



15 HRMS Calculated for $C_{33}H_{38}N_3O_5$: m/z 556.2811. Found: 556.2819

Examples 333

3-{4-[2-[3-(4-Bromo-phenyl)-5-methyl-pyrazol-1-yl]-ethoxy]-2-[(isopropoxy-carbonyl-methyl-amino)-methyl]-phenyl}-propionic acid



20


HRMS Calculated for $C_{27}H_{33}BrN_3O_5$: m/z 558.1604. Found: 558.1608

-205-

5

Example 334

3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(1-methyl-4-phenyl-1H-imidazol-2-yl)-ethoxy]-phenyl}-propionic acid




The title compound was prepared from 3-[4-hydroxy-2-(isopropoxycarbonyl-amino-methyl)-phenyl]-propionic acid tert-butyl ester (Prep 16) and 2-(1-Methyl-4-phenyl-1H-imidazol-2-yl)-ethanol (*J. Med. Chem.* 1998, 41(25), 5037-5054) using Standard coupling procedure B followed by Standard hydrolysis procedure D affording an off-white finely-crystalline solid. HRMS Calculated for C₂₆H₃₁N₃O₅: m/z 466.2342. Found: 466.2355.

15

Tail Piece Aryl Ring Modifications

Example 335

3-[4-{2-[2-(4-Bromo-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonyl-amino-methyl)-phenyl]-propionic acid tert-butyl ester



20

Toluene-4-sulfonic acid 2-[2-(4-bromo-phenyl)-5-methyl-oxazol-4-yl]-ethyl ester (5.06g, 11.6mmol) and 3-[4-hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (3.01g, 8.92mmol) were coupled using Standard procedure A to give the title compound as a white solid (3.41 g, 64%). MS [ES] m/z 601,603 (M+H).


25

-206-

5 The following Examples 336 to 337 are prepared by following a substantially similar procedure as described in Example 335.

Example 336


10 3-[4-{2-[2-(3-Bromo-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxy-carbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester



MS [ES] m/z 601,603 (M+H).

Example 337


15 3-[4-{2-[2-(2-Bromo-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxy-carbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester.



5

Example 338

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-pyridin-3-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



Suzuki Coupling Using Aryl Boronic Acids: A mixture of 3-[4-{2-[2-(4-bromo-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (450 mg, 0.75 mmol), 3-pyridineboronic acid (120 mg, 0.97 mmol), triphenylphosphine (6 mg, 0.024 mmol), 2M aqueous sodium carbonate (3mL) in n-propanol (15mL) was sparged with nitrogen for 5 min. Palladium acetate (2 mg, 0.008 mmol) was added, and the reaction was heated to reflux under a blanket of nitrogen for 4 h. The reaction mixture was concentrated and then partitioned between water/EtOAc. The organic layer was dried (MgSO₄) and concentrated to crude product (290 mg). This material was purified by silica gel column chromatography to yield 3-(2-(isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-pyridin-3-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid tert-butyl ester was obtained as a white solid (230 mg, 51%). ¹H NMR (CDCl₃) δ 1.25 (d, 6H), 1.42 (s, 9H), 2.42 (s, 3H), 2.51 (t, 2H), 2.89 (t, 2H), 3.05 (t, 2H), 4.26 (t, 2H), 4.36 (d, 2H), 4.96 (m, 2H), 6.78 (dd, 1H), 6.87 (d, 1H), 7.21 (d, 1H), 7.47 (dd, 1H), 7.67 (d, 2H), 8.00 (dt, 1H), 8.12 (d, 2H), 8.65 (dd, 1H), 8.94 (s, 1H). MS [EI+] m/z 600 (M+H).


This ester (225 mg, 0.375 mmol) was dissolved in THF (15mL) and treated with 1N HCl (1mL). The mixture was heated at reflux for 7 h, cooled, and concentrated. The residue was azeotroped twice with acetonitrile and dried in a vacuum oven to give the title compound, isolated as the HCl salt (140 mg, 64%). ¹H NMR (DMSO-d₆) δ 1.15 (d, 6H), 2.41 (s, 3H), 2.45 (t, 2H), 2.76 (t, 2H), 2.95 (t, 2H), 4.17 (overlapping t and d, 4H), 4.75 q, 1H), 6.77 (m, 2H), 7.07 (dd, 1H), 7.55 (t, 1H), 8.02 (m, 4H), 8.73 (d, 1H), 8.85 (d, 1H), 9.25 (s, 1H). MS [EI+] m/z 544 (M+H).

-208-

5 The following Examples 339 to 367 are prepared by following a
substantially similar procedure as described in Example 338

Example 339


10 3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-pyridin-3-ylphenyl)oxazol-
4-yl]ethoxy} phenyl) propionic acid



MS (ES) m/z 544 (M+1).

Example 340

15 3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-pyridin-4-yl-phenyl)oxazol-
4-yl]ethoxy} phenyl) propionic acid



MS (ES) m/z 544 (M+1).


20

-209-

5

Example 341

3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[5-methyl-2-(3-pyridin-3-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid




MS (ES) m/z 544 (M+1).

10

Example 342


3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[5-methyl-2-(3-pyridin-4-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid



15 MS (ES) m/z 544 (M+1).

Example 343

3-[4-{2-[2-(4'-Fluorobiphenyl-4-yl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminoethyl)phenyl] propionic acid



20


MS (ES) m/z 559 (M+1).

-210-

5

Example 344

3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4'-trifluoromethylbiphenyl-4-yl)oxazol-4-yl]ethoxy}phenyl) propionic acid




MS (ES) m/z 609 (M+1).

10

Example 345


3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(2'-trifluoromethylbiphenyl-4-yl)oxazol-4-yl]ethoxy}phenyl) propionic acid



15 MS (ES) m/z 609 (M+1).

Example 346

3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[2-(4'-methoxybiphenyl-4-yl)-5-methyloxazol-4-yl]ethoxy}phenyl) propionic acid



20


MS (ES) m/z 571 (M+1).

-211-

5

Example 347


3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[2-(3'-methoxybiphenyl-4-yl)-5-methyloxazol-4-yl]ethoxy}phenyl) propionic acid



¹H-NMR (DMSO-d₆) δ 1.16 (d, 6H), 2.39 (s, 3H), 2.45 (t, 2H), 2.76 (t, 2H), 2.95 (t, 2H),
 10 3.85 s, 3H), 4.17 (m, 4H), 4.77 (m, 1H), 6.74-7.55 (m, 7H, 7.83 (d, 2H), 7.99 (d, 2H).

Example 348

3-[4-{2-[2-(4'-Fluorobiphenyl-3-yl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid




15

MS (ES) m/z 559 (M+1).

5

Example 349

3-[4-{2-[2-(3'-Fluorobiphenyl-3-yl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid




MS (ES) m/z 546 (M+1).

10

Example 350

3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4'-trifluoromethylbiphenyl-3-yl)oxazol-4-yl]ethoxy}phenyl) propionic acid




15 ¹H-NMR (DMSO-d₆) δ 1.15 (d, 6H), 2.40 (s, 3H), 2.45 (t, 2H), 2.79 (t, 2H), 2.96 (t, 2H), 4.17 (m, 4H), 4.75 (m, 1H), 6.77 (m, 2H), 7.08 (d, 1H), 7.53 (m, 1H), 7.66 (m, 1H), 7.83-8.05 (m, 5H), 8.20 (s, 1H).

-213-

5

Example 351

3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[5-methyl-2-(3'-trifluoromethylbiphenyl-3-yl)oxazol-4-yl]ethoxy}phenyl) propionic acid




¹H-NMR (DMSO-d₆) δ 1.14 (d, 6H), 2.40 (s, 3H), 2.45 (t, 2H), 2.67 (t, 2H), 2.95 (t, 2H), 4.16 (m, 4H), 4.74 (m, 1H), 6.77 (m, 2H), 7.09 d, 1H), 7.47-8.20 (m, 8H).

10

Example 352

3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[5-methyl-2-(2'-trifluoromethylbiphenyl-3-yl)oxazol-4-yl]ethoxy}phenyl) propionic acid



15


¹H-NMR (DMSO-d₆) δ 1.13 (d, 6H), 2.36 (s, 3H), 2.44 (t, 2H), 2.77 (t, 2H), 2.93 (t, 2H), 4.15 (m, 4H), 4.74 (m, 1H), 6.76 (m, 2H), 7.06 (d, 1H), 7.40-8.00 (m, 8H).

-214-

5

Example 353

3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[2-(3'-methoxybiphenyl-3-yl)-5-methyloxazol-4-yl]ethoxy}phenyl) propionic acid




MS (ES) m/z 573 (M+1).

10

Example 354

3-(2-(Isopropoxycarbonylaminoethyl)-4-{2-[2-(2'-methoxybiphenyl-3-yl)-5-methyloxazol-4-yl]ethoxy}phenyl) propionic acid




15 $^1\text{H-NMR}$ (DMSO- d_6) δ 1.13 (d, 6H), 2.40 (s, 2H), 2.45 (t, 2H), 2.78 (t, 2H), 2.93 (t, 2H), 3.80 (s, 3H), 4.16 (m, 4H), 4.75 (m, 1H), 6.76 (m, 2H), 7.07 (m, 2H), 7.16 (d, 1H), 7.32-8.05 (m, 6H).

-215-

5

Example 355

3-[4-{2-[2-(3'-Fluorobiphenyl-4-yl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid




MS (ES) m/z 559 (M+1).

10

Example 356

3-[4-{2-[2-(2'-Fluoro-biphenyl-4-yl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylaminomethyl)-phenyl]-propionic acid




15 ¹H-NMR (CDCl₃) δ 1.25 (d, 6H), 2.41 (s, 3H), 2.65 (t, 2H), 2.95 (t, 2H), 3.01 (t, 2H), 4.24 (t, 2H), 4.38 (br d, 2H), 4.96 (m, 1H), 5.03 (br s, 1H), 6.78 (dd, 1H), 6.84 (d, 1H), 7.19-7.55 (m, 6H), 7.65 (dd, 2H), 8.06 dd, 2H).

-216-

5

Example 357

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(3'-trifluoromethylbiphenyl-4-yl)oxazol-4-yl]ethoxy}phenyl) propionic acid




MS (ES) m/z 561 (M+1).

10

Example 358


3-[4-[2-(2-Biphenyl-2-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



15 MS (ES) m/z 543.2 (M+1)⁺.

Example 359

4'-(4-{2-[4-(2-Carboxy-ethyl)-3-(isopropoxycarbonylamino-methyl)-phenoxy]-ethyl}-5-methyl-oxazol-2-yl)-biphenyl-3-carboxylic acid



20


The above compound is prepared after an additional hydrolysis step using Standard procedure E. MS (ES) m/z 587.3 (M+1)⁺.

-217-

5

Example 360


4'-(4-{2-[4-(2-Carboxy-ethyl)-3-(isopropoxycarbonylamino-methyl)-phenoxy]-ethyl}-5-methyl-oxazol-2-yl)-biphenyl-4-carboxylic acid



- 10 The above compound is prepared after an additional hydrolysis step using Standard procedure E. MS (ES) m/z 587.2 ($M+1$)⁺.

Example 361


- 15 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-3-(4-pyridin-4-yl-phenyl)-pyrazol-1-yl]-ethoxy}-phenyl)-propionic acid hydrochloride



HRMS Calculated for $C_{31}H_{35}N_4O_5$: m/z 543.2607. Found: 543.2614.

Example 362

- 20 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-3-(4-pyridin-3-yl-phenyl)-pyrazol-1-yl]-ethoxy}-phenyl)-propionic acid hydrochloride



HRMS Calculated for $C_{31}H_{35}N_4O_5$: m/z 543.2607. Found: 543.2612.


25

-218-

5

Example 363

3-[4-{2-[3-(4'-Fluoro-biphenyl-4-yl)-5-methyl-pyrazol-1-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




HRMS Calculated for $C_{28}H_{35}FN_3O_5$: m/z 560.2561. Found: 560.2575.

10

Example 364


3-[4-{2-[3-(4'-Methoxy-biphenyl-4-yl)-5-methyl-pyrazol-1-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



15 HRMS Calculated for $C_{33}H_{38}N_3O_6$: m/z 572.2761. Found: 572.2752.

Example 365

3-[4-{2-[3-(3'-Methoxy-biphenyl-4-yl)-5-methyl-pyrazol-1-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



20


HRMS Calculated for $C_{33}H_{38}N_3O_6$: m/z 572.2761. Found: 572.2776.

-219-

5

Example 366

3-[4-{2-[3-(2'-Fluoro-biphenyl-4-yl)-5-methyl-pyrazol-1-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




HRMS Calculated for $C_{32}H_{35}FN_4O_5$: m/z 560.2561. Found: 560.2540.

10

Example 367

3-[4-{2-[3-(2'-Methyl-biphenyl-4-yl)-5-methyl-pyrazol-1-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




15 HRMS Calculated for $C_{33}H_{38}N_4O_5$: m/z 556.2811. Found: 556.2802.

-220-


5

Example 368

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-pyrazin-2-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



10 Step A: 3-[2-(Isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid



A solution of 3-[4-{2-[2-(4-bromo-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (4.00 g, 6.65 mmol) in DMSO (50 mL) was treated with bis(pinacolato)diborane (2.20 g, 8.64 mmol) and potassium acetate (1.96 g, 19.95 mmol). The solution was sparged with nitrogen for 10min, Pd(dppf)Cl₂ (980 mg, 1.20 mmol; 1:1 complex with CH₂Cl₂) was added, and the reaction mixture was heated at 80 °C for 4 h. The mixture was partitioned between water and ethyl acetate. The organic layer was dried (MgSO₄), filtered, and concentrated. Purification by silica gel chromatography (9:1 hexanes:ethyl acetate) yielded the title compound as a yellow oil (2.81 g). ¹H NMR (300MHz CDCl₃) δ 1.25 (d, 6H), 1.26 (s, 9H), 1.37 (s, 12H), 2.40 (s, 3H), 2.51 (t, 2H), 2.88 (t, 2H), 2.98 (t, 2H), 4.26 (t, 2H), 4.35 (d, 2H), 4.97 (br m, 2H), 6.77 (dd, 1H), 6.85 (d, 1H), 7.10 (d, 1H), 7.87 (d, 2H), 7.99 (d, 2H). MS [EI+] 649 (M+H).

-221-

- 5 Step B: 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-pyrazin-2-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid


Suzuki coupling using haloaryl compounds: 3-[2-(Isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[4-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid (500 g, 0.771 mmol), CsF (258 g, 1.69 mmol), PdCl₂(dppf) (0.057 g, 0.07 mmol), and 2-chloropyrazine (0.0971g, 0.85 mmol) were added to a 3-neck flask and were dissolved in anhydrous dioxane (25 mL). The reaction mixture was stirred at 100°C under a stream of N₂ for about 12 h. The reaction mixture was diluted with ethyl acetate (200 mL) and washed with saturated NaCl (100 mL) and water (100 mL). The organic layer was filtered
15 through Celite, dried (Na₂SO₄), filtered, and concentrated. The crude product (1.2 g) was purified by radial chromatography (10-70% EtOAc/hexanes) to give the intermediate ester (0.312g): MS (ES) m/z 601 (M+H)⁺. The ester was dissolved in of 4M HCl/dioxane (10 mL). The mixture was stirred N₂ for 16 h and concentrated to give the title compound (305 mg, 72%): MS (ES) m/z 545 (M+H)⁺. Anal. Calculated for C
20 66.16%, H 5.92%, N 10.29%. Found C 65.97%, H 6.07%, N 10.45%.

The following Examples 369 to 382 are prepared by following a substantially similar procedure as described in Example 368.

25

Example 369

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-pyridin-2-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid



MS (ES) m/z 544 (M+1).


30

-222-

5

Example 370

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(5-methylpyridin-2-yl)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




MS (ES) m/z 558 (M+1).

10

Example 371


3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(3-methylpyridin-2-yl)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid



15 MS (ES) m/z 558 (M+1).

Example 372

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(6-methylpyridin-2-yl)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid



20


MS (ES) m/z 492 (M+1).

-223-

5

Example 373

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(4-methylpyridin-2-yl)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




MS (ES) m/z 558 (M+1).

10

Example 374

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(4-trifluoromethylpyridin-2-yl)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




15 MS (ES) m/z 612 (M+1).

5

Example 375

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(5-trifluoromethylpyridin-2-yl)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




MS (ES) m/z 612 (M+1).

10

Example 376

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{2-[4-(6-methoxypyridin-3-yl)phenyl]-5-methyloxazol-4-yl}ethoxy)phenyl] propionic acid




15 MS (ES) m/z 574 (M+1).

-225-

5

Example 377

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{2-[4-(6-methoxypyridin-2-yl)phenyl]-5-methyloxazol-4-yl}ethoxy)phenyl] propionic acid




MS (ES) m/z 574 (M+1).

10

Example 378

3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-quinolin-4-ylphenyl)oxazol-4-yl]ethoxy} phenyl) propionic acid




15 MS (ES) m/z 594 (M+1).

-226-

5

Example 379

3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-pyrazin-2-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid




MS (ES) m/z 545 (M+1).

10

Example 380

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{2-[4-(4-methoxypyridin-2-yl)phenyl]-5-methyloxazol-4-yl}ethoxy)phenyl] propionic acid




15 MS (ES) m/z 574 (M+1).

5

Example 381

3-[4-(2-{2-[4-(5-Cyanopyridin-2-yl)phenyl]-5-methyloxazol-4-yl}ethoxy)-2-(isopropoxycarbonylaminoethyl)phenyl] propionic acid




MS (ES) m/z 569 (M+1).

10

Example 382

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-pyrimidin-2-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid:




15 MS (ES) m/z 545 (M+H)⁺. Anal. Calculated for C 66.16%, H 5.92%, N 10.29%, Found C 65.87%, H 6.17%, N 10.53%.

-228-

5

Example 383

3-[4-(2-{2-[4-(4-Fluoro-phenylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




- 3-[4-{2-[2-(4-bromo-phenyl)-5-methoxy-oxazol-4-yl]-ethoxy}-2-(isopropoxy-carbonylamino-methyl)-propionic acid tert-butyl ester (0.10 g, 0.17 mmol) was dissolved in DME (5 mL) in a sealed tube apparatus. N₂ was bubbled through the solution, and 4-fluoroaniline (24 mg, 0.20 mmol), 2-(di-t-butylphosphino)-biphenyl (20 mg), Pd₂(dba)₃ (10 mg), and K₃PO₄ (50 mg, 0.24 mmol) were added. The tube was sealed and heated at 100 °C for 16 h. The reaction mixture was cooled, diluted with EtOAc, and washed with water. The organic layer was dried (MgSO₄), filtered, and concentrated. The residue was purified using silica gel chromatography (hexanes/EtOAc 2/1 to 1/1). Using the Standard hydrolysis procedure C, this material was converted into the title compound as an oil (25 mg): MS (ESI) m/z 594 (M+H)⁺.

-229-

5 The following Examples 384 to 387 are prepared by following a substantially similar procedure as described in Example 383.

10 Example 384


3-[4-(2-{2-[4-(4-Cyano-phenylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



MS (ES) m/z 583 (M+1).

15 Example 385

3-[4-(2-{2-[4-(3,5-Difluoro-phenylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




MS (ES) m/z 594 (M+1).

20

5

Example 386

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-p-tolylamino-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid




MS (ES) m/z 572 (M+1).

10

Example 387

3-[2-(Isopropoxycarbonylamino-methyl)-4-(2-{2-[4-(4-methoxy-phenylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid




15 MS (ES) m/z 588 (M+1).

-231-

5

Example 390

3-[4-(2-{2-[3-Benzylamino-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




- 3-[4-{2-[2-(3-bromo-phenyl)-5-methoxy-oxazol-4-yl]-ethoxy}-2-(isopropoxy-carbonylamino-methyl)-propionic acid tert-butyl ester (0.10 g, 0.17 mmol) was dissolved in toluene (5 mL) in a sealed tube apparatus. N₂ was bubbled through the solution, and benzylamine (46 μ L, 0.42 mmol), 2-(dicyclohexylphosphino)-biphenyl (12 mg), Pd₂(dba)₃ (16 mg) and t-BuONa (23 mg, 0.24 mmol) were added. The tube was sealed and heated at 100 °C for 16 h. The reaction mixture was cooled, diluted with EtOAc, and washed with water. The organic layer was dried (MgSO₄), filtered, and concentrated. The residue was purified using silica gel chromatography (hexanes/EtOAc 2/1 to 1/1). Using the Standard hydrolysis procedure C, this material was converted into the corresponding carboxylic acid; the title compound was isolated as the HCl salt by treatment with 1M HCl in ether (15 mg). MS (ES) m/z 572 (M+H)⁺.

20

Example 391

3-[4-{2-[2-(4-Diethylamino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




- The above compound is prepared by following a substantially similar procedure as described in Example 390 except that the reaction is carried out in DME at 80°C.

5 MS (ES) m/z 538.2 (M+H)⁺.

Example 392

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-morpholin-4-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



10

A solution of 3-[4-{2-[2-(4-bromo-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (60 mg, 0.100 mmol) in toluene (3.0 mL) in a sealed tube apparatus under a flow of N₂ was treated with Pd(OAc)₂ (5 mg), 2-(di-*t*-butylphosphino)biphenyl (10 mg), morpholine (17 mg, 0.20 mmol), and sodium *t*-butoxide (19 mg, 0.200 mmol). The tube was sealed and heated at 105 °C for 14 h. The mixture was cooled and purified directly using silica gel chromatography (30-50% EtOAc/hexanes) to yield the intermediate ester. A solution of ester in TFA (1.0 mL)/CH₂Cl₂ (1.5 mL)/H₂O (0.1 mL) and stirred 14 h and concentrated. The residue was and purified using silica gel chromatography (hexanes/EtOAc/HOAc, 5/5/0.02) to afford the title compound (36 mg, 65%): MS (ESI) m/z 552.3 (M+H)⁺.

15


20

-233-

5 The following Examples 393 to 395 are prepared by following a substantially similar procedure as described in Example 392.

Example 393


10 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(3-morpholin-4-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



MS (ESI) m/z 552.2 (M+H)⁺.

Example 394

15 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-piperidin-1-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



MS (ESI) m/z 550.3 (M+H)⁺.


20

-234-

5

Example 395

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(3-piperidin-1-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid




MS (ESI) m/z 550.3 ($M+H$)⁺.

10

Example 396

3-[2-(Isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[4-(morpholin-4-ylamino)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid




15

A solution of 3-[4-{2-[2-(4-bromo-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (120 mg, 0.200 mmol) in toluene (2.0 mL) in a sealed tube under a flow of N₂ was treated Pd₂(dba)₃ (10 mg), 2-(di-*t*-butylphosphino)biphenyl (10 mg), N-aminomorpholine (29 mg, 0.28 mmol), and sodium *t*-butoxide (38 mg, 0.40 mmol). The tube was sealed and heated at 100 °C for 60 min. The mixture was cooled and purified using silica gel chromatography (5-10% MeOH/EtOAc) to yield the title compound directly (22 mg, 21%): MS (ESI) m/z 567.3 ($M+H$)⁺.

20

5 Example 397

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-phenoxy-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



A mixture of 3-[4-{2-[2-(4-bromo-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl] -propionic acid tert-butyl ester (0.44 mmol, 0.24 g), phenol (0.53 mmol, 0.050 g), K_3PO_4 (0.88 mmol, 0.19g), 2-(di-t-butylphosphino)biphenyl (0.066 mmol, 0.020g), and $Pd(OAc)_2$ (0.044 mmol, 0.010g) in toluene (7 mL) was degassed under vacuum and backfilled with nitrogen (3x) and heated at 110 °C for 18 h. Additional $Pd(OAc)_2$ (10 mg) and 2-(di-t-butylphosphino)biphenyl (20 mg) were added to ensure complete reaction, and the mixture was heated 5 h. Upon cooling, the mixture was placed directly onto a silica gel column and eluted with 30%-50% EtOAc/hexanes to give 3-(2-(isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-phenoxy-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid tert-butyl ester (0.11 g).


The tert-butyl ester (0.18 mmol, 0.11 g) was dissolved in CH_2Cl_2 (3 mL), and 90% TFA/water (5 mL) was added. The mixture was stirred for 3 h and concentrated. The residue was purified by silica gel chromatography (50% ethyl acetate/hexanes) to give an oil. Trituration with Et_2O gave the title compound as a white solid (90 mg, 90%): 1H NMR (250 MHz, $CDCl_3$) δ 8.11 (d, 2H, $J=8.8$ Hz), 7.56 (t, 2H, $J=7.9$ Hz), 7.37-7.20 (m, 6H), 7.0-6.93 (m, 2H), 5.11 (heptet, 1H, $J=6.3$ Hz), 4.53 (d, 2H, $J=5.1$ Hz), 4.38 (t, 2H, $J=6.7$ Hz), 3.12 (q, 4H, $J=7.3$ Hz), 2.80 (t, 2H, $J=7.6$ Hz), 2.54 (s, 3H), 1.41 (d, 6H, $J=6.3$ Hz).

-236-

5 The following Examples 398 to 405 are prepared by following a substantially similar procedure as described in Examples 396 and 397.

Example 398


10 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(3-phenoxy-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



MS (ESI) m/z 559.2 (M+H)⁺.

Example 399

15 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(2-phenoxy-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



MS (ESI) 559 (M+H)⁺. Anal Calculated for C₃₂H₃₄N₂O₇: C, 68.8; H, 6.1; N, 5.0. Found: C, 67.9; H, 6.2; N, 5.3.


20

-237-

5

Example 400

3-[4-(2-{2-[4-(4-Cyano-phenoxy)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




MS (ES) m/z 584.

10

Example 401


3-[2-(Isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[4-(4-trifluoromethyl-phenoxy)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid



15 MS (ES) m/z 627 (M+1).

Example 402

3-[4-(2-{2-[4-(4-Fluoro-phenoxy)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




20 MS (ES) m/z 577 (M+1).

-238-

5

Example 403

3-[4-(2-{2-[4-(3,4-Difluoro-phenoxy)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




MS (ES) m/z 595 (M+1).

10

Example 404


3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-m-tolyloxy-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



15 MS (ES) m/z 573 (M+1).

Example 405

3-[4-(2-{2-[4-(4-Acetyl-phenoxy)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



20


MS (ES) m/z 601 (M+1).

-239-

5

Example 406

3-[4-{2-[2-(4-Hydroxy-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester




According to the procedure in *Organic Synthesis*, vol. V, p. 918, a solution
 10 of 3-[2-(isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[4-(4,4,5,5-tetramethyl-
 [1,3,2]dioxaborolan-2-yl)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid tert-butyl
 ester (1.4g, 2.16mmol, Example 368, Step A) in THF (8mL) was treated with glacial
 acetic acid (194mg, 3.24mmol) at 0°C. A solution of 30% H₂O₂ in water (4.75mL) was
 diluted with H₂O (1 mL) and added to the reaction mixture. The mixture was warmed to
 15 room temperature and treated with 1M aqueous Na₂S₂O₃ solution (50mL). The THF was
 removed under vacuum and the aqueous phase was extracted with ethyl acetate. The
 organic layer was washed with 1M Na₂S₂O₃ (2 x 50mL), dried (MgSO₄), filtered, and
 concentrated to an oil containing (1.21 g). Purification using silica gel chromatography
 (1:1 ηεξανεσ:ετηψλ acetate) yielded the title compound as a white solid (1.10 g, 95%).
 20 ¹H NMR (300MHz CDCl₃) δ 1.15 (d, 6H), 1.19 (s, 9H), 2.26 (s, 3H), 2.39 (t, 2H), 2.75 (t,
 2H), 2.85 (t, 2H), 4.10 (t, 2H), 4.28 (d, 2H), 4.85 (m, 1H), 5.01 (br d, 1H), 6.63 (dd, 1H),
 6.74 (m, 3H), 6.96 (d, 1h), 7.70 (d, 2H); MS [EI+] m/z 539 (M+H).

-240-

5

Example 407

3-[4-{2-[2-(3-Hydroxy-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester:




10 The above compound is prepared by following a substantially similar procedure as described in Example 406. MS [EI+] m/z 539 (M+H).

The following Examples 408 to 409 are prepared by following a substantially similar procedure as described in Example 406 and the corresponding carboxylic acids are obtained from Standard hydrolysis procedure C:

15

Example 408

3-[4-{2-[2-(4-Hydroxyphenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylamino-methyl)phenyl] propionic acid




20 MS (ES) m/z 483 (M+1).

-241-

5

Example 409

3-[4-{2-[2-(3-Hydroxyphenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminoethyl)phenyl] propionic acid:




MS (ES) m/z 483 (M+1).

10

Example 410

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[2-(4-isopropoxy-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



15

A solution of 3-[4-{2-[2-(4-hydroxy-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (300mg, 0.557mmol, Example 406) in ethanol (20mL) was treated with 2-iodopropane (473mg, 2.78mmol) and K_2CO_3 (231mg, 1.67mmol) and was heated at reflux overnight. The reaction mixture was cooled and concentrated. The residue was diluted with H_2O and ethyl acetate. The layers were separated, and the aqueous layer was extracted with ethyl acetate (2 x). The combined organic layers were dried ($MgSO_4$), filtered, and concentrated to yield an oil. Purification using silica gel chromatography (3:1 hexane:ethyl acetate) yielded 3-(2-(isopropoxycarbonylamino-methyl)-4-{2-[2-(4-isopropoxy-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid tert-butyl ester as a yellow oil (210mg). 1H NMR (300MHz $CDCl_3$) δ 1.28 (d, 6H), 1.40 (d, 6H), 1.45 (s, 9H), 2.38 (s, 3H), 2.53 (t, 2H), 2.91 (t, 2H), 2.99 (t, 2H), 4.25 (t, 2H), 4.39 (d,

20

25

-242-

5 2H), 4.66 (m, 1H), 5.00 (br m, 2H), 6.80 (dd, 1H), 6.87 (d, 1H), 6.95 (d, 2H), 7.13 (d, 1H), 7.95 (d, 2H) MS [ES] 581 (M+H).


This ester was converted to the title compound according to Standard Procedure C; a white solid (134 mg). ¹H NMR (300MHz CDCl₃) δ 1.15 (d, 6H), 1.29 (d, 6H), 2.31 (s, 3H), 2.53 (t, 2H), 2.82 (t, 2H), 2.97 (t, 2H), 4.15 (t, 2H), 4.26 (d, 2H), 4.55 (m, 1H), 4.84 (m, 1H), 4.99 (br s, 1H), 6.68 (dd, 1H), 6.73 (d, 1H), 6.87 (d, 2H), 7.00 (d, 1H), 7.90 (d, 2H) MS [ES] m/z 525 (M+H).

The following Examples 411 to 422 are prepared by following a substantially similar procedure as described in Example 410.

15

Example 411


3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-propoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid



20 MS (ES) m/z 525 (M+1).

Example 412

3-[4-{2-[2-(4-Ethoxyphenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid



25

MS (ES) m/z 511 (M+1).

-243-

5

Example 413

3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[2-(4-methoxyphenyl)-5-methyloxazol-4-yl]ethoxy}phenyl) propionic acid

MS (ES) m/z 497 (M+1).

10

Example 414


3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[2-(3-methoxyphenyl)-5-methyloxazol-4-yl]ethoxy}phenyl) propionic acid

MS (ES) m/z 497 (M+1)

15

Example 415


3-[4-{2-[2-(3-Ethoxyphenyl)-5-methyloxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid



20 MS (ES) m/z 511 (M+1).

Example 416

3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[2-(3-isopropoxyphenyl)-5-methyloxazol-4-yl]ethoxy}phenyl) propionic acid



25


MS (ES) m/z 525 (M+1).

-244-

5

Example 417

3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(3-propoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid




MS (ES) m/z 525 (M+1).

10

Example 418


3-[4-{2-[2-(3-Butoxyphenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid



15 MS (ES) m/z 539 (M+1).

Example 419

3-[4-{2-[2-(3-Cyclopentyloxyphenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid



20


MS (ES) m/z 551 (M+1).

-245-

5

Example 420

3-[4-{2-[2-(3-Cyclohexyloxyphenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)-phenyl] propionic acid




MS (ES) m/z 565 (M+1).

10

Example 421


3-[4-{2-[2-(4-Cyclopentyloxyphenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid



15 MS (ES) m/z 551 (M+1).

Example 422

3-[4-{2-[2-(4-Cyclohexyloxyphenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid



20


MS (ES) m/z 565 (M+1).

-246-

5

Example 423

3-[2-(isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[4-(tetrahydro-pyran-4-yloxy)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid



A mixture of 3-[4-{2-[2-(4-hydroxy-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (120 mg, 0.223 mmol, Example 406), tetrahydro-pyran-4-ol (22.7 mg, 0.223 mmol), triphenylphosphine (58.4 mg, 0.223 mmol) and toluene (10 mL) was treated dropwise with DIAD (45 mg, 0.223 mmol). The mixture was stirred under N₂ at ambient temperature for 16 h and concentrated. The crude product was purified by radial chromatography (10-70% EtOAc/hexanes) to give 3-[2-(isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[4-(tetrahydro-pyran-4-yloxy)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid tert-butyl ester. The ester product was dissolved 4M HCl/dioxane (5 mL), stirred for 16 h and concentrated to give the title compound: MS [ES] m/z 525 (M+H).


20

-247-

5 The following Examples 424 to 431 are prepared by following a
substantially similar procedure as described in Example 423.

10 Example 424


3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(1-methylpiperidin-4-
yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid



MS (ES) m/z 580 (M+1).

15 Example 425

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[3-(tetrahydropyran-4-
yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid



MS (ES) m/z 567 (M+1).


20

-248-

5

Example 426

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[3-(1-methylpiperidin-4-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




MS (ES) m/z 580 (M+1).

10

Example 427

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[3-(piperidin-4-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




MS (ES) m/z 566 (M+1).

15

Example 428

3-[4-(2-{2-[3-(3-Dimethylaminopropoxy)phenyl]-5-methyloxazol-4-yl}ethoxy)-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid




20 MS (ES) m/z 568 (M+1).

-249-

5

Example 429

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(piperidin-4-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




MS (ES) m/z 566 (M+1).

10

Example 430


3-[4-(2-{2-[4-(2-Dimethylaminoethoxy)phenyl]-5-methyloxazol-4-yl}ethoxy)-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid



15 MS (ES) m/z 554 (M+1).

Example 431

3-[4-(2-{2-[4-(3-Dimethylaminopropoxy)phenyl]-5-methyloxazol-4-yl}ethoxy)-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid



20


MS (ES) m/z 568 (M+1).

-250-

5

Example 432

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(pyridin-2-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




10 3-[4-{2-[2-(4-hydroxy-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxy-carbonylamino-methyl)-phenyl]-propionic acid tert-butyl ester (Example 406) was reacted with 2-bromopyridine by the procedure in Example 397 to give the title compound. MS (ES) m/z 560 (M+1).

15 The following Examples 433 to 435 are prepared by following a substantially similar procedure as described in Example 432.

Example 433

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(pyridin-4-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid



20


MS (ES) m/z 560 (M+1).

-251-

5

Example 434

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(pyridin-3-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




MS (ES) m/z 560 (M+1).

10

Example 435

3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(pyrimidin-2-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid




15 MS (ES) m/z 561 (M+1).


5

Example 436


3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-carbamoyl-phenyl)-oxazol-4yl]-ethoxy}-phenyl)-propionic acid



10 Step A: 4-(4-{2-[4-(2-tert-Butoxycarbonyl-ethyl)-3-(isopropoxycarbonylamino-methyl)-phenoxy]-ethyl}-5-methyl-oxazol-2-yl)-benzoic acid methyl ester



15 A mixture of 3-[4-{2-[2-(4-bromo-phenyl)-5-methoxy-oxazol-4-yl]-ethoxy}-2-(isopropoxy-carbonylamino-methyl-phenyl)-propionic acid tert-butyl ester (0.25 g, 0.42 mmol), 1,1'-bis(diphenylphosphino)-ferrocene palladium (II) chloride (50 mg), MeOH (0.1 mL) and triethylamine (0.12 mL, 0.66 mmol) in acetonitrile (2 mL) was stirred and heated at 70°C under CO gas (balloon) for 16 h. The reaction mixture was cooled and concentrated. The residue was purified by silica gel chromatography (hexanes/EtOAc, 5/1 to 1/1) to afford a the title compound as a white solid (0.12 g, 50%): MS (ESI) m/z 581 (M+H)⁺.



20

A solution of 4-(4-{2-[4-(2-tert-Butoxycarbonyl-ethyl)-3-(isopropoxycarbonyl-amino-methyl)-phenoxy]-ethyl}-5-methyl-oxazol-2-yl)-benzoic acid

-253-

5 methyl ester (100 mg, 0.17 mmol) in methanol (2 mL) was treated with 1.5 N aqueous LiOH (1.0 mL) and stirred at ambient temperature for 3 h. The reaction mixture was acidified with 1N HCl and extracted with CH₂Cl₂. The organic layer was dried (MgSO₄), concentrated, and purified by silica gel chromatography (hexanes/EtOAc, 1/1 to 0/1) to afford the title compound (55 mg, 57%). MS (ESI) m/z 567 (M+H)⁺.

10

Step C: 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-carbamoyl-phenyl)-oxazol-4yl]-ethoxy}-phenyl)-propionic acid


A solution of 4-(4-{2-[4-(2-tert-Butoxycarbonyl-ethyl)-3-(isopropoxycarbonyl-amino-methyl)-phenoxy]-ethyl}-5-methyl-oxazol-2-yl)-benzoic acid
15 (40 mg, 0.07 mmol) in CH₂Cl₂ (2 mL) was treated with oxalyl chloride (1.0 μL, 0.12 mmol) and one drop of DMF. The reaction mixture was stirred at ambient temperature for 30 min, concentrated, and co-evaporated with toluene. The residue was dissolved in CH₂Cl₂ (2 mL) and was added to a flask charged with 2.0 M methylamine in THF (2M, 0.04 mL) and triethylamine (20 μL, 0.12 mmol). The mixture was stirred at ambient
20 temperature for 16 h, diluted with CH₂Cl₂ (10 mL), and washed with H₂O. The organic layer was dried (MgSO₄), concentrated, and purified by silica gel chromatography (hexanes/EtOAc, 3:1 to 0:1) to afford the intermediate *tert*-butyl ester. The ester was converted to the acid by Standard procedure C; purification by mass directed HPLC yielded the title compound. MS (ESI) m/z 524 (M+H)⁺.

25

5 The following Examples 437 to 448 are prepared by following a substantially similar procedure as described in Example 436.

Example 437


10 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(3-methylcarbamoyl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



MS (ES) m/z 524 (M+1).

Example 438

15 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(3-propylcarbamoyl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



MS (ES) m/z 552 (M+1).


20

-255-

5

Example 439

3-[4-{2-[2-(3-Cyclobutylcarbamoyl-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




MS (ES) m/z 564 (M+1).

10

Example 440


3-[4-{2-[2-(3-Isobutylcarbamoyl-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



15 MS (ES) m/z 566 (M+1).

Example 441

3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(3-phenylcarbamoyl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



20


MS (ES) m/z 586 (M+1).

-256-

5

Example 442

3-[2-(Isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[3-(morpholine-4-carbonyl)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid




MS (ES) m/z 580 (M+1).

10

Example 443


3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-propylcarbamoyl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



15 MS (ES) m/z 552 (M+1).

Example 444

3-[4-{2-[2-(4-Cyclobutylcarbamoyl-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



20


MS (ES) m/z 564 (M+1).

-257-

5

Example 445

3-[4-{2-[2-(4-Cyclohexylcarbamoyl-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




MS (ES) m/z 592 (M+1).

10

Example 446


3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-phenylcarbamoyl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



15 MS (ES) m/z 586 (M+1).

Example 447

3-[2-(Isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[4-(pyridin-3-ylcarbamoyl)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid



20


MS (ES) m/z 587 (M+1).

-258-

5

Example 448

3-[2-(Isopropoxycarbonylamino-methyl)-4-(2-{5-methyl-2-[4-(pyrrolidine-1-carbonyl)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid




MS (ES) m/z 564 (M+1).

10

Example 449

4-(4-{2-[4-(2-Carboxy-ethyl)-3-(isopropoxycarbonylamino-methyl)-phenoxy]-ethyl}-5-methyl-oxazol-2-yl)-benzoic acid



15

4-(4-{2-[4-(2-tert-Butoxycarbonyl-ethyl)-3-(isopropoxycarbonylamino-methyl)-phenoxy]-ethyl}-5-methyl-oxazol-2-yl)-benzoic acid (50 mg, 0.09 mmol, Example 436, Step B) was converted to the title compound using Standard Procedure C (40 mg, 95%): MS (ESI) m/z 525 (M+H)⁺.


20

-259-

5

Example 450


3-(4-{2-[4-(2-Carboxy-ethyl)-3-(isopropoxycarbonylamino-methyl)-phenoxy]-ethyl}-5-methyl-oxazol-2-yl)-benzoic acid:



10 The above compound is prepared by following a substantially similar procedure as described in Example 449. MS (ESI) m/z 525 ($M+H$)⁺.

Example 451

15 3-[4-{2-[2-(3-Cyclohexylcarbamoyl-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



A solution of 3-[4-{2-[2-(4-bromo-phenyl)-5-methoxy-oxazol-4-yl]-ethoxy}-2-(isopropoxy-carbonylamino-methyl)-propionic acid tert-butyl ester (230 mg, 0.38 mmol) in acetonitrile (20 ml) in a dry 3-neck flask was treated with 2-hydroxypyridine (44 mg, 0.46 mmol) and 1,1'-bis(diphenylphosphino)-ferrocene palladium (II) chloride (47 mg, 0.057 mmol). The mixture was stirred for about 5 minutes under N₂, and triethylamine (58 mg, 0.57 mmol) was added dropwise. CO gas was bubbled through the mixture, and the reaction was heated at 70°C for 4 h. The CO bubbling was replaced by a balloon filled with CO, and the reaction was stirred an additional 16 h. The mixture was cooled and partitioned equally into 2 flasks. One portion was treated with cyclohexylamine (46 mg, 0.46 mmol) and triethylamine (58 mg, 0.57 mmol). The mixture was stirred at 70°C for 16 h, cooled, filtered through Celite,


-260-

- 5 and concentrated to a brown solid (196 mg). The solid was purified by radial chromatography (10-70% EtOAc/hexanes) to give the penultimate *tert*-butyl ester (78 mg; MS (ESI) m/z 648.6 ($M+H$)⁺). The ester was converted to the title compound by Standard procedure D (72 mg, 64%). MS (ESI) m/z 592.0 ($M+H$)⁺.


10

Example 452

3-[4-(2-{2-[4-(3-Fluoro-benzoylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




- 15 Step A: 3-(2-(Isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-nitro-phenyl)-oxazol-4-yl]-ethoxy}-phenyl) -propionic acid *tert*-butyl ester



- 20 4-Methyl-3-nitro-benzenesulfonic acid 2-[5-methyl-2-(4-nitro-phenyl)-oxazol-4-yl]-ethyl ester (Preparation 7) and 3-[4-hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid *tert*-butyl ester were coupled by Standard procedure A to give the title compound.

-261-

- 5 Step B: 3-[4-{2-[2-(4-Amino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid *tert*-butyl ester



- A mixture of 3-(2-(isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(4-nitro-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid *tert*-butyl ester (207 mg, 0.365 mmol) and 10% Pd-C (27 mg) in EtOAc (20 mL) was stirred under H₂ (1 atm) for 18 h. The reaction mixture was filtered through Celite and concentrated (75°C) to give the title compound as a colorless oil (196 mg, 100%): MS (ESI) *m/z* 538 (M+H)⁺.

- 15 Step C: 3-[4-(2-{2-[4-(3-Fluoro-benzoylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid


- 3-Fluorobenzoyl chloride (31 mg, 0.20 mmol, 1.8 equiv) was added to a solution of 3-[4-{2-[2-(4-amino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonyl-amino-methyl)-phenyl]-propionic acid *tert*-butyl ester (58 mg, 0.11 mmol, 1 equiv) and triethylamine (30 µL, 22 mg, 0.22 mmol, 2.0 equiv) in CH₂Cl₂ (5 mL). After 16 h, the reaction solution was washed with 1 M aqueous HCl (5 mL) and saturated aq NaHCO₃ (5 mL). The organic layer was dried (MgSO₄) and concentrated (75 °C) to the intermediate ester as a light yellow oil. The oil was diluted with 4 M HCl in 1,4-dioxane (5 mL), stirred for 64 h, and concentrated to give the title compound (73 mg, 110%). HRMS Calculated for C₃₃H₃₅FN₃O₇: *m/z* 604.2459. Found: 604.2453.

-262-

5 The following Examples 453 to 475 are prepared by following a substantially similar procedure as described in Examples 451 and 452.

Example 453


10 3-[4-(2-{2-[4-(4-Fluoro-benzoylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



HRMS Calculated for $C_{33}H_{35}FN_3O_7$: m/z 604.2459. Found: 604.2454.

Example 454

15 3-[4-(2-{2-[4-(3,5-Difluoro-benzoylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



HRMS Calculated for $C_{33}H_{34}F_2N_3O_7$: m/z 622.2365. Found: 622.2352.


20

-263-

5

Example 455

3-[4-(2-{2-[4-(3,4-Difluoro-benzoylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




HRMS Calculated for $C_{33}H_{34}F_2N_3O_7$: m/z 622.2365. Found: 622.2382.

10

Example 456


3-(4-{2-[2-(4-Acetylamino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-[(pyridine-2-carbonyl)-amino]-methyl}-phenyl)-propionic acid



15 MS (ES) m/z 543 (M+1).

Example 457

3-(4-{2-[2-(4-Amino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-[(pyridine-2-carbonyl)-amino]-methyl}-phenyl)-propionic acid




20

MS (ES) m/z 501 (M+1).

5

Example 458

3-(4-{2-[2-(4-Isobutoxycarbonylamino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-
{[(pyridine-2-carbonyl)-amino]-methyl}-phenyl)-propionic acid




MS (ES) m/z 601 (M+1).

10

Example 459


3-[4-{2-[2-(4-Amino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-
(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



15 MS (ES) m/z 482 (M+1).

Example 460

3-[4-{2-[2-(4-Benzoylamino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-
(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



20


MS (ES) m/z 586 (M+1).

-265-

5

Example 461

3-(4-{2-[2-(4-Benzoylamino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-[[pyridine-2-carbonyl]-amino]-methyl}-phenyl)-propionic acid




MS (ES) m/z 661 (M+1).

10

Example 462


3-(4-(2-{2-[4-(4-Methoxy-benzoylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-[[pyridine-2-carbonyl]-amino]-methyl)-phenyl)-propionic acid



15 MS (ES) m/z 635 (M+1).

Example 463

3-(4-[2-(5-Methyl-2-{4-[(pyridine-3-carbonyl)-amino]-phenyl}-oxazol-4-yl)-ethoxy]-2-[[pyridine-2-carbonyl]-amino]-methyl)-phenyl)-propionic acid



20


MS (ES) m/z 606 (M+1).

-266-

5

Example 464

3-(4-[2-(2-{4-[(2,5-Dichloro-thiophene-3-carbonyl)-amino]-phenyl}-5-methyl-oxazol-4-yl)-ethoxy]-2-[[pyridine-2-carbonyl)-amino]-methyl]-phenyl)-propionic acid




MS (ES) m/z 680 (M+1).

10

Example 465


3-(4-(2-{2-[4-(*N,N*-Di-(butane-1-sulfonyl)amino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-[[pyridine-2-carbonyl)-amino]-methyl]-phenyl)-propionic acid



15 MS (ES) m/z 741 (M+1).

Example 466

3-(4-(2-{2-[4-(Butane-1-sulfonylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-[[pyridine-2-carbonyl)-amino]-methyl)-phenyl)-propionic acid



20


MS (ES) m/z 621 (M+1).

-267-

5

Example 467

3-[4-{2-[2-(4-Acetylamino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




MS (ES) m/z 524 (M+1).

10

Example 468


3-[4-{2-[2-(4-Butyrylamino-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



15 MS (ES) m/z 552 (M+1).

Example 469

3-[4-(2-{2-[4-(Cyclobutanecarbonyl-amino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




20

MS (ES) m/z 564 (M+1).

5

Example 470

3-[2-(Isopropoxycarbonylamino-methyl)-4-(2-{2-[4-(4-methoxy-benzoylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid




MS (ES) m/z 616 (M+1).

10

Example 471


3-[4-(2-{2-[4-(3,5-Dimethoxy-benzoylamino)-phenyl]-5-methyl-oxazol-4-yl}-ethoxy)-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



15 MS (ES) m/z 646 (M+1).

Example 472

3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-2-{4-[(pyridine-3-carbonyl)-amino]-phenyl}-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



20


MS (ES) m/z 587 (M+1).

-269-

5

Example 473

3-[4-[2-(2-{4-[(2,5-Dichloro-thiophene-3-carbonyl)-amino]-phenyl}-5-methyl-oxazol-4-yl)-ethoxy]-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid




MS (ES) m/z 661 (M+1).

10

Example 474


3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-2-{4-[(pyridine-2-carbonyl)-amino]-phenyl}-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



15 MS (ES) m/z 587 (M+1).

Example 475

3-[4-[2-(2-{4-[(Furan-2-carbonyl)-amino]-phenyl}-5-methyl-oxazol-4-yl)-ethoxy]-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid



20


MS (ES) m/z 576 (M+1).

-270-

5

Example 476

3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(pyrimidin-2-ylaminomethyl)-phenyl]-propionic acid




A solution of 3-{2-Aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester: acetic acid salt (0.40 mmol, 0.20 g, Example 1 Step B) in DMF (2 mL) was treated with K_2CO_3 (1.0 mmol, 0.14 g) and 2-chloropyrimidine (1.2 mmol, 0.14 g). The mixture was heated at 60 °C for 18 h, cooled, diluted with water (50 mL, and extracted with EtOAc (3 x 15 mL). The combined organic layers were dried (Na_2SO_4), concentrated and purified by silica gel chromatography to give the tert-butyl ester intermediate. This material was converted into the title compound (80 mg) using Standard procedure C. MS (ESI) m/z 459.2 ($M+H$)⁺.

20

Example 477

3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(2-methylsulfanyl-pyrimidin-4-ylamino)-methyl]-phenyl}-propionic acid



The above compound is prepared by following a substantially similar procedure as described in Example 476. MS (ESI) m/z 505.1 ($M+H$)⁺.


25

-271-

5

Example 478

3-{2-(Benzothiazol-2-ylaminomethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



A solution of 3-{2-aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid *tert*-butyl ester acetic acid salt (150 mg, 0.302 mmol; Example 1 Step B) and 2-chloro-benzothiazole (154 mg, 0.907 mmol) in toluene (5 mL) was treated with K_2CO_3 (42 mg). The suspension was heated at 110°C for 48 h and concentrated. The residue was purified by silica gel chromatography (25-50% EtOAc/hexanes) to afford the *tert*-butyl ester intermediate (15 mg). This material was converted into the title compound (80 mg) using Standard procedure C. The intermediate was treated with TFA (0.25 ml)/ CH_2Cl_2 (1.0 ml)/ H_2O (0.1 mL), stirred for 3 h, and concentrated. The residue was dissolved in CH_2Cl_2 (10 ml), washed with aqueous buffer (pH=7), dried (Na_2SO_4), and concentrated to afford the title compound (12 mg, 8%). MS (ESI) m/z 514.3 ($M+H$)⁺.


20

-272-

5

Example 479

3-{2-[(2-Benzoyl-phenylamino)-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid




A solution of 3-{2-aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester acetic acid salt (290 mg, 0.665 mmol, Example 1, Step B) and 2-benzoyl-cyclohexanone (161 mg, 0.797 mmol) in anisole (20 mL) was treated with a slurry of Pd/C (60 mg) in anisole (2 mL). The mixture was refluxed at 200°C with azeotropic removal of water for 2 h and cooled to room temperature. The catalyst was filtered and a fresh slurry of Pd/C (60 mg) in anisole was added. The mixture was heated at 110°C for 72 h and filtered through a pad of Celite. The filtrate was concentrated, purified by silica gel chromatography (25% EtOAc/hexanes) to afford a mixture of starting material and desired tert butyl ester intermediate. The mixture was treated with TFA (1.0 ml)/CH₂Cl₂(1.0 ml)/H₂O (0.1 mL), stirred for 3 h, and concentrated. The residue was and purified using silica gel chromatography (MeOH/EtOAc 1/9) to give the title compound (38 mg, 10%). MS (ESI) m/z 561.3 (M+H)⁺.

-273-

5

Example 480

3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(4-trifluoromethyl-phenylamino)-methyl]-phenyl}-propionic acid



10 A solution of 3-{2-aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester (174 mg, 0.40 mmol, Example 1 and Procedure I) in toluene (5.0 mL) in a sealed tube apparatus under a flow of N₂ was added Pd(OAc)₂ (15 mg), 2-(di-*t*-butylphosphino)biphenyl (10 mg), sodium *t*-butoxide (54 mg, 0.56 mmol) and 4-trifluoromethylchlorobenzene (29 mg, 0.16 mmol). The tube was sealed and heated at 110 °C for 14 h. The reaction mixture was cooled, quenched with
15 water (1.0 mL), and extracted with EtOAc (2 x 15 mL). The combined organics were concentrated and purified using silica gel chromatography column (10-50% EtOAc/hexanes) to yield the *tert*-butyl ester intermediate (80 mg). This material was converted into the title compound (40 mg, 48%) using Standard procedure C. MS (ESI) *m/z* 525.4 (M+H)⁺.


20

-274-

5 The following Examples 481 to 484 are prepared by following a substantially similar procedure as described in Examples 478 to 480.

Example 481


10 3-{2-[(4-Methanesulfonyl-phenylamino)-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



MS (ESI) m/z 535.1 (M+H)⁺.

Example 482

15 3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(4-propionyl-phenylamino)-methyl]-phenyl}-propionic acid



MS (ESI) m/z 513.2 (M+H)⁺.


20

-275-

5

Example 483

3-{2-{[Bis-(4-methoxy-phenyl)-amino]-methyl}-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid




MS (ESI) m/z 593.3 (M+H)⁺.

10

Example 484

3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(pyridin-2-ylaminomethyl)-phenyl]-propionic acid




15 MS (ESI) m/z 458.2 (M+H)⁺.


5

Example 485

3-{2-[[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl]-propionic acid




10 Step A: 3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(2,2,2-trifluoroacetyl-amino)-methyl]-phenyl}-propionic acid tert-butyl ester



3-{2-Aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester: acetic acid salt (894 mg, 1.80 mmol) was dissolved in CH₂Cl₂ (25 mL) and washed with saturated NaHCO₃ solution (15 mL). The organic layer was dried (NaSO₄), filtered, and concentrated to a yellow oil (681 mg). The crude amine was dissolved in CH₂Cl₂ (25 mL) and treated with trifluoroacetic anhydride (0.66 mL, 4.7 mmol) then pyridine (0.37 mL, 4.6 mmol). The reaction mixture was stirred at ambient temperature for 4 h and concentrated. The residue was partitioned between EtOAc and 1N HCl, and the organic layer was washed with saturated NaHCO₃ solution then brine. The organic layer was dried (Na₂SO₄) and concentrated to give a pale yellow solid (839 mg, 88%).


-277-

- 5 Step B: 3-(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[[methyl-(2,2,2-trifluoro-acetyl)-amino]-methyl]-phenyl)-propionic acid tert-butyl ester



- A solution of 3-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(2,2,2-trifluoro-acetyl-amino)-methyl]-phenyl}-propionic acid tert-butyl ester (151 mg, 0.28 mmol) in dry DMF (10 mL) was cooled in an ice bath and treated with NaH (22 mg, 0.55 mmol, 60% oil dispersion). After the reaction mixture was stirred for 25 min, iodomethane (0.15 mL, 3.0 mmol) was added, and the reaction was allowed to warm to ambient temperature gradually. After 4 h, more iodomethane (0.10 mL, 2 mmol) was added. The mixture was stirred overnight and was partitioned between EtOAc and aqueous LiCl solution. The organic layer was washed with brine, dried (Na₂SO₄), and concentrated. The residue was purified by silica gel radial chromatography (CH₂Cl₂/EtOAc 100/0 to 95/5) to give the title compound (111 mg, 73%).
- 15

- Step C: 3-{2-Methylaminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester
- 20




- A solution of 3-(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[[methyl-(2,2,2-trifluoro-acetyl)-amino]-methyl]-phenyl)-propionic acid tert-butyl ester (111 mg, 0.20 mmol) in methanol (5 mL) and THF (5 mL) was treated with 2N NaOH (1.0 mL, 2.0 mmol) and heated at 55°C for 1 h. The reaction mixture was cooled, concentrated, neutralized with 1N HCl and extracted into EtOAc. The organic layer was washed with saturated NaHCO₃ solution and brine, dried (Na₂SO₄), and concentrated.
- 25

-278-

- 5 The residue was purified by silica gel radial chromatography (CH₂Cl₂/methanol 95/5 to 90/10) to the title compound (37 mg, 42%). MS (ES) m/z 451.3 [M+1].

Step D: 3-{2-[[2-(5-Dichloro-thiophene-3-carbonyl)-methyl-amino]-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert butyl ester



10

A solution of 3-{2-methylaminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert-butyl ester (37 mg, 0.083 mmol) in CH₂Cl₂ (5 mL) was treated with triethylamine (0.035 mL, 0.25 mmol) then 2,5-dichloro-thiophene-3-carbonyl chloride (0.054 mL, 0.25 mmol) and stirred at ambient temperature overnight.

- 15 The mixture was diluted with EtOAc (25 mL) and washed with brine (3 x 10 mL). The organic layer was dried (Na₂SO₄) and concentrated to the title compound (50 mg, 96%). MS (ES) m/z 629, 631 [M+1].

- 20 **Step E:** 3-{2-[[2-(5-Dichloro-thiophene-3-carbonyl)-methyl-amino]-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid

- A solution of 3-{2-[[2-(5-dichloro-thiophene-3-carbonyl)-methyl-amino]-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid tert butyl ester (50 mg, 0.079 mmol) in CH₂Cl₂ (2 mL) was treated with anisole (1.0 mL) then TFA (0.6 mL). The solution was stirred at ambient temperature 2 h, and more TFA (1.0 mL) was added. After 15 min, the reaction was concentrated and co-evaporated with CCl₄ (3 x). The residue was triturated with hexanes to yield a foam (43 mg, 95%). ¹H NMR (400 MHz, CDCl₃) δ 2.36 (s, 3H), 2.48 (t, 0.6H, J = 7.8 Hz), 2.58 (t, 1.4H, J = 7.8 Hz), 2.72 (t, 0.6H, J = 7.8 Hz), 2.84 (s, 2H), 2.90 (t, 1.4H, J = 7.8 Hz), 2.93 (s, 1H), 2.98 (brs, 2H), 4.17 (t, 2H, J = 6.4 Hz), 4.47 (s, 0.6H), 4.71 (s, 1.4H), 6.61 (brs, 0.3H), 6.71-6.76 (m, 2H), 6.79 (s, 0.7H), 7.04 (d, 0.3H, J = 8.3 Hz), 7.08 (d, 0.7H, J = 8.3 Hz), 7.41-7.43 (m, 3H), 7.94 (brs, 2H).


30

-279-

- 5 The following Examples 486 to 488 are prepared by following a substantially similar procedure as described in Example 485.

Example 486


- 10 3-{2-[(Butyryl-methyl-amino)-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



MS (ES) m/z 465 (M+1).

Example 487


- 15 3-{2-[(Cyclobutanecarbonyl-methyl-amino)-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



MS (ES) m/z 477 (M+1).

Example 488

- 20 3-{2-[(Benzyloxycarbonyl-methyl-amino)-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid




MS (ES) m/z 529 (M+1).

-280-

Example 489

3-{4-[2-(2-Biphenyl-3-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-[(isopropoxycarbonyl-methyl-amino)-methyl]-phenyl}-propionic acid




3-{4-Hydroxy-2-[(isopropoxycarbonyl-methyl-amino)-methyl]-phenyl}-propionic acid tert-butyl ester (500 mg, 1.4 mmol; Preparation 22) and toluene-4-sulfonic acid 2-(2-biphenyl-3-yl-5-methyl-oxazol-4-yl)ethyl ester (617 mg, 1.4 mmol; Preparation 1/2) were coupled using Standard procedure A to give the penultimate *tert*-butyl ester. This ester was converted to the title compound using Standard procedure D (326 mg): MS (ESI) m/z 557 ($M+H$)⁺.

The following Examples 490 to 501 are prepared by following a substantially similar procedure as described in Example 489.

Example 490

3-{4-[2-[2-(3-Bromophenyl)-5-methyloxazol-4-yl]ethoxy]-2-[(isopropoxycarbonylmethylamino)methyl]phenyl} propionic acid




MS (ES) m/z 560 ($M+1$).

-281-

5

Example 491

3-{4-[2-(2-Biphenyl-3-yl-5-methyloxazol-4-yl)ethoxy]-2-[(isopropoxycarbonylmethylamino)methyl]phenyl} propionic acid




MS (ES) m/z 577 (M+1).

10

Example 492


3-{4-[2-(2-Cyclohexyl-5-methyloxazol-4-yl)ethoxy]-2-[(isopropoxycarbonylmethylamino)methyl]phenyl} propionic acid



15 MS (ES) m/z 487 (M+1).

Example 493

3-{2-[(Isopropoxycarbonylmethylamino)methyl]-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid



20


MS (ES) m/z 506 (M+1).

-282-

5

Example 494

3-{4-{2-[2-(4-Bromophenyl)-5-methyloxazol-4-yl]ethoxy}-2-[(isopropoxycarbonylmethylamino)methyl]phenyl} propionic acid




MS (ES) m/z 560 (M+1).

10

Example 495


3-{2-[(Isopropoxycarbonylmethylamino)methyl]-4-[2-(5-methyl-2-phenylthiazol-4-yl)ethoxy]phenyl} propionic acid



15 MS (ES) m/z 497 (M+1).

Example 496

3-{4-[2-(2-Biphenyl-4-yl-5-methylthiazol-4-yl)ethoxy]-2-[(isopropoxycarbonylmethylamino)methyl]phenyl} propionic acid



20


MS (ES) m/z 573 (M+1).

-283-

5

Example 497

3-(2-[(Isopropoxycarbonylmethylamino)methyl]-4-{2-[5-methyl-2-(1-methyl-cyclohexyl)oxazol-4-yl]ethoxy}phenyl) propionic acid




MS (ES) m/z 501 (M+1).

10

Example 498


3-{2-[(Isopropoxycarbonylmethylamino)methyl]-4-[2-(5-methyl-2-phenethyl-oxazol-4-yl)ethoxy]phenyl} propionic acid



15 MS (ES) m/z 509 (M+1).

Example 499

3-{2-[(Isopropoxycarbonylmethylamino)methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)ethoxy]phenyl} propionic acid



20


MS (ES) m/z 481 (M+1).

-284-

5

Example 500

3-(4-[2-(2-Biphenyl-3-yl)-5-methyloxazol-4-yl]ethoxy)-2- {[methyl(pyridine-2-carbonyl)amino]methyl} phenyl) propionic acid




MS (ES) m/z 576 (M+1).

10

Example 501

3-[4-{2-[2-(2-Bromophenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxy-carbonylaminomethyl)phenyl] propionic acid




15 MS (ES) m/z 546 (M+1).

-285-


5

Example 502

3-{4-[2-(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-[(ethyl-isopropoxycarbonyl-amino)-methyl]-phenyl}-propionic acid



10 Step A: 3-[4-[2-(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid methyl ester



15 3-[4-Hydroxy-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid methyl ester (Preparation 17) and toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)ethyl ester (Preparation 1) were combined according to Standard procedure A to give the title compound.

Step B: 3-{4-[2-(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-[(ethyl-isopropoxycarbonyl-amino)-methyl]-phenyl}-propionic acid

20 A solution of 3-[4-[2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-(isopropoxycarbonylamino-methyl)-phenyl]-propionic acid methyl ester (200 mg, 0.36 mmol) in DMF (15 mL) was treated with sodium bis(trimethylsilyl)amide (132 mg, 0.719 mmol). Ethyl iodide (112 mg, 0.719 mmol) was added, and the reaction mixture was stirred overnight at room temperature under N₂. The mixture was diluted with EtOAc (100 mL) and washed with brine (100mL), then water (100 mL). The organic layer was
25 dried (MgSO₄) and concentrated to the penultimate ester as white solid (179 mg). The material was dissolved in EtOH (20 mL), treated with 5N NaOH (20 mL), and stirred at


-286-

5 room temperature overnight. The mixture was acidified with 1N HCl(10mL) and extracted with EtOAc (50mL). The organics were dried (MgSO₄) and concentrated to give a white solid (201mg). MS [EI+] m/z 571 (M+H)⁺. Anal. Calculated for C₃₄H₃₈N₂O₆: C, 71.6; H, 6.7; N, 4.9. Found: C, 70.8; H, 6.8; N, 5.0.

10 The following Examples 503 to 504 are prepared by following a substantially similar procedure as described in Example 502.

Example 503


15 3-{4-[2-(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-[(isopropoxycarbonyl-methyl-amino)-methyl]-phenyl}-propionic acid



MS [EI+] m/z 557 (M+H)⁺. Anal. Calculated for C₃₃H₃₆N₂O₆: C, 71.2; H, 6.5; N, 5.0. Found: C, 70.6; H, 6.6; N, 5.1.

Example 504

20 3-{2-[(Isopropoxycarbonyl-methyl-amino)-methyl]-4-[2-(5-methyl-3-phenyl-pyrazol-1-yl)-ethoxy]-phenyl}-propionic acid




MS [EI+] m/z 480.1 (M+H)⁺.

-287-

5

Example 505

3-(2-[(Isopropoxycarbonyl-methyl-amino)-methyl]-4-{2-[5-methyl-2-(3-pyridin-3-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid




N-Methylation of carbamates: In a small screw cap vial was placed 3-
10 (2(isopropoxycarbonylamino-methyl)-4-{2-[5-methyl-2-(3-pyridin-3-yl-phenyl)-oxazol-
4-yl]-ethoxy}-phenyl)-propionic acid tert-butyl ester (0.5 mmol). A 1/1 mixture of
CHCl₃/TFA (1.6 mL) was added followed by 37% HCHO (50 μ L). The reaction mixture
was shaken at ambient temperature for 0.5 h and triethylsilane (110 μ L, 0.7 mmol) was
added. The mixture was shaken for another 0.5 h and was concentrated. The product
15 mixture was purified using mass guided reverse-phase HPLC to afford the title compound
(85%). MS [EI+] m/z 571 (M+H)⁺.

The following Examples 506 to 509 are prepared by following a
substantially similar procedure as described in Example 505.

20

Example 506

3-(2-[(Isopropoxycarbonyl-methyl-amino)-methyl]-4-{2-[5-methyl-2-(4-trifluoromethyl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid




25 MS (ES) m/z 549 (M+H)⁺.

-288-

5

Example 507

3-{4-{2-[2-(4-Butoxy-phenyl)-5-methyl-oxazol-4-yl]-ethoxy}-2-[(isopropoxycarbonyl-methyl-amino)-methyl]-phenyl}-propionic acid




MS (ES) m/z 553 (M+H)⁺.

10

Example 508


3-(2-[(Isopropoxycarbonyl-methyl-amino)-methyl]-4-{2-[5-methyl-2-(3-pyridin-3-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



15 MS (ES) m/z 558 (M+H)⁺.

Example 509

3-{2-[(Isopropoxycarbonyl-methyl-amino)-methyl]-4-[2-(5-methyl-2-pyridin-4-yl-thiazol-4-yl)-ethoxy]-phenyl}-propionic acid




20

MS (ES) m/z 498 (M+H)⁺.


5

Example 510

3-{3-[[[(2,5-Dichloro-thiophene-3-carbonyl)-amino]-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



Step A: 3-(3-Allyl-4-benzyloxy-phenyl)-propionic acid methyl ester




10


A solution of 3-(3-allyl-4-hydroxy-phenyl)-propionic acid methyl ester (20.5 g, 93.0 mmol; Brown GR, et al. *Bioorg. and Med. Chem. Lett.* 1997, 7, 597) in DMF (50 mL) was treated with Cs_2CO_3 (32.6 g, 100 mmol) then benzyl bromide (12.8 mL, 108 mmol) and heated at 55 °C for 16 h. Cs_2CO_3 (16.3 g, 50 mmol) and benzyl bromide (6.4 mL, 54 mmol) were added. The mixture was stirred at 55°C for 23 h, cooled, and partitioned between EtOAc (250 mL) and water (100 mL). The organic layer was washed with brine (75 mL). The combined aqueous layers were back-extracted with EtOAc (100 mL). The combined organic layers were dried (Na_2SO_4), and concentrated. The residue was purified by silica gel column chromatography (CH_2Cl_2 /hexanes 1/1 to 100/0) to the title compound (27.0 g, 93%).

20

-290-

5 Step B: 3-(4-Benzyloxy-3-carboxymethyl-phenyl)-propionic acid methyl ester

A solution of 3-(3-allyl-4-benzyloxy-phenyl)-propionic acid methyl ester (25.0 g, 80.5 mmol) in acetone (300 mL) and water (30 mL) was treated with 4-methylmorpholine 4-oxide (12.96 g, 95.8 mmol) then osmium (IV) oxide (5 chips). The flask
10 was covered with foil and stirred 20 h. The solution was diluted with EtOAc (1 L) and washed with 1N Na₂S₂O₃ (2 x 150 mL) and brine (125 mL). The organic layer was concentrated to a yellow oil (28.9 g). The oil was dissolved in THF (190 mL) and water (125 mL), and sodium periodate (49.0 g, 229 mmol) was added. THF (190 mL) and water (125 mL) were added. The thick white slurry was stirred for 2 h and filtered. The
15 filtrate was extracted with EtOAc (1 L). The organic layer was washed successively with brine, 1N Na₂S₂O₃, and brine (150 mL each) and was concentrated to an orange oil (25.3 g). The oil was diluted with *tert*-butanol (400 mL) and 2-methyl-2-butene (100 mL) and was cooled in an ice bath. The mixture was treated with sodium chlorite (68 g, 0.76 mol), and a solution of NaH₂PO₄ (68 g, 0.49 mol) in water (250 mL) was added over 5 min.
20 After 15 min, the ice bath was removed. The mixture was stirred for 2 h and was partitioned between EtOAc (1 L) and water (125 mL). The organic layer was washed with 1N Na₂S₂O₃ (125 mL) and brine (125 mL), dried (NaSO₄), and concentrated to the title compound as a tan solid (31.4 g, 119%). This material was used in subsequent reactions without further purification.



25


3-(4-Benzyloxy-3-carboxymethyl-phenyl)-propionic acid methyl ester (80.5 mmol), ammonium chloride (7.75 g, 145 mmol), EDC (27.7 g, 144 mmol), and *N*-hydroxybenzotriazole hydrate (19.6 g, 145 mmol) were combined in a flask and diluted

-291-

- 5 with DMF (320 mL). Ethyl-diisopropyl-amine (51 mL, 293 mmol) was added. The solution was stirred for 20 h and partitioned between EtOAc (1.2 L) and 1N HCl (250 mL). The organic layer was washed with saturated NaHCO₃ solution (200 mL) and brine (200 mL), dried (NaSO₄), and concentrated. The crude product was slurred with ethyl ether and filtered to give the title compound (15.9 g, 60%).

10

Step C: 3-[4-Benzyloxy-3-(*tert*-butoxycarbonylamino-methyl)-phenyl]-propionic acid methyl ester




- 3-(4-benzyloxy-3-carbamoylmethyl-phenyl)-propionic acid methyl ester
15 (15.9 g, 48.6 mmol) was dissolved in CH₃CN (850 mL) and DMF (8 mL) with warming. Water was added, the solution was cooled to ambient temperature, and [bis(trifluoroacetoxy)iodo]benzene (31.32 g, 72.8 mmol) was added. After 30 min, pyridine (8.2 mL) was added, and the solution was stirred for 17 h. Triethylamine (28 mL, 200 mmol) and di-*tert*-butyl dicarbonate (16.0 g, 73.3 mmol) were added. The
20 mixture was stirred for 2.5 h and was concentrated. The residue was partitioned between EtOAc (1 L) and brine (150 mL). The organic layer was washed with ice-cold 1N HCl (150 mL), saturated NaHCO₃ solution (150 mL) and brine (100 mL); dried (NaSO₄); and concentrated. The residue was purified by silica gel column chromatography (CH₂Cl₂/EtOAc 100/0 to 95/5) to a yellow oil (11.0 g, 57%).

25


-292-

- 5 Step D: 3-[3-(*tert*-Butoxycarbonylamino-methyl)-4-hydroxy-phenyl]-propionic acid methyl ester



- A solution of 3-[4-benzyloxy-3-(*tert*-butoxycarbonylamino-methyl)-phenyl]-propionic acid methyl ester (8.50 g, 21.3 mmol) in THF (100 mL) was treated
10 with 5% Pd-on-carbon (1.1 g) and shaken under a hydrogen atmosphere (60 psi) at ambient temperature for 6 h. The mixture was filtered through Celite and concentrated to a pale yellow solid (5.63 g, 85%).


- Step E: 3-{3-(*tert*-Butoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester
15



- 3-[3-(*tert*-Butoxycarbonylamino-methyl)-4-hydroxy-phenyl]-propionic acid methyl ester (5.50 g, 17.8 mmol), toluene-4-sulfonic acid 2-(5-methyl-2-phenyl-oxazol-4-yl)-ethyl ester (7.93 g, 22.2 mmol) and Cs₂CO₃ (7.23 g, 22.2 mmol) were
20 suspended in DMF (40 mL) and stirred at 55°C for 18 h. Additional toluene-4-sulfonic acid 2-(5-methyl-2-phenyl-oxazol-4-yl)-ethyl ester (3.00 g, 8.39 mmol) and Cs₂CO₃ (2.73 g, 8.39 mmol) were added and stirring was continued for 22 h. The reaction mixture was cooled and partitioned between EtOAc (125 mL) and water (50 mL). The organic layer was washed with brine (50 mL), dried (NaSO₄), and concentrated. The residue was
25 purified by silica gel column chromatography (CH₂Cl₂/EtOAc 97/3 to 80/20) to give pure product (3.33 g, 38%) and slightly impure product (7.08g).

-293-

- 5 Step F: 3-{3-Aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester: trifluoroacetic acid salt



- A solution of 3-{3-(*tert*-butoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester (3.33 g, 6.73 mmol) in
 10 CH₂Cl₂ (24 mL) was treated at ambient temperature with TFA (10 mL) and stirred for 3 h. The solution was concentrated and co-evaporated with CCl₄ (3x) to yield the title compound as a foam (4.13 g, quantitative)






- Step G: 3-{3-[(2,5-Dichloro-thiophene-3-carbonyl)-amino]-methyl}-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
 15

- General Parallel Synthesis Procedure: 3-{3-Aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester: trifluoroacetic acid salt (565 mg, 1.1 mmol) was dissolved in CH₂Cl₂ (25 mL) and washed with saturated NaHCO₃ solution (15 mL). The organic layer was dried (NaSO₄), filtered,
 20 and concentrated to 3-{3-aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester (368 mg, 0.93 mmol, 84%). A portion of this free base (26 mg, 0.071 mmol) in CH₂Cl₂ (0.5 mL) was treated with triethylamine (0.075 mL, 0.54 mmol) then 2,5-dichloro-thiophene-3-carbonyl chloride (0.049 mL, 0.22 mmol). The reaction mixture was shaken overnight and dimethylethylenediamine (0.15 mL, 1.4
 25 mmol) was added. The reaction mixture was shaken for 2 h and passed through an SCX column (1g, equilibrated with 2 mL MeOH then 2 mL MeOH/CH₂Cl₂ 1/1). The methyl ester-amide product was eluted with MeOH/CH₂Cl₂ (1:1, 10 mL) and concentrated. The residue was dissolved in THF (2 mL) and MeOH (2 mL) and treated with 5N NaOH (1 mL). The solution was heated at 55°C for 2.5 h, cooled, and acidified with 5N HCl (1.5
 30 mL). The mixture was transferred to a ChemElute cartridge and eluted with CH₂Cl₂. The solvent was removed under a stream of N₂. The crude product was dried under vacuum







- 5 and purified by mass-directed HPLC to give the title compound as a foam (9.9mg, 26%).
MS (ESI) m/z 559.1 ($M+H$)⁺.

Examples 511-543

- Examples 511 to 543 are prepared by following a substantially similar
10 procedure as described in Example 510.







No.	Compounds	MS (ES+)
511		541.3
512		463.3
513		559.1
514		513.3
515		499.3







-295-

No.	Compounds	MS (ES+)
516		485.3
517		515.3
518		481.3
519		493.3
520		486.2
521		487.2





No.	Compounds	MS (ES+)
522		475.2
523		535.2
524		536.2
525		499.2
526		515.2
527		477.3

-297-

No.	Compounds	MS (ES+)
528		463.3
529		543.2
530		505.3
531		525.2
532		491.2
533		547.3

No.	Compounds	MS (ES+)
534		521.3
535		505.3
536		547.3
537		505.3
538		505.3
539		491.3


-299-

No.	Compounds	MS (ES+)
540		505.3
541		525.2
542		521.2
543		527.1


-300-

Example 544

3-{2-(2-Isopropoxycarbonylamino-ethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



Step A: 5-Benzyloxy-2-bromo-(2-nitrovinyl)benzene




A solution of 2-bromo-5-hydroxy-benzaldehyde (5.99 g, 29.8 mmol, Preparation 13 Step A) in DMF (50 mL) was treated with benzyl bromide (7.65 g, 44.7 mmol) and Cs_2CO_3 (15.1 g, 44.7 mmol). The resulting mixture was heated at 80°C for 60 min and was quenched with water (200 mL). The mixture was extracted with EtOAc (2 x 100 mL), dried (Na_2SO_4), and concentrated to a residue, which was purified on silica gel chromatography (hexanes/EtOAc 9/1) to afford 5-benzyloxy-2-bromo-benzaldehyde as white solid in quantitative yield.

A solution of nitromethane (2.02 mL, 37.3 mmol) in ethanol (10 mL) was treated with 10N NaOH (3.0 mL) at ambient temperature. A white solid precipitated immediately, and a solution of 5-benzyloxy-2-bromo-benzaldehyde (8.67 g, 29.8 mmol) in CH_2Cl_2 (50 mL) was added. The reaction mixture was stirred for 18 h and quenched with water (100 mL). The mixture was extracted with CH_2Cl_2 (2 x 50 mL), dried (Na_2SO_4), and concentrated to a residue, which was then purified by silica gel chromatography (hexanes/EtOAc 8/2) to afford 1-(5-benzyloxy-2-bromo-phenyl)-2-nitro-ethanol as yellow oil.

-301-


5 This intermediate was dissolved in CH_2Cl_2 (150 mL) and cooled to 0°C . Methanesulfonyl chloride (2.55 mL, 60.0 mmol) was added, and the mixture was stirred for 15 min. Triethylamine (8.41 mL) was added, and the mixture was stirred for 60 min. The reaction was quenched with water (200 mL) and extracted with CH_2Cl_2 (150 mL). The organic layer was dried (Na_2SO_4) and concentrated. The residue was purified by
10 silica gel chromatography (hexanes/EtOAc 8/2) to afford the title compound (8.60 g, 86%). ^1H NMR (400 MHz, CDCl_3): δ 5.04 (s, 2 H), 6.93 (dd, 1H, $J = 2.7$ Hz, 9.0 Hz), 7.11 (d, 1H, $J = 3.2$ Hz), 7.34-7.37 (m, 5H), 7.48 (d, 1H, $J = 13.7$ Hz), 7.51 (d, 1H, $J = 9.0$ Hz), 8.28 (d, 1H, $J = 13.7$ Hz).

15 Step B: 2-(5-Benzyloxy-2-bromo-phenyl)-ethylamine




To a solution of 5-benzyloxy-2-bromo-(2-nitrovinyl)benzene (8.60 g, 25.6 mmol) in THF (100 mL) at -78°C was added LAH (4.1 g, 102 mmol) portion wise over 30 min. The reaction mixture was allowed to warm gradually to ambient temperature.
20 After 18 h, the mixture was carefully quenched with water (4 mL), 2N NaOH (4 mL), and water (12 mL) successively. The slurry was filtered and the precipitate was washed with CH_2Cl_2 (3 x 150 mL), dried (Na_2SO_4), and concentrated. The residue was purified by silica gel chromatography (EtOAc, then $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 10/1/0.01) to afford the title compound as an oil (3.36 g, 43%). ^1H NMR (400 MHz, CDCl_3): δ 2.84 (t, 2H, $J =$
25 6.7 Hz), 2.96 (t, 2H, $J = 6.7$ Hz), 5.03 (s, 2 H), 6.71 (dd, 1H, $J = 2.7$ Hz, 8.6 Hz), 6.86 (d, 1H, $J = 2.4$ Hz), 7.33-7.43 (m, 6H).

-302-

5 Step C: 2-(5-Benzyloxy-2-bromo-phenyl)-ethyl]-carbamic acid tert-butyl ester

A solution of 2-(5-benzyloxy-2-bromo-phenyl)-ethylamine (3.36 g, 10.97 mmol) in CH₂Cl₂ (50mL) at 0°C was treated with Et₃N (3.08 mL, 21.9 mmol) and di-*tert*-butyl dicarbonate (2.87 g, 13.2 mmol). The mixture was stirred for 2 h while being
10 allowed to warm gradually to ambient temperature and was concentrated. The residue was purified by silica gel chromatography (EtOAc/hexanes, 65:35) to afford the title compound as an oil (4.45 g, 100%). ¹H NMR (400 MHz, CDCl₃): δ 1.44 (s, 9H), 2.90 (t, 2H, *J* = 6.7 Hz), 3.39 (dt, 2H, *J* = 6.3 Hz, 6.7 Hz), 5.02 (s, 2 H), 6.72 (dd, 1H, *J* = 3.1 Hz, 8.6 Hz), 6.87 (br s, 1H), 7.33-7.43 (m, 6H).


15

Step D: 3-[4-Benzyloxy-2-(2-*tert*-butoxycarbonylamino-ethyl)-phenyl]-acrylic acid methyl ester

A solution of 2-(5-benzyloxy-2-bromo-phenyl)-ethyl]-carbamic acid *tert*-butyl ester (4.40 g, 10.8 mmol) in propionitrile (100 mL) was degassed (vacuum/ Ar
20 purge, 3x). Tri-*ortho*-tolylphosphine (0.660 g, 2.17 mmol), methyl acrylate (2.93 mL, 32.5 mmol), and diisopropylethyl amine (3.76 mL, 21.6 mmol) were added. The mixture was de-gassed (vacuum/ Ar purge, 3x). Pd(OAc)₂ (242 mg, 1.08 mmol) was added, and the reaction mixture was degassed again. The mixture was stirred at 95 °C for 18 h and
25 concentrated to a residue, which was purified by silica gel chromatography (EtOAc/hexanes, 7/3) to afford the title compound (4.15 g, 93%). ¹H NMR (400 MHz, CDCl₃) (major isomer): δ 1.43 (s, 9H), 2.94 (t, 2H, *J* = 6.8 Hz), 3.33 (dt, 2H, *J* = 6.3 Hz, 6.8 Hz), 3.79 (s, 3 H), 5.08 (s, 2 H), 6.28 (d, 1H, *J* = 15.6 Hz), 6.84 (br s, 1 H), 6.86 (d, 1H, *J* = 8.3 Hz), 7.34-7.43 (m, 5 H), 7.55 (d, 1H, *J* = 8.3 Hz), 7.91 (d, 1H, *J* = 15.6 Hz).


-303-

- 5 Step E: 3-[2-(2-tert-Butoxycarbonylamino-ethyl)-4-hydroxy-phenyl]-propionic acid methyl ester



To a solution of 3-[4-benzyloxy-2-(2-tert-butoxycarbonylamino-ethyl)-phenyl]-acrylic acid methyl ester (4.15 g, 10.1 mmol) in THF (100 mL) and MeOH (10 mL) was added 5% Pd-C (200 mg) in THF (10 mL). The resulting suspension was treated with hydrogen under balloon pressure for 18 h. The mixture was filtered through a pad of Celite and concentrated. The residue was purified by silica gel chromatography (EtOAc/hexanes, 6/4) to afford the title compound (1.20 g, 34%). ¹H NMR (400 MHz, CDCl₃): δ 1.44 (s, 9H), 2.57 (t, 2H, *J* = 7.8 Hz), 2.77 (t, 2H, *J* = 7.3 Hz), 2.88 (t, 2H, *J* = 7.8 Hz), 3.33 (br s, 2H), 3.68 (s, 3 H), 6.65 (br s, 1 H), 6.67 (d, 1H, *J* = 8.3 Hz), 7.02 (d, 1H, *J* = 8.3 Hz). MS (ES⁺) *m/z* 324.1 [M+H]⁺.

Step F: 3-{2-(2-tert-Butoxycarbonylamino-ethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester




20

To a solution of 3-[2-(2-tert-butoxycarbonylamino-ethyl)-4-hydroxy-phenyl]-propionic acid methyl ester (0.807 g, 2.50 mmol) and toluene-4-sulfonic acid 2-(5-methyl-2-phenyl-oxazol-4-yl)-ethyl ester (1.34 g, 3.74 mmol) in DMF (5 mL) was added Cs₂CO₃ (1.22 g, 3.74 mmol). The suspension was stirred at 65°C for 48 h, quenched with H₂O (200 mL), and extracted with EtOAc (3 x 100 mL). The combined organics were dried (Na₂SO₄) and concentrated to a residue, which was purified on silica gel chromatography (hexanes/EtOAc 9/1 to 8/2 to 6/4) to afford the title compound as a colorless oil (0.90 g, 71%). ¹H NMR (400 MHz, CDCl₃): δ 1.42 (s, 9H), 2.37 (s, 3H), 2.54 (t, 2H, *J* = 7.8 Hz), 2.77 (t, 2H, *J* = 7.8 Hz), 2.88 (t, 2H, *J* = 6.8 Hz), 2.96 (t, 2H, *J* =

- 5 6.8 Hz), 3.33 (~dt, 2H, $J = 6.8$ Hz), 3.65 (s, 3 H), 4.20 (t, 2H, $J = 6.8$ Hz), 6.70 (br s, 1 H), 6.72 (d, 1H, $J = 8.3$ Hz), 7.05 (d, 1H, $J = 8.3$ Hz), 7.39-7.44 (m, 3H), 7.95-7.98 (m, 2H).


MS (ES^+) m/z $[\text{M}+\text{H}]^+$.

- 10 Step G: 3-{2-(2-Amino-ethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester



- A solution of 3-{2-(2-tert-butoxycarbonylamino-ethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester (0.90 g, 1.77 mmol) in CH_2Cl_2 (10mL) at 0°C was treated with TFA (5.0 mL) and water (0.2 mL). The resulting solution was allowed to warm gradually to ambient temperature. After 18 h, the mixture was concentrated. The residue was partitioned between CH_2Cl_2 (30mL) and 5N NaOH (2 mL), and the aqueous layer was extracted with CH_2Cl_2 (3 x 30mL). The combined organics were dried (K_2CO_3) and concentrated to a residue, which was purified by silica gel chromatography (CH_2Cl_2 /MeOH/ NH_4OH , 8/2/0.05) to afford the title compound as a colorless oil (0.42 g, 58%). ^1H NMR (400 MHz, CDCl_3): δ 2.34 (s, 3H), 2.52 (t, 2H, $J = 7.8$ Hz), 2.81 (t, 2H, $J = 7.8$ Hz), 2.91 (t, 4H, $J = 6.4$ Hz), 3.06 (t, 2H, $J = 7.3$ Hz), 3.60 (s, 3 H), 4.20 (t, 2H, $J = 6.8$ Hz), 6.68 (br s, 1 H), 6.71 (d, 1H, $J = 8.3$ Hz), 7.01 (d, 1H, $J = 8.3$ Hz), 7.37-7.39 (m, 3H), 7.89-7.92 (m, 2H).

- 25 Step H: 3-{2-(2-Isopropoxycarbonylamino-ethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



A solution of 3-{2-(2-amino-ethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid methyl ester (26 mg, 0.064 mmol) in CH_2Cl_2 (2 mL) at 0°C was treated with TEA (0.2 mL) and isopropyl chloroformate (0.13 mL, 1.0 M in

-305-

5 ether). The resulting mixture was allowed to warm gradually to ambient temperature. After 2 h, the mixture was concentrated to a residue, which was purified by silica gel chromatography (hexanes/EtOAc 7/3) to afford the isopropyl carbamate as a colorless oil. ¹H NMR (400 MHz, CDCl₃): δ 1.19 (d, 6H, *J* = 5.9 Hz), 2.42 (s, 3H), 2.54 (t, 2H, *J* = 7.8 Hz), 2.79 (t, 2H, *J* = 6.8 Hz), 2.89 (t, 2H, *J* = 7.8 Hz), 3.05 (br s, 2H), 3.38 (br s, 2H),
10 3.60 (s, 3 H), 4.26 (t, 2H, *J* = 6.8 Hz), 4.68 (br s, 1H), 4.89 (q, 1H, *J* = 6.4 Hz), 6.70 (br s, 1 H), 6.71 (d, 1H, *J* = 8.3 Hz), 7.05 (d, 1H, *J* = 8.3 Hz), 7.47 (m, 3H), 8.10 (m, 2H).


This compound was dissolved in THF (1.0 mL) and MeOH (0.5 mL). The solution was treated with 2N NaOH (0.32 mL) and heated at 60 °C for 60 min. The reaction mixture was concentrated, neutralized with 2N HCl (0.40 mL), and extracted
15 with EtOAc (3 x 5 mL). The combined organics were dried (Na₂SO₄), and concentrated to a residue, which was purified by silica gel chromatography (hexanes/EtOAc 9/1, then EtOAc/MeOH 95/5) to afford the title compound as a white solid (12 mg, 32% over two steps). ¹H NMR (400 MHz, CDCl₃): δ 1.17 (d, 6H, *J* = 5.9 Hz), 2.36 (s, 3H), 2.52 (br s, 2H), 2.79 (t, 2H, *J* = 6.8 Hz), 2.89 (t, 2H, *J* = 7.8 Hz), 2.93 (t, 2H, *J* = 6.8 Hz), 3.31 (br s, 2H), 4.16 (t, 2H, *J* = 6.8 Hz), 4.68 (br s, 1H), 4.89 (q, 1H, *J* = 6.4 Hz), 6.67 (br s, 1 H),
20 6.71 (d, 1H, *J* = 8.3 Hz), 7.05 (d, 1H, *J* = 8.3 Hz), 7.47 (m, 3H), 8.10 (m, 2H). MS [ES] *m/z* 481.4 (M+1).

-306-

5 The following Examples 545 to 552 are prepared by following a substantially similar procedure as described in Example 544.

10 Example 545


3-{2-[2-(Butane-1-sulfonylamino)-ethyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



MS [ES] m/z 515.3 (M+1).

15 Example 546

3-(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-{2-[(pyridine-2-carbonyl)-amino]-ethyl}-phenyl)-propionic acid



MS [ES] m/z 500.3 (M+1).


20

-307-

5

Example 547

3-{2-{2-[(2,5-Dichloro-thiophene-3-carbonyl)-amino]-ethyl}-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid




MS [ES-] m/z 573.0(M-1).

10

Example 548

3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(2-phenylacetyl-amino-ethyl)-phenyl]-propionic acid




15 MS [ES] m/z 513.2 (M+1).

-308-

5

Example 549

3-{2-[2-(Cyclobutanecarbonyl-amino)-ethyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid




MS [ES] m/z 477.1 (M+1).

10

Example 550

3-{2-(2-Benzoylamino-ethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid




15 MS [ES] m/z 499.3 (M+1).

-309-

5

Example 551

3-{2-(2-Isobutoxycarbonylamino-ethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid




MS [ES] m/z 495.3 (M+1).

10

Example 552

3-{2-(2-Benzoyloxycarbonylamino-ethyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid




15 MS [ES] m/z 529.0 (M+1).

-310-


5

Example 553

3-(2-(2-Isopropoxycarbonylamino-ethyl)-4-{2-[5-methyl-2-(4-morpholin-4-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid



10 Step A: 3-[4-Hydroxy-2-(2-isopropoxycarbonylamino-ethyl)-phenyl]-propionic acid methyl ester



15 A solution of 3-[2-(2-tert-butoxycarbonylamino-ethyl)-4-hydroxy-phenyl]-propionic acid methyl ester (282 mg, 0.876 mmol; Example 544, Step E) in CH₂Cl₂ (10 mL) at ambient temperature was treated with TFA (5.0 mL), stirred for 60 min, and concentrated. The residue in CH₂Cl₂ (10 mL) was treated with triethyl amine (2.0 mL) and iso-propyl chloroformate (0.97 mL, 1.0 M in toluene). The reaction mixture was stirred at ambient temperature for 16 h and concentrated. The crude material was purified using silica gel chromatography (50% EtOAc/hexanes) to yield the title compound (180 mg, 67%).

20

Step B: 3-(2-(2-Isopropoxycarbonylamino-ethyl)-4-{2-[5-methyl-2-(4-morpholin-4-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid

25 A solution of toluene-4-sulfonic acid 2-[5-methyl-2-(4-morpholin-4-yl-phenyl)-oxazol-4-yl]-ethyl ester (44 mg, 0.10 mmol; Preparation 5) and 3-[4-hydroxy-2-(2-isopropoxycarbonylamino-ethyl)-phenyl]-propionic acid methyl ester (31 mg, 0.10 mmol) in DMF (1.0 mL) was treated with K₂CO₃ (30 mg). The resulting suspension was stirred at 65 °C for 16 h and diluted with water (10 mL). The mixture was extracted with EtOAc (3 x 10 mL). The combined organic layers were dried (Na₂SO₄), concentrated,

-311-

- 5 and purified using silica gel chromatography (50% EtOAc/hexanes to yield 3-(2-(2-isopropoxycarbonylamino-ethyl)-4-{2-[5-methyl-2-(4-morpholin-4-yl-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid methyl ester.


This ester intermediate was dissolved in THF (0.6 mL) and MeOH (0.4 mL) and treated with aqueous 2M LiOH (1.0 mL, 2.0 mmol). The mixture was stirred for
10 16 h at ambient temperature, neutralized with HCl (1.0 mL, 2.0 M), and concentrated. The residue was extracted with EtOAc (3 x 5 mL). The combined organic layers were dried (Na₂SO₄), concentrated, and purified using silica gel chromatography column (hexanes/EtOAc/HOAc, 5/5/0.02) to afford the title compound (10 mg, 18%). MS (ES+) m/z 566.2 (M+H)⁺.

15

The following Examples 554 to 560 are prepared by following a substantially similar procedure as described in Example 553.

Example 554

- 20 3-[4-[2-(2-Biphenyl-3-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-(2-isopropoxycarbonyl-amino-ethyl)-phenyl]-propionic acid



MS [ES] m/z 557.5 (M+1).


25

-312-

5

Example 555

3-[4-[2-(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-(2-isopropoxycarbonyl-amino-ethyl)-phenyl]-propionic acid




MS [ES] m/z 557.2 (M+1).

10

Example 556

3-{2-(2-Isopropoxycarbonyl-amino-ethyl)-4-[2-(5-methyl-2-morpholin-4-yl-thiazol-4-yl)-ethoxy]-phenyl}-propionic acid




15 MS [ES] m/z 506.2 (M+1).

-313-

5

Example 557

3-{2-(2-Isopropoxycarbonylamino-ethyl)-4-[2-(5-methyl-2-pyridin-2-yl-thiazol-4-yl)-ethoxy]-phenyl}-propionic acid




MS [ES] m/z 498.3 (M+1).

10

Example 558

3-{2-(2-Isopropoxycarbonylamino-ethyl)-4-[2-(5-methyl-3-phenyl-pyrazol-1-yl)-ethoxy]-phenyl}-propionic acid




15 MS [ES] m/z 480.3 (M+1).

-314-

5

Example 559

3-(2-(2-Isopropoxycarbonylamino-ethyl)-4-{2-[5-methyl-2-(4-phenylamino-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid




MS [ES] m/z 572.2 (M+1).

10

Example 560

3-[2-(2-Isopropoxycarbonylamino-ethyl)-4-(2-{5-methyl-2-[4-(methyl-phenyl-amino)-phenyl]-oxazol-4-yl }-ethoxy)-phenyl]-propionic acid




15 MS [ES] m/z 586.2 (M+1).

-315-


5

Example 561

2-{2-(tert-Butoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid



Step A: 2-(2-Allyl-4-benzyloxy-phenoxy)-2-methyl-propionic acid ethyl ester




10

2-Allyl-4-benzyloxy-phenol (WO 9728137 A1 19970807, Adams et al.) (6.04 g, 25.1 mmol) in dry DMF (75 mL) was cooled in an ice bath and treated with NaH (1.78 g, 44.5 mmol, 60 % oil dispersion). After 30 min, the red reaction mixture was treated over 5 min with ethyl bromoisobutyrate (7.4 mL, 50 mmol). The cold bath was removed after 5 min. The reaction was stirred 20 min and placed in an oil bath (T=85°C). After 18 h, the mixture was cooled and partitioned between brine (75 mL) and ether (200 mL). The organic layer was washed with brine (50 mL), dried (Na₂SO₄), and concentrated to a brown oil (12g). The crude product was purified by flash chromatography using hexanes:ethyl acetate (100:0 to 5:1) to give the desired product (8.47 g, 95%).

20

Step B: 2-(4-Benzyloxy-2-carboxymethyl-phenoxy)-2-methyl-propionic acid ethyl ester




A solution of 2-(2-allyl-4-benzyloxy-phenoxy)-2-methyl-propionic acid ethyl ester (8.07 g, 22.8 mmol) in acetone (85 mL) and water (8.5 mL) was treated with 4-methyl-morpholine 4-oxide (3.68 g, 27.2 mmol) then osmium (IV) oxide (2 chips). The flask was covered with foil and stirred 5 h. The solution was diluted with EtOAc (600

25

-316-

5 mL) and washed with 1N $\text{Na}_2\text{S}_2\text{O}_3$ (2 x 75 mL) and brine (75 mL). The organic layer was concentrated to a tan oil (8.74 g). The oil was dissolved in THF (54 mL) and water (36 mL), and sodium periodate (15.4 g, 72.0 mmol) was added. THF (54 mL) and water (36 mL) were added. The thick white slurry was stirred for 3 h and filtered. The filtrate was extracted with EtOAc (500 mL). The organic layer was washed successively with brine, 10 1N $\text{Na}_2\text{S}_2\text{O}_3$, and brine (75 mL each) and was concentrated to an orange oil (7.65 g). The oil was diluted with *tert*-butanol (180 mL) and 2-methyl-2-butene (60 mL) and was cooled in an ice bath. The mixture was treated with sodium chlorite (19 g, 0.21 mol), and a solution of NaH_2PO_4 (19 g, 0.14 mol) in water (120 mL) was added over 5 min. After 15 min, the ice bath was removed. The mixture was stirred for 2 h and was partitioned 15 between EtOAc (500 mL) and water (50 mL). The organic layer was washed with 1N $\text{Na}_2\text{S}_2\text{O}_3$ (75 mL) and brine (75 mL), dried (NaSO_4), and concentrated to the title compound as a pale yellow solid (6.52 g, 77%). This material was used in subsequent reactions without further purification.


20 Step C: 2-(4-Benzyloxy-2-carboxymethyl-phenoxy)-2-methyl-propionic acid ethyl ester



2-(4-Benzyloxy-2-carboxymethyl-phenoxy)-2-methyl-propionic acid ethyl ester (500 mg, 1.34 mmol), ammonium chloride (108 mg, 2.02 mmol), EDC (3.86 mg, 2.01 mmol), and *N*-hydroxybenzotriazole hydrate (272 mg, 2.01 mmol) were combined in a flask and diluted with DMF (5 mL). Ethyl diisopropyl amine (0.70 mL, 4.0 mmol) was added. The solution was stirred for 18 h and partitioned between EtOAc (25 mL) and 1N HCl (10 mL). The organic layer was washed with saturated NaHCO_3 solution (10 mL) then brine (10 mL), dried (Na_2SO_4), and concentrated. The crude product was purified by 30 radial chromatography (hexanes/EtOAc 1/2 to 1/4) to give the title compound (430 mg, 86%).


-317-

- 5 Step D: 2-[4-Benzyloxy-2-(tert-butoxycarbonylamino-methyl)-phenoxy]-2-methyl-propionic acid ethyl ester



- 2-(4-Benzyloxy-2-carbamoylmethyl-phenoxy)-2-methyl-propionic acid ethyl ester (430 mg, 1.16 mmol) was dissolved in CH₃CN (4.5 mL) and water (4.5 mL), and [bis(trifluoroacetoxy)iodo]benzene (745 mg, 1.73 mmol) was added. After 10 min, pyridine (0.19 mL, 2.3 mmol) was added, and the solution was stirred for 17 h. Triethylamine (0.60 mL, 4.3 mmol) and di-tert-butyl dicarbonate (506 mg, 2.32 mmol) were added. The mixture was stirred for 1 h and was concentrated. The residue was partitioned between ether (50 mL) and brine (15 mL). The organic layer was washed with ice-cold 1N HCl, saturated NaHCO₃ solution, and brine (15 mL each); dried (NaSO₄); and concentrated. The residue was purified by silica gel column chromatography (hexanes/EtOAc 4/1 to 3/2) to a yellow oil (345 g, 67%).

- 20 Step E: 2-[2-(tert-Butoxycarbonylamino-methyl)-4-hydroxy-phenoxy]-2-methyl-propionic acid ethyl ester



- A solution of 2-[4-benzyloxy-2-(tert-butoxycarbonylamino-methyl)-phenoxy]-2-methyl-propionic acid ethyl ester (0.38 g, 0.86 mmol) in THF (15 mL) was treated with 5% Pd-on-carbon (47 mg) and shaken under a hydrogen atmosphere (60 psi) at ambient temperature for 18 h. The mixture was filtered through Celite and concentrated. The crude product was purified by radial chromatography (hexanes/EtOAc 2.5/1) to give the title compound (208 mg g, 69%).


-318-

- 5 Step F: 2-{2-(tert-Butoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid

2-[2-(tert-Butoxycarbonylamino-methyl)-4-hydroxy-phenoxy]-2-methyl-propionic acid ethyl ester (200 mg, 0.566 mmol), toluene-4-sulfonic acid 2-(5-methyl-2-phenyl-oxazol-4-yl)-ethyl ester (245 mg, 686 mmol) and Cs_2CO_3 (185 mg, 0.567 mmol)
 10 were suspended in DMF (5 mL) and stirred at 55°C for 39 h. The reaction mixture was cooled and partitioned between Et_2O (50 mL) and water (10 mL). The organic layer was washed with brine (10 mL), dried (NaSO_4), and concentrated. The residue was purified by radial chromatography (hexanes/ EtOAc 4/1) to give an oil (280 mg). A solution of the product in MeOH (6 mL) and THF (3 mL) was treated with 2.5N aqueous NaOH solution
 15 (2 mL) and heated at 55 °C for 2 h. The mixture was cooled and concentrated. Ice chips were added (~5) and the mixture was acidified using 5N aqueous HCL solution (2 mL). The reaction mixture was partitioned between EtOAc (30 mL) and brine (15 mL). The organic layer was washed with brine (2 x 10 mL), dried (NaSO_4), and concentrated. The residue was purified by silica gel column chromatography (hexanes/ EtOAc/HOAc 3/2/0
 20 to 3/2/0.05) to give the title compound as a white foam (156 mg, 54%). MS (ES^-) m/z 509.1 $[\text{M}-\text{H}]^-$. Anal. Calcd. For $\text{C}_{28}\text{H}_{34}\text{N}_2\text{O}_7$: C: 65.88; H: 6.71; N: 5.49. Found: C: 65.67; H: 6.79; N: 5.52.

Example 562

- 25 {2-(tert-Butoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-acetic acid



The above compound is prepared by following a substantially similar procedure as described in Example 561. MS [ES] m/z 483.2 ($\text{M}+1$).


30

-319-


5

Example 563

2-{2-(Ethoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid



10 Step A: 2-{2-Aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid ethyl ester: trifluoroacetic acid salt



15 A solution of 2-{2-(tert-butoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid ethyl ester (350 mg, 0.650 mmol) in CH₂Cl₂ (10 mL) was treated at ambient temperature with TFA (4 mL) and stirred for 2.5 h. The solution was concentrated and co-evaporated with CCl₄ (3x) to yield the title compound as a foam (381 mg, quantitative).

Step B: 2-{2-(Ethoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid

20 General Parallel Synthesis Procedure: 2-{2-Aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid ethyl ester: trifluoroacetic acid salt (381 mg, 0.650 mmol max.) was dissolved in CH₂Cl₂ (25 mL) and washed with saturated NaHCO₃ solution (15 mL). The organic layer was dried (NaSO₄), filtered, and concentrated to 2-{2-aminomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid ethyl ester (245 mg, 0.59 mmol, 86%). A portion of this free base (13 mg, 0.030 mmol) in CH₂Cl₂ (0.8 mL) was treated with triethylamine (0.050 mL, 0.36 mmol) then ethyl chloroformate (0.020 mL, 0.21 mmol). The reaction mixture was shaken overnight and dimethylethylenediamine (0.050 mL,

-320-


5 0.47 mmol) was added. The reaction mixture was shaken for 2 h, diluted with MeOH (0.5 mL), and passed through an SCX column (1g, equilibrated with 4 mL MeOH/CH₂Cl₂ 1/1). The methyl ester-amide product was eluted with MeOH/CH₂Cl₂ (1:1, 8 mL) and concentrated. The residue was dissolved in THF (1 mL) and EtOH (1 mL) and treated with 2N NaOH (0.5 mL). The solution was heated at 45°C for 45 min, cooled, 10 concentrated, and acidified with 1N HCl (2.5 mL). The mixture was diluted with diluted with CH₂Cl₂ (0.5 mL), transferred to a 3-mL ChemElute cartridge, and eluted with CH₂Cl₂ (8 mL). The solvent was removed under a stream of N₂. The crude product was dried under vacuum and purified by mass-directed HPLC to give a foam (2.7 mg, 19%). MS (ES+) m/z 483.5 [M+H]⁺.

15

The following Examples 564 to 570 are prepared by following a substantially similar procedure as described in Example 563.

Example 564

20 {2-(Benzyloxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-acetic acid



MS [ES] m/z 517.2 (M+1).


25

-321-

5

Example 565

{2-[[2-(5-Dichloro-thiophene-3-carbonyl)-amino]-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-acetic acid




MS [ES] m/z 561.1 (M+1).

10

Example 566


{2-[(Cyclobutanecarbonyl)-amino]-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-acetic acid



15 MS [ES] m/z 465.2 (M+1).

Example 567

2-{2-(Butyrylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid



20


MS [ES] 481 m/z (M+1).

-322-

5

Example 568

2-{2-[[2,5-Dichloro-thiophene-3-carbonyl]-amino]-methyl}-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid




MS [ES] m/z 481 (M+1).

10

Example 569


2-{2-(Ethoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid



15 MS [ES] m/z 483 (M+1).

Example 570

2-{2-[(Cyclobutanecarbonyl-amino)-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid



20



MS [ES] m/z 493 (M+1).

-323-

5

Example 571


3-(2-Cyano-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid

Step A: 2-Bromo-5-hydroxy-benzonitrile

10

To a stirred solution of 3-cyanophenol (2.00 g, 18.0 mmol) in anhydrous acetonitrile (20 mL) under N₂ at -30°C was added dropwise 54% HBF₄.Et₂O (2.48 mL, 18.0 mmol). The temperature was maintained below -20° C during addition. To this stirred solution was added NBS (3.27 g, 18.0 mmol) portion wise keeping the temperature below -10°C. After the addition was complete, the solution was allowed to warm to 19°C. The reaction mixture was diluted with 38% NaHSO₃ (10 mL) and extracted with MTBE (2 x 25 mL). The organic layer was washed with H₂O (2 x 25 mL) and brine (25 mL), dried (MgSO₄), filtered, and concentrated to give a white solid (3.15 g). Purification of a portion of this product (1.0 g) by radial chromatography (30% EtOAc/hexane) gave 0.51g of the title compound as a white solid. mp 183-184°C. Anal. Calculated for C₇H₄BrNO: C, 42.46; H, 2.04; N, 7.07. Found: C, 42.44; H, 1.93; N, 6.90.

20

Step B: 2-(3-Ethoxy-buta-1,3-dienyl)-5-hydroxy-benzonitrile


25

To a stirred solution of 2-bromo-5-hydroxy-benzonitrile (1.4 g, 7.1 mmol), ethyl acrylate (2.12g, 21.2 mmol), palladium(II) acetate (0.159g, 7.0 mmol), diisopropylethylamine (0.82 g, 14.1 mmol), and tri-*ortho*-tolylphosphine (0.430g, 14.1 mmol) was added propionitrile (50 mL). The reaction was stirred overnight at 80°C under a stream of N₂. The mixture was cooled, filtered, and concentrated. The crude

-324-

- 5 material (3.0g) was purified by radial chromatography (10-70% EtOAc/hexanes) to give the title compound as a white solid (1.1 g). MS [EI] m/z 216 (M+H)⁺.

Step C: 2-(3-Ethoxy-but-3-enyl)-5-hydroxy-benzonitrile



- 10 To a solution of 2-(3-ethoxy-buta-1,3-dienyl)-5-hydroxy-benzonitrile (600mg) in EtOAc (50mL) was added 5% Pd/C (0.3g). The reaction was exposed to hydrogen gas in a Parr apparatus at 60psi overnight at ambient temperature. The reaction mixture was filtered and concentrated to give the title compound (410 mg). MS [EI] m/z 218 (M+H)⁺.

15

Step D: 3-(2-Cyano-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid

- A mixture of 2-(3-ethoxy-but-3-enyl)-5-hydroxy-benzonitrile (220 mg, 1.00 mmol), toluene-4-sulfonic acid 2-[5-methyl-2-(4-phenoxy-phenyl)-oxazol-4-yl]-ethyl ester (450 mg, 1.00 mmol; Preparation 6), and CsCO₃ (978 mg, 3.00 mmol) in DMF (10mL) was stirred overnight at 60°C under an N₂. The mixture was cooled, diluted with EtOAc (20 mL) and washed with brine (20 mL). The organic layer was dried (Na₂SO₄) and concentrated. The product mixture was purified by radial chromatography (10-70% EtOAc/hexanes) to give 3-(2-cyano-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid ethyl ester (201 mg). A solution of this ester in EtOH (10 mL) and 5N NaOH (10 mL) was heated at 70°C overnight, concentrated, acidified with 1N HCl, and extracted with EtOAc. The organic layer dried (Na₂SO₄) and concentrated to give the title compound (192 mg). MS [ES] m/z 469 (M+1).


30

-325-

5 The following Examples 572 to 573 and 575 to 581 are prepared by following a substantially similar procedure as described in Example 571.

Example 572

3-{4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-cyanophenyl} propionic acid:




10

MS [ES] m/z 453 (M+1).

Example 573

3-{4-[3-(Biphenyl-4-yloxy)propoxy]-2-cyanophenyl} propionic acid




15

MS [ES] m/z 402 (M+1).

Example 575

3-(4-{2-[2-(4-Bromo-phenyl)-5-methyloxazol-4-yl]ethoxy}-2-cyanophenyl) propionic acid

20




MS [ES] m/z 456 (M+1).

-326-

5

Example 576

3-{4-[2-(2-Biphenyl-3-yl-5-methyloxazol-4-yl)ethoxy]-2-cyanophenyl} propionic acid




MS [ES] m/z 453 (M+1).

10

Example 577

3-{2-Cyano-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid




MS [ES] m/z 402 (M+1).

15

Example 578

3-{2-Cyano-4-[2-(5-methyl-3-phenylpyrazol-1-yl)ethoxy]phenyl} propionic acid:




MS [ES] m/z 376 (M+1).

20

Example 579

3-(4-{2-[2-(4-Butoxyphenyl)-5-methyloxazol-4-yl]ethoxy}-2-cyanophenyl) propionic acid




MS [ES] m/z 449 (M+1).

-327-

5

Example 580

3-(2-Cyano-4-{2-[5-methyl-2-(6-phenylpyridin-3-yl)thiazol-4-yl]ethoxy}phenyl) propionic acid




MS [ES] m/z 470 (M+1).

10

Example 581

3-{2-Cyano-4-[2-(5-methyl-2-phenylthiazol-4-yl)ethoxy]phenyl} propionic acid




MS [ES] m/z 393 (M+1).


15

Example 582

3-{2-Benzylcarbamoyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid



20 Step A: 3-(4-Benzyloxy-2-formyl-phenyl)-acrylic acid *tert*-butyl ester




To a solution of 3-(2-formyl-4-hydroxy-phenyl)-acrylic acid *tert*-butyl ester (4.97 g, 20.0 mmol) in DMF (40 mL) was added benzyl bromide (2.85 mL, 24.0

-328-

5 mmol) and Cs_2CO_3 (7.82 g, 24.0 mmol). The suspension was stirred at 80°C for 30 min, and TLC indicated the reaction was complete. The mixture was quenched with H_2O (500 mL) and extracted with EtOAc (2 x 100 mL). The combined organics were dried (Na_2SO_4) and concentrated to a residue, which was purified by silica gel chromatography (hexanes/EtOAc 9/1) to afford the title compound as yellow solid (6.70 g, 99%).

10

Step B: 5-Benzyloxy-2-(2-tert-butoxycarbonyl-vinyl)-benzoic acid




A solution of 3-(4-benzyloxy-2-formyl-phenyl)-acrylic acid tert-butyl ester (6.70 g, 19.8 mmol) in *tert*-butanol (150 mL) was treated with 2-methyl-2-butene (50 mL) and cooled to 0°C . A solution of NaClO_2 (17.0 g, 188 mmol) and NaH_2PO_4 (17.0 g, 142 mmol) in H_2O (200 mL) was added, and the mixture was stirred at 0°C for 15 min. The ice bath was removed, and the mixture was stirred at room temperature for 2 h. The organic layer was separated, and the aqueous layer was extracted with EtOAc (2 x 100 mL). The combined organic layers were dried (Na_2SO_4) and concentrated. The residue was purified using silica gel chromatography (hexanes/EtOAc 9/1 to 0/10) to afford the title compound as white solid (6.78 g, 96%). ^1H NMR (400 MHz, CDCl_3): δ 1.52 (s, 9 H), 5.12 (s, 2 H), 6.20 (d, 1H, $J = 16.2$ Hz), 7.13 (dd, 1H, $J = 3.6, 9.0$ Hz), 7.34-7.45 (m, 5H), 7.58 (d, 1H, $J = 8.8$ Hz), 7.64 (d, 1H, $J = 2.4$ Hz), 8.41 (d, 1H, $J = 16.2$ Hz).

25


-329-

- 5 Step C: 5-Benzyloxy-2-(2-tert-butoxycarbonyl-vinyl)-benzoic acid 2-trimethylsilanyl-ethyl ester



- To a solution of 5-benzyloxy-2-(2-tert-butoxycarbonyl-vinyl)-benzoic acid (5.50 g, 15.5 mmol) in CH₂Cl₂ (100 mL) was added 2-trimethylsilyl-ethanol (3.67 g, 31.0 mmol), EDC (5.36 g, 27.9 mmol), and DMAP (3.67 g, 31.0 mmol). The resulting mixture was stirred at room temperature for 12 h, washed with aqueous NH₄Cl (150 mL), and extracted with CH₂Cl₂ (100 mL). The combined organics were dried (Na₂SO₄) and concentrated. The residue was purified using silica gel chromatography (hexanes/EtOAc 9/1) to afford the title compound as an oil (5.90 g, 84%). ¹H NMR (400 MHz, CDCl₃):
- 15 δ 0.07 (s, 9H), 1.14 (t, 2H, *J* = 8.8 Hz), 1.52 (s, 9H), 4.41 (t, 2H, *J* = 8.8 Hz), 5.10 (s, 2H), 6.20 (d, 1H, *J* = 15.6 Hz), 7.13 (dd, 1H, *J* = 3.0, 8.3 Hz), 7.34-7.44 (m, 5H), 7.50 (d, 1H, *J* = 3.0 Hz), 7.53 (d, 1H, *J* = 8.8 Hz), 8.41 (d, 1H, *J* = 16.1 Hz).

- 20 Step D: 2-(2-tert-Butoxycarbonyl-ethyl)-5-hydroxy-benzoic acid 2-trimethylsilylanyl-ethyl ester




- A solution of 5-benzyloxy-2-(2-tert-butoxycarbonyl-vinyl)-benzoic acid 2-trimethylsilylanyl-ethyl ester (6.20 g, 13.6 mmol) in EtOH (95 mL) was treated with Pd/C (5%, 0.775 g). The resulting suspension was treated with hydrogen at 60 psi for 6 h at room temperature. The catalyst was filtered through a pad of Celite, and the filtrate was concentrated to an oil (4.30 g, 86%). ¹H NMR (400 MHz, CDCl₃): δ 0.07 (s, 9H), 0.96 (t,
- 25

-330-

- 5 1H, $J = 8.8$ Hz), 1.12 (t, 1H, $J = 8.8$ Hz), 1.40 (s, 9 H), 2.52 (t, 1H, $J = 7.8$ Hz), 3.13 (t, 1H, $J = 7.8$ Hz), 3.73 (t, 1H, $J = 8.5$ Hz), 4.36 (t, 1H, $J = 8.5$ Hz), 5.00 (br s, 1 H), 6.88 (dd, 1H, $J = 2.4, 8.3$ Hz), 7.14 (d, 1H, $J = 8.3$ Hz), 7.35 (d, 1H, $J = 2.4$ Hz).

- Step E: 2-(2-tert-Butoxycarbonyl-ethyl)-5-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-
10 benzoic acid 2-trimethylsilanyl-ethyl ester.




- To a solution of 2-(2-tert-butoxycarbonyl-ethyl)-5-hydroxy-benzoic acid
2-trimethylsilanyl-ethyl ester (4.30 g, 11.7 mmol) and toluene-4-sulfonic acid 2-(5-
methyl-2-phenyl-oxazol-4-yl)-ethyl ester (4.61 g, 12.9 mmol) in DMF (30 mL) was added
15 Cs_2CO_3 (4.59 g, 14.1 mmol). The suspension was stirred at 55°C for 12 h. The reaction
mixture was quenched with H_2O (200 mL) and extracted with EtOAc (3 x 100 mL). The
combined organics were dried (Na_2SO_4) and concentrated to a residue, which was
purified using silica gel chromatography (hexanes/EtOAc 9/1 to 8/2) to afford the title
compound as a colorless oil (5.04 g, 78%). ^1H NMR (400 MHz, CDCl_3): δ 0.06 (s, 9H),
20 1.11 (t, 2H, $J = 8.8$ Hz), 1.40 (s, 9 H), 2.36 (s, 3 H), 2.51 (t, 1H, $J = 7.8$ Hz), 2.97 (t, 2H, J
= 6.8 Hz), 3.12 (t, 1H, $J = 7.8$ Hz), 4.24 (t, 2H, $J = 6.8$ Hz), 4.35 (t, 2H, $J = 8.8$ Hz), 6.88
(dd, 1H, $J = 3.0, 8.3$ Hz), 7.15 (d, 1H, $J = 8.3$ Hz), 7.37-7.44 (m, 4H), 7.97 (d, 2H, $J = 7.8$
Hz).

25

-331-

5. Step F: 2-(2-tert-Butoxycarbonyl-ethyl)-5-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzoic acid



- A solution of 2-(2-tert-butoxycarbonyl-ethyl)-5-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzoic acid 2-trimethylsilanylethyl ester (5.04 g, 9.13 mmol) in THF (100 mL) was treated with TBAF (20 mL, 1.0 M) at room temperature for 1 h. The reaction mixture was concentrated and purified using silica gel chromatography (hexanes/EtOAc 1/1) to afford the title compound as a colorless oil (4.04 g, 98%). ¹H NMR (400 MHz, CDCl₃): δ 1.38 (s, 9 H), 2.37 (s, 3 H), 2.55 (t, 2H, *J* = 7.3 Hz), 2.98 (t, 2H, *J* = 6.5 Hz), 3.15 (t, 2H, *J* = 7.3 Hz), 4.26 (t, 2H, *J* = 6.5 Hz), 4.35 (t, 2H, *J* = 8.8 Hz), 6.99 (dd, 1H, *J*=2.9, 8.8 Hz), 7.18 (d, 1H, *J*=8.3 Hz), 7.38-7.43 (m, 3H), 7.49 (d, 1H, *J*=2.9 Hz), 7.97 (d, 2H, *J*=7.8 Hz). MS (ES) *m/z* 522.3 [M+H]⁺.

Step G: 3-{2-Benzylcarbamoyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid


- To a solution of 2-(2-tert-butoxycarbonyl-ethyl)-5-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzoic acid (93 mg, 0.200 mmol) in CH₂Cl₂ (2 mL) was added benzylamine (52 mg, 0.49 mmol), EDC (54 mg, 0.28 mmol), triethylamine (0.057 mL, 0.40 mmol), and DMAP (catalyst). The resulting mixture was stirred at room temperature for 12 h and washed with aqueous NH₄Cl (2 mL). The organic layer was purified using a silica gel column (Sep-Pak column, 10 g; hexanes/EtOAc 1/1) to afford the *tert*-butyl ester intermediate. The ester was treated a mixture of CH₂Cl₂ (1.0 mL), TFA (0.8 mL), and water (0.1 mL) at room temperature for 2 h. The reaction mixture was concentrated and dried under vacuum to afford the title product as a white solid (35 mg, 36%). ¹H NMR (400 MHz, CDCl₃): δ 2.32 (s, 3 H), 2.59 (t, 2H, *J* = 7.3 Hz), 2.88-2.98 (m, 4H, *J* = 6.5 Hz), 4.16 (t, 2H, *J* = 6.5 Hz), 4.52 (s, 2H), 6.83 (dd, 1H, *J*=2.6, 8.4 Hz), 6.87 (d, 1 H, *J*=2.4 Hz), 7.10 (d, 1H, *J*=8.3 Hz), 7.20-7.30 (m, 3H), 7.37-7.40 (m, 3 H), 7.88-7.91 (m, 2H). MS (ES) *m/z* 485.2 [M+H]⁺, *m/z* 483.4 [M-H]⁻.

-332-


5

Example 583

3-{2-Benzylcarbamoyl-4-[2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-phenyl}-
propionic acid




Step A: 3-(2-Benzylaminocarbonyl-4-benzyloxy-phenyl)-acrylic acid *tert*-butyl ester



10

To a solution of 5-benzyloxy-2-(2-*tert*-butoxycarbonyl-vinyl)-benzoic acid (1.45 g, 4.09 mmol) in CH_2Cl_2 (40 mL) was added benzylamine (0.613 g, 5.73 mmol), EDC (1.254 g, 6.54 mmol), and DMAP (1.00 g, 8.18 mmol). The resulting mixture was stirred at room temperature for 12 h, washed with aqueous NH_4Cl (50 mL), and extracted with CH_2Cl_2 (2X50 mL). The combined organic layers were dried (Na_2SO_4) and concentrated. The residue was then purified using silica gel chromatography (hexanes/EtOAc 8/2) to afford the title compound as an oil (720 mg, 40%). ^1H NMR (400 MHz, CDCl_3): δ 1.52 (s, 9 H), 4.63 (d, 2H, $J = 5.5$ Hz), 5.09 (s, 2 H), 5.97 (br s, 1 H), 6.22 (d, 1H, $J = 15.6$ Hz), 7.01 (dd, 1H, $J = 2.7, 8.6$ Hz), 7.10 (d, 1H, $J = 2.8$ Hz), 7.28-7.42 (m, 10H), 7.55 (d, 1H, $J = 9.0$ Hz), 7.88 (d, 1H, $J = 16.0$ Hz).

20



A solution of 3-(2-benzylaminocarbonyl-4-benzyloxy-phenyl)-acrylic acid *tert*-butyl ester (720 mg, 1.625 mmol) in EtOH (15 mL) and THF (5 mL) was treated with

-333-

- 5 Pd/C (5%, 70 mg). The resulting suspension was treated with hydrogen using a balloon for 4 h at room temperature. The catalyst was filtered through a pad of celite and the filtrate was concentrated to an oil (450 mg, 78%). ¹H NMR (400 MHz, CDCl₃): δ 1.34 (s, 9 H), 2.56 (t, 2H, *J* = 7.0 Hz), 2.90 (t, 2H, *J* = 7.0 Hz), 4.59 (d, 1H, *J* = 5.9 Hz), 6.74 (dd, 1H, *J* = 2.7, 8.2 Hz), 6.89 (d, 1H, *J* = 2.7 Hz), 7.01 (d, 1H, *J* = 8.2 Hz), 7.25-7.36 (m, 5H).
- 10 MS (ES) *m/z* 356.2 [M+H]⁺, *m/z* 354.0 [M-H]⁻.

Step C: 3-{2-Benzylcarbamoyl-4-[2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid






- To a solution of 3-(2-benzylaminocarbonyl-4-hydroxy-phenyl)-propionic acid *tert*-butyl ester (50 mg, 0.140 mmol) and toluene-4-sulfonic acid 2-(5-methyl-2-(4-phenylphenyl)-oxazol-4-yl)-ethyl ester (73 mg, 0.168 mmol) in DMF (1.0 mL) was added K₂CO₃ (100 mg, 0.724 mmol). The suspension was stirred at 65°C for 12 h. The reaction mixture was quenched with H₂O (10 mL) and extracted with EtOAc (3x10 mL). The combined organic layers were dried (Na₂SO₄) and concentrated to a residue, which was
- 20 purified using silica gel chromatography (hexanes/EtOAc 7/3) to afford the *tert*-butyl ester intermediate as an oil. The ester was treated a mixture of CH₂Cl₂ (1.0 mL), TFA (0.8 mL) and water (0.1 mL) at room temperature for 2 h. The reaction mixture was concentrated and dried under vacuum to afford the title product as a white solid (37 mg, 47%). ¹H NMR (400 MHz, CDCl₃): δ 2.36 (s, 3 H), 2.59 (t, 2H, *J* = 7.3 Hz), 2.94 (t, 4H, *J* = 7.2 Hz), 4.20 (t, 2H, *J* = 6.5 Hz), 4.55 (s, 2H), 6.85 (dd, 1H, *J* = 2.9, 8.2 Hz), 6.90 (d, 1H, *J* = 2.4 Hz), 7.12 (d, 1H, *J* = 8.6 Hz), 7.20-7.36 (m, 6H), 7.42 (t, 2H, *J* = 7.4 Hz), 7.59 (d, 2H, *J* = 8.6 Hz), 7.64 (d, 2H, *J* = 8.2 Hz), 8.00 (d, 2H, *J* = 8.6 Hz). MS (ES) *m/z* 561.3 [M+H]⁺, *m/z* 559.5 [M-H]⁻.
- 25

5







Examples 584-643

Examples 584 to 643 are prepared by following a substantially similar procedure as described in Examples 582 and 583. Examples 584 to 628 are prepared by following a substantially similar procedure as described in Example 582. Examples 629 to 643 are prepared by following a substantially similar procedure as described in Example 583.

10







No.	Compounds	Name	MS (ES+)
584		3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-phenyl-carbamoyl-phenyl}-propionic acid	471.2
585		3-{2-Benzylcarbamoyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	485.2
586		3-{2-(3,4-Dichloro-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	553
587		3-{2-(4-Methoxy-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	515.2
588		3-{2-[(Biphenyl-3-yl methyl)-carbamoyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	561.2







-335-

No.	Compounds	Name	MS (ES+)
589		3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-phenethyl-carbamoyl-phenyl}-propionic acid	499.3
590		3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(3-phenyl-propylcarbamoyl)-phenyl]-propionic acid	513.3
591		3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(thiophen-2-ylmethyl)-carbamoyl]-phenyl}-propionic acid	491.2
592		3-{2-Hexylcarbamoyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	479.3
593		3-{2-Methylcarbamoyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	409.1
594		3-{2-Butylcarbamoyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	451.1







No.	Compounds	Name	MS (ES+)
595		3-{2-Isopropylcarbamoyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	437.2
596		3-{2-(Cyclohexylmethylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	491.2
597		3-{2-tert-Butylcarbamoyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	451.2
598		3-{2-Carbamoyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	395.1
599		3-{2-(2-Fluorobenzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	503.1
600		3-{2-(2-Chlorobenzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	519.1

-337-







No.	Compounds	Name	MS (ES+)
601		3-{2-(2,4-Dichloro-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	553
602		3-{2-(2-Methoxy-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	515.1
603		3-{2-(Indan-1-ylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	511.1
604		3-{2-(3-Fluoro-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	503.1
605		3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(4-trifluoromethyl-benzylcarbamoyl)-phenyl]-propionic acid	553
606		3-{2-(3-Methyl-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	499.1







No.	Compounds	Name	MS (ES+)
607		3-{2-(4-Fluorobenzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	503.1
608		3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(naphthalen-1-ylmethyl)-carbamoyl]-phenyl}-propionic acid	535.1
609		3-{2-(4-Methanesulfonylbenzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	563.1
610		3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(2-trifluoromethylbenzylcarbamoyl)-phenyl]-propionic acid	553.1
611		3-{2-(4-Nitrobenzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	530.1
612		3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(4-sulfamoylbenzylcarbamoyl)-phenyl]-propionic acid	564.1







-339-

No.	Compounds	Name	MS (ES+)
613		3-{2-(3,5-Dimethyl-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	513.2
614		3-{2-(4-tert-Butyl-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	541.2
615		3-{2-(2-Methyl-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	499.2
616		3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(pyridin-4-ylmethyl)-carbamoyl]-phenyl}-propionic acid	486.2
617		3-{2-(3-Methoxy-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	515.2
618		3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(3-trifluoromethyl-benzylcarbamoyl)-phenyl]-propionic acid	553.1

-340-


No.	Compounds	Name	MS (ES+)
619		3-{2-(3,5-Bis-trifluoro methyl-benzyl-carbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	621.1
620		3-{2-(3-Chloro-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	519.1
621		3-{2-(3-Fluoro-5-trifluoro methyl-benzyl carbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	571.1
621		3-{2-(3,5-Difluoro-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	521.1
623		3-{2-(3,5-Dichloro-benzylcarbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	553
624		(R)-3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(1-phenyl-ethylcarbamoyl)-phenyl]-propionic acid	499.2

No.	Compounds	Name	MS (ES+)
625		3-{2-(Benzyl-ethyl-carbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	513.2
626		3-{2-(Benzyl-methyl-carbamoyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	499.2
627		(S)-3-{2-[(Carboxy-phenyl-methyl) carbamoyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	529.1
628		(S)-3-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-(1-phenyl-ethylcarbamoyl)-phenyl]-propionic acid	499.2
629		3-{2-Benzylcarbamoyl-4-[2-(2-biphenyl-3-yl-5-methyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	561.3
630		3-{2-Benzylcarbamoyl-4-[2-(2-biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	561.3

No.	Compounds	Name	MS (ES+)
631		3-{2-Benzylcarbamoyl-4-[2-(2-cyclohexyl-5-methyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid	491.3
632		3-(2-Benzylcarbamoyl-4-{2-[5-methyl-2-(1-methyl-cyclohexyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid	505.4
633		3-{2-Benzylcarbamoyl-4-[2-(5-methyl-2-morpholin-4-yl-thiazol-4-yl)-ethoxy]-phenyl}-propionic acid	510.3
634		(R)-3-[4-[2-(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-(1-phenyl-ethylcarbamoyl)-phenyl]-propionic acid	575.3
635		(R)-3-[4-[2-(2-Biphenyl-3-yl-5-methyl-oxazol-4-yl)-ethoxy]-2-(1-phenyl-ethylcarbamoyl)-phenyl]-propionic acid	575.3
636		(R)-3-[4-[2-(5-Methyl-2-morpholin-4-yl-thiazol-4-yl)-ethoxy]-2-(1-phenyl-ethylcarbamoyl)-phenyl]-propionic acid	524.2

No.	Compounds	Name	MS (ES+)
637		3-[4-[2-(2-Biphenyl-4-yl)-5-methyl-oxazol-4-yl)-ethoxy]-2-(3,5-difluorobenzylcarbamoyl)-phenyl]-propionic acid	597.3
638		3-[4-[2-(2-Biphenyl-3-yl)-5-methyl-oxazol-4-yl)-ethoxy]-2-(3,5-difluorobenzylcarbamoyl)-phenyl]-propionic acid	597.3
639		3-{2-(3,5-Difluorobenzylcarbamoyl)-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)-ethoxy]-phenyl}-propionic acid	546.1
640		3-{2-(3,5-Difluorobenzylcarbamoyl)-4-[2-(5-methyl-2-phenylthiazol-4-yl)-ethoxy]-phenyl}-propionic acid	537.1
641		3-(2-Benzylcarbamoyl-4-{2-[5-methyl-2-(4-phenoxy-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid	577.3
642		3-[2-Benzylcarbamoyl-4-(2-{5-methyl-2-[4-(methylphenyl-amino)-phenyl]-oxazol-4-yl}-ethoxy)-phenyl]-propionic acid	590.2


-344-

No.	Compounds	Name	MS (ES+)
643		3-(2-Benzylcarbamoyl-4-{2-[5-methyl-2-(4-phenylamino-phenyl)-oxazol-4-yl]-ethoxy}-phenyl)-propionic acid	576.2


5

Example 644

3-{2-Cyclopentylcarbamoyloxymethyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}-propionic acid



10 Step A: 3-{2-Formyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}acrylic acid *tert*-butyl ester




A flame-dried 100 mL round bottomed flask was charged with toluene-4-sulfonic acid 2-(5-methyl-2-phenyloxazol-4-yl) ethyl ester (3.74 g, 10.5 mmol), 3-(2-
15 formyl-4-hydroxyphenyl)acrylic acid *tert*-butyl ester (2.0 g, 8.05 mmol), and anhydrous DMF (40 mL). Cesium carbonate (3.94 g, 12.1 mmol) was added, and the reaction was heated to 55° C under a nitrogen atmosphere for 18 h. The volatiles were removed *in vacuo*, and the crude residue was dissolved in EtOAc (250 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and concentrated to a yellow oil (1.70 g, 49%).

20 ¹H NMR (400 MHz, CDCl₃) δ 1.54 (s, 9H), 2.38 (s, 3H), 3.01 (t, J= 6.6 Hz, 2H), 4.34 (t,

-345-


- 5 J= 6.6 Hz, 2H), 6.24 (d, J= 15.6 Hz, 1H), 7.12 (dd, J= 8.8, 2.9 Hz, 1H), 7.38 (d, J= 2.9 Hz, 1H), 7.40-7.45 (m, 3H), 7.58 (d, J= 8.8 Hz, 1H), 7.98 (dd, J= 6.4, 2.0 Hz, 2H), 8.33 (d, J= 15.6 Hz, 1H), 10.3 (s, 1H). MS (ES) m/e 434 (M+1).

Step B: 3-{2-Formyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}acrylic acid
10 *tert*-butyl ester



- A 100 mL round bottomed flask was charged with 3-{2-formyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)-ethoxy]phenyl}acrylic acid *tert*-butyl ester (1.69 g, 3.90 mmol), THF (40 mL), and then 10% Pd/C catalyst (0.17 g). The reaction was stirred
15 vigorously under a hydrogen atmosphere at one atmosphere for 18 h. The mixture was filtered through Celite and concentrated to a yellow solid (1.70 g, quantitative). ^1H NMR (400 MHz, CDCl_3) δ 1.39 (d, J= 4.9 Hz, 9H), 2.37 (d, J= 2.9 Hz, 3H), 2.51 (t, J= 7.6 Hz, 2H), 2.99 (t, J= 6.6 Hz, 2H), 3.23 (t, J= 7.1 Hz, 2H), 4.29 (t, J= 6.6 Hz, 2H), 7.04-7.10 (m, 1H), 7.21 (d, J= 8.3 Hz, 1H), 7.34 (d, J= 2.4 Hz, 1H), 7.38-7.45 (m, 3H), 7.97 (dd, J= 7.3,
20 2.0 Hz, 2H), 10.2 (s, 1H). MS (ES) m/e 436 (M+1).

Step C: 3-{2-Hydroxymethyl-4-[2-(2-phenyl-oxazol-4-yl)ethoxy]phenyl}propionic acid
tert-butyl ester




- 25 A 100 mL round bottomed flask was charged with 3-{2-formyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester (1.82 g, 4.18 mmol) and absolute ethanol (20 mL). The stirring solution was cooled in an ice/ethanol bath and treated with sodium borohydride (0.31 g, 8.36 mmol). The cold bath was removed, and the mixture was stirred at ambient temperature under a nitrogen atmosphere for 2 h. The

-346-

5 reaction mixture was poured into EtOAc (100 mL) and ice water (100 mL). The organic layer was washed with brine, dried (Na_2SO_4), and concentrated to a yellow oil (1.62 g, 88%). ^1H NMR (400 MHz, CDCl_3) δ 1.38 (s, 9H), 2.37 (s, 3H), 2.54 (t, $J=7.6$ Hz, 2H), 2.88 (t, $J=7.6$ Hz, 2H), 2.97 (t, $J=6.6$ Hz, 2H), 4.23 (t, $J=6.6$ Hz, 2H), 4.66 (s, 2H), 6.78 (dd, $J=8.3, 2.9$ Hz, 1H), 6.94 (d, $J=2.4$ Hz, 1H), 7.09 (d, $J=8.3$ Hz, 1H), 7.38-7.45 (m, 3H), 7.97 (dd, $J=8.1, 1.7$ Hz, 2H). MS (ES) m/e 438 ($M+1$).

Step D: 3-{2-Cyclopentylcarbamoyloxymethyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}-propionic acid *tert*-butyl ester



15 A 15 mL round bottomed flask was charged with 3-{2-hydroxymethyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester (0.10 g, 0.23 mmol), cyclopentylisocyanate (0.15 mL, 1.38 mmol), and anhydrous CH_2Cl_2 (0.75 mL). A 1.0 M HCl solution in ether (0.115 mL, 0.115 mmol) was added, and the reaction mixture was stirred at ambient temperature under a nitrogen atmosphere for 24 h. The reaction mixture was diluted with CH_2Cl_2 (30 mL), washed with brine, dried (Na_2SO_4), and concentrated to an oil. The crude product was purified using radial chromatography (EtOAc:hexanes 10:90 to 35:65) to a colorless oil (0.105 g, 83%). ^1H NMR (400 MHz, CDCl_3) δ 1.42 (s, 9H), 1.56-1.65 (m, 4H), 1.94-2.04 (m, 2H), 2.37 (s, 3H), 2.46 (t, $J=7.6$ Hz, 2H), 2.87 (t, $J=8.1$ Hz, 2H), 2.96 (t, $J=6.6$ Hz, 2H), 3.97 (br s, 1H), 4.22 (t, $J=6.6$ Hz, 2H), 4.76 (br s, 1H), 5.07 (s, 2H), 6.90 (dd, $J=8.3, 2.4$ Hz, 1H), 6.89 (s, 1H), 7.10 (d, $J=8.8$ Hz, 1H), 7.40-7.45 (m, 3H), 7.96-7.98 (m, 2H). MS (ES) m/e 549 ($M+1$).

-347-

- 5 Step E: 3-{2-Cyclopentylcarbamoyloxymethyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}-propionic acid

A 15 mL round bottom flask was charged with 3-{2-cyclopentylcarbamoyloxymethyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester (0.097 g, 0.18 mmol), CH₂Cl₂ (1.2 mL), and then trifluoroacetic acid (1.2 mL). The solution was stirred at ambient temperature under a nitrogen atmosphere for 4 h and was concentrated. The residue was purified using radial chromatography (McOH:CH₂Cl₂ 2:98 to 10:90) to give a white solid (0.085 g, 93%). ¹H NMR (400 MHz, CDCl₃) δ 1.37 (br s, 2H), 1.63 (br s, 4H), 1.93-2.05 (m, 2H), 2.39 (s, 3H), 2.64 (br s, 2H), 2.91-3.18 (m, 4H), 3.99 (br s, 1H), 4.21 (s, 2H), 4.87 (br s, 1H), 5.10 (br s, 2H), 6.80 (s, 1H), 6.89 (s, 1H), 7.00-7.25 (br s, 1H), 7.36-7.52 (m, 3H), 7.95 (s, 2H). MS (ES) *m/e* 493 (M+1).

The following Examples 645 to 651 are prepared by following a substantially similar procedure as described in Example 644.

20


Example 645

3-{2-Isopropylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester
MS (ES) *m/e* 523 (M+1).

25

Example 646

3-{2-Isopropylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



30 ¹H NMR (400 MHz, CDCl₃) δ 1.13 (d, J= 5.4 Hz, 6H), 2.37 (s, 3H), 2.61 (t, J= 7.8 Hz, 2H), 2.92-2.98 (m, 4H), 3.80 (br s, 1H), 4.21 (t, J= 6.6 Hz, 2H), 4.66 (br s, 1H), 5.08 (s,

-348-


5 2H), 6.81 (dd, $J = 8.6, 2.7$ Hz, 1H), 6.90 (s, 1H), 7.11 (d, $J = 8.3$ Hz, 1H), 7.41-7.44 (m, 3H), 7.97 (dd, $J = 4.4, 2.9$ Hz, 2H). MS (ES) m/e 467 (M+1).

Example 647

10 3-{2-Benzylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester
MS (ES) m/e 571 (M+1).

Example 648


15 3-{2-Benzylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



^1H NMR (400 MHz, CDCl_3) δ 2.37 (s, 3H), 2.60 (t, $J = 7$ Hz, 2H), 2.92-2.97 (m, 4H), 4.20 (t, $J = 7$ Hz, 2H), 4.36 (d, $J = 5.9$ Hz, 2H), 5.13 (s, 2H), 5.18 (br s, 1H), 6.80 (d, $J = 8.3$ Hz, 1H), 6.91 (s, 1H), 7.10 (d, $J = 8.3$ Hz, 1H), 7.24-7.30 (m, 5H), 7.42 (s, 3H), 7.97 (s, 2H).
20 MS (ES) m/e 515 (M+1).

Example 649

Morpholine-4-carboxylic acid 2-(2-carboxyethyl)-5-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]benzyl ester



25 MS (ES) m/e 495 (M+1).

-349-

5

Example 650


3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester

MS (ES) *m/e* 563 (M+1)

10

Example 651

3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



15

¹H NMR (400 MHz, CDCl₃) δ 1.08-1.18 (m 3H), 1.21-1.42 (m, 2H), 1.57 (br s, 1H), 1.66-1.70 (m, 2H), 1.90 (br s, 2H), 2.40 (s, 3H), 2.61 (t, J= 7.7 Hz, 2H), 2.94 (t, J= 7.8 Hz, 2H), 3.01 (t, J= 6.4 Hz, 2H), 3.48 (br s, 1H), 4.22 (t, J= 6.4 Hz, 2H), 4.74 (br s, 1H), 5.07 (s, 2H), 6.79-6.89 (m, 1H), 6.90 (s, 1H), 7.11 (d, J= 8.3 Hz, 1H), 7.44-7.46 (m, 3H), 7.98 (dd, J= 6.7, 2.8 Hz, 2H). MS (ES) *m/e* 507 (M+1).


20

-350-


5

Example 652

3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}-
propionic acid



Step A: 3-[4-(*tert*-Butyldiphenylsilyloxy)-2-formylphenyl]acrylic acid *tert*-butyl ester




10

A flame-dried 200 mL round bottomed flask was charged with 3-(2-formyl-4-hydroxyphenyl)acrylic acid *tert*-butyl ester (10.0 g, 40.3 mmol), *tert*-butylchlorodiphenylsilane (12.6 mL, 48.3 mmol), and dry CH₂Cl₂ (150 mL). Triethylamine (11.2 mL, 80 mmol) and N,N-dimethylaminopyridine (1.0 g, 1.0 mmol) were added, and the mixture was stirred at ambient temperature under a nitrogen atmosphere for 16 h. The reaction mixture was diluted with additional CH₂Cl₂, washed with brine (2x), dried (Na₂SO₄), and concentrated to an oil. The crude product was purified using the Biotage medium pressure chromatography system (EtOAc:hexanes 5:95) to give a pale yellow oil (18.3 g, 93%). ¹H NMR (400 MHz, CDCl₃) δ 1.12 (s, 9H), 1.51 (d, J= 15.1 Hz, 9H), 6.15 (d, J= 5.1 Hz, 1H), 6.87 (dd, J= 8.6, 2.7 Hz, 1H), 7.34-7.47 (m, 7H), 7.68-7.73 (m, 5H), 8.28 (d, J= 15.6 Hz, 1H), 10.1 (s, 1H). MS (ES) *m/e* 487 (M+1).

20


-351-

- 5 Step B: 3-[4-(*tert*-Butyldiphenylsilanyloxy)-2-hydroxymethylphenyl]propionic acid *tert*-butyl ester



- A 500 mL Parr hydrogenation bottle was charged with 3-[4-(*tert*-butyldiphenylsilanyloxy)-2-formylphenyl]acrylic acid *tert*-butyl ester (18.3 g, 37.6 mmol), THF (60 mL), and methanol (120 mL). Triethylamine (2 mL) and then 5% Pd/C (5.9 g) were added. The mixture was shaken with hydrogen at 60 psi pressure for 48 h. The mixture was filtered through Celite and concentrated to an oil. This oil was purified using the Biotage medium pressure chromatography (EtOAc:hexanes 15:85) to a pale yellow oil (12.8 g, 69%). ¹H NMR (400 MHz, CDCl₃) δ 1.09 (s, 9H), 1.35 (s, 9H), 2.49 (t, J= 7.6 Hz, 2H), 2.83 (t, J= 7.6 Hz, 2H), 4.55 (s, 2H), 6.56 (dd, J= 8.3, 2.4 Hz, 1H), 6.83 (d, J= 2.9 Hz, 1H), 6.87 (d, J= 8.3 Hz, 1H), 7.33-7.44 (m, 6H), 7.70 (dd, J= 7.8, 1.5 Hz, 4H). MS (ES) *m/e* 508 (M+NH₄).
- 15

- Step C: 3-[4-(*tert*-Butyldiphenylsilanyloxy)-2-cyclohexylcarbamoyloxymethylphenyl]propionic acid *tert*-butyl ester
- 20




- A 100 mL round bottomed flask under N₂ were charged with 3-[4-(*tert*-butyldiphenylsilanyloxy)-2-hydroxymethylphenyl]propionic acid *tert*-butyl ester (3.0 g, 6.11 mmol), cyclohexylisocyanate (4.7 mL, 36.7 mmol), and anhydrous CH₂Cl₂ (25 mL). A 1.0 M HCl solution in ether (3.06 mL, 3.06 mmol) was added, and the reaction mixture was stirred at ambient temperature under a nitrogen atmosphere for 16 h. The mixture was diluted with CH₂Cl₂ (100 mL), washed with brine, dried (Na₂SO₄), and concentrated
- 25

-352-

5 to a brown oil. This crude oil was purified using the Biotage medium pressure chromatography system (EtOAc:hexanes 5:95) to a colorless oil (3.1 g, 82%). ¹H NMR (400 MHz, CDCl₃) δ 1.08-1.24 (m, 1H), 1.26-1.40 (m, 11H), 1.56-1.62 (m, 2H), 1.68-1.72 (m, 2H), 2.41 (t, J= 8.3 Hz, 2H), 2.80 (t, J= 7.8 Hz, 2H), 3.47 (br s, 1H), 4.56 (br s, 1H), 4.95 (s, 2H), 6.57 (dd, J= 8.3, 2.7 Hz, 1H), 6.80 (s, 1H), 6.87 (d, J= 8.6 Hz, 1H),
10 7.33-7.44 (m, 6H), 7.69-7.72 (m, 4H). MS (ES) *m/e* 616 (M+1).


Step D: 3-(2-Cyclohexylcarbamoyloxymethyl-4-hydroxyphenyl)propionic acid *tert*-butyl ester



15 A 500 mL round bottomed flask was charged with 3-[4-(*tert*-butyl-diphenylsilanyloxy)-2-cyclohexylcarbamoyloxymethylphenyl]propionic acid *tert*-butyl ester (3.1 g, 5.03 mmol) and anhydrous THF (180 mL). Tetrabutylammonium fluoride (15.1 mL, 15.1 mmol, 1.0M in THF) was added, and the reaction mixture was stirred at ambient temperature under a nitrogen atmosphere for 4 h. The mixture was concentrated,
20 and the residue was diluted with EtOAc (100 mL), washed with brine, dried over (Na₂SO₄), and concentrated to give an oil. This crude oil was purified using the Biotage medium pressure chromatography system (EtOAc:hexanes 10:90 to 50:50) to a colorless oil (1.67 g, 88%). ¹H NMR (400 MHz, CDCl₃) δ 1.04-1.19 (m, 3H), 1.24-1.38 (m, 2H), 1.42 (s, 9H), 1.57-1.62 (m, 1H), 1.66-1.71 (m, 2H), 1.91-2.05 (m, 2H), 2.47 (t, J= 7.8 Hz, 2H), 2.88 (t, J= 7.8 Hz, 2H), 3.48-3.50 (m, 1H), 4.77-4.78 (m, 1H), 5.07 (s, 2H), 5.40 (s, 2H), 6.73 (dd, J= 8.3, 2.9 Hz, 1H), 6.84 (d, J= 2.4 Hz, 1H), 7.07 (d, J= 8.3 Hz, 1H). MS (ES) *m/e* 378 (M+1).
25

-353-

- 5 Step E: 3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester



- A 40 mL Carousel tube was charged with 3-(2-cyclohexyl-carbamoyloxymethyl-4-hydroxyphenyl)propionic acid *tert*-butyl ester (0.10 g, 0.26 mmol) in anhydrous DMF (1.0 mL). Toluene-4-sulfonic acid 2-(5-methyl-2-phenyloxazol-4-yl)ethyl ester (0.104g, 0.292 mmol) and cesium carbonate (0.13 g, 0.40 mmol) were added. The mixture was stirred and heated at 55° C under a nitrogen atmosphere for 30 h and was concentrated. The residue was diluted with EtOAc (50 mL) and washed twice with brine (2x), dried (Na₂SO₄), and concentrated. The crude residue was purified using radial chromatography (EtOAc: CH₂Cl₂ 2:98 to 5:95) to give a white solid (0.085 g, 57%). ¹H NMR (400 MHz, CDCl₃) δ 1.07-1.19 (m, 3H), 1.24-1.38 (m, 2H), 1.42 (s, 9H), 1.56-1.60 (m, 1H), 1.67-1.70 (m, 2H), 1.91-1.93 (m, 2H), 2.37 (s, 3H), 2.46 (t, J= 7.8 Hz, 2H), 2.88 (t, J= 7.8 Hz, 2H), 2.96 (t, J= 7.8 Hz, 2H), 3.48-3.50 (m, 1H), 4.22 (t, J= 6.8 Hz, 2H), 4.69-4.71 (m, 1H), 5.07 (s, 2H), 6.80 (dd, J= 8.3, 2.4 Hz, 1H), 6.90 (d, J= 2.4 Hz, 1H), 7.10 (d, J= 8.3 Hz, 1H), 7.39-7.45 (m, 3H), 7.98 (dd, J= 4.2, 2.2 Hz, 2H). MS (ES) *m/e* 563 (M+1).

Step F: 3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid

- 25 A 25 mL round bottomed flask was charged with 3-{2-cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}-propionic acid *tert*-butyl ester (0.080 g, 0.14 mmol) and CH₂Cl₂ (2 mL). Trifluoroacetic acid (2mL) was added, and the solution was stirred at ambient temperature under a nitrogen atmosphere for 1.5 h. The solution was concentrated to give a white solid (0.068 g, 94%). ¹H NMR (400 MHz, CDCl₃) δ 1.08-1.18 (m 3H), 1.21-1.42 (m, 2H), 1.57 (br s,
- 30


-354-

5 1H), 1.66-1.70 (m, 2H), 1.90 (br s, 2H), 2.40 (s, 3H), 2.61 (t, J= 7.7 Hz, 2H), 2.94 (t, J= 7.8 Hz, 2H), 3.01 (t, J= 6.4 Hz, 2H), 3.48 (br s, 1H), 4.22 (t, J= 6.4 Hz, 2H), 4.74 (br s, 1H), 5.07 (s, 2H), 6.79-6.89 (m, 1H), 6.90 (s, 1H), 7.11 (d, J= 8.3 Hz, 1H), 7.44-7.46 (m, 3H), 7.98 (dd, J= 6.7, 2.8 Hz, 2H). MS (ES) *m/e* 507 (M+1).

10 The following Examples 653 to 661 are prepared by following a substantially similar procedure as described in Example 652.

Example 653


15 3-{4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-cyclohexylcarbamoyloxymethylphenyl}propionic acid *tert*-butyl ester



MS (ES) *m/e* 639 (M+1).

Example 654

20 3-{4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-cyclohexylcarbamoyloxymethylphenyl}propionic acid



¹H NMR (400 MHz, CDCl₃) δ 1.14-1.25 (m, 3H), 1.31-1.34 (m, 2H), 1.56-1.66 (m, 1H), 1.69-1.70 (m, 2H), 1.89 (br s, 2H), 2.46 (s, 3H), 2.62 (t, J= 7.6 Hz, 2H), 2.93 (t, J= 7.8 Hz, 2H), 3.11 (t, J= 5.9 Hz, 2H), 3.47 (br s, 1H), 4.24 (t, J= 5.9 Hz, 2H), 4.83 (br s, 1H), 5.08 (s, 2H), 6.80 (d, J= 6.8 Hz, 1H), 6.90 (d, J= 2.9 Hz, 1H), 7.11 (d, J= 8.3 Hz, 1H),

25

-355-

- 5 7.41 (d, J= 7.3 Hz, 1H), 7.48 (t, J= 7.6 Hz, 2H), 7.64 (d, J= 7.3 Hz, 2H), 7.74 (d, J= 8.3 Hz, 2H), 8.09 (d, J= 8.3 Hz, 2H). MS (ES) *m/e* 583 (M+1).

Example 655

- 10 3-{4-[2-(2-Biphenyl-3-yl-5-methyloxazol-4-yl)ethoxy]-2-cyclohexylcarbamoyloxymethylphenyl}propionic acid *tert*-butyl ester
MS (ES) *m/e* 639 (M+1)

Example 656

- 15 3-{4-[2-(2-Biphenyl-3-yl-5-methyloxazol-4-yl)ethoxy]-2-cyclohexylcarbamoyloxymethylphenyl}propionic acid
¹H NMR (400 MHz, CDCl₃) δ 1.26-1.41 (m, 3H), 1.44-1.56 (m, 2H), 1.75 (br s, 1H), 1.84-1.88 (m, 2H), 2.08-2.10 (m, 2H), 2.63 (s, 3H), 2.80 (t, J= 7.8 Hz, 2H), 3.12 (t, J= 7.9 Hz, 2H), 3.27 (t, J= 6.1 Hz, 2H), 3.66 (br s, 4.42 (t, J= 6.1 Hz, 2H), 4.97 (br s, 1H), 5.27 (s, 2H), 7.04 (d, J= 6.4 Hz, 1H), 7.08 (d, J= 2.4 Hz, 1H), 7.30 (d, J= 8.3 Hz, 1H), 7.55-
20 7.76 (m, 4H), 7.85-7.86 (m, 2H), 7.92 (d, J= 7.8 Hz, 1H), 8.15 (d, J= 7.8 Hz, 1H), 8.44 (d, J= 1.5 Hz, 1H). MS (ES) *m/e* 583 (M+1).

Example 657


- 25 3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester
MS (ES) *m/e* 588 (M+1)

-356-

5

Example 658

3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl}propionic acid



¹H NMR (400 MHz, CDCl₃) δ 1.11-1.37 (m, 5H), 1.57-1.61 (m, 1H), 1.68-1.72 (m, 2H),
 10 1.90-1.93 (m, 2H), 2.29 (s, 3H), 2.58 (t, J= 7.8 Hz, 2H), 2.90 (t, J= 7.6 Hz, 2H), 3.06 (t,
 J= 5.1 Hz, 2H), 3.47-3.50 (m, 1H), 3.66 (t, J= 4.6 Hz, 4H), 3.85 (t, J= 4.9 Hz, 4H), 4.20 (t,
 J= 5.4 Hz, 2H), 5.06 (s, 2H), 6.74 (d, J= 8.3 Hz, 1H), 6.86 (s, 1H), 7.08 (d, J= 8.3 Hz,
 1H). MS (ES) *m/e* 532 (M+1).

15

Example 659


3-(2-Cyclohexylcarbamoyloxymethyl-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl)propionic acid *tert*-butyl ester

MS (ES) *m/e* 655 (M+1)

20

Example 660

3-(2-Cyclohexylcarbamoyloxymethyl-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl)propionic acid




¹H NMR (400 MHz, CDCl₃) δ 1.15-1.24 (m, 3H), 1.26-1.34 (m, 2H), 1.57 (br s, 1H),
 25 1.67-1.70 (m, 2H), 1.90 (br s, 2H), 2.43 (s, 3H), 2.61 (t, 7.6 Hz, 2H), 2.93 (t, J= 7.3 Hz,
 2H), 3.08 (t, J= 5.6 Hz, 2H), 3.47-3.52 (m, 1H), 4.22 (t, J= 5.6 Hz, 1H), 4.84 (br s, 1H),

-357-

- 5 1H), 6.79 (d, 2.4 Hz, 1H), 6.88 (d, J= 2.4 Hz, 1H), 7.00-7.12 (m, 5H), 7.21 (t, J= 7.6 Hz, 1H), 7.41 (t, J= 7.8 Hz, 2H), 7.99 (d, 8.8 Hz, 2H). MS (ES) *m/e* 599 (M+1).

Example 661

- 10 3-(4-{2-[2-(4-Bromophenyl)-5-methyloxazol-4-yl]ethoxy}-2-cyclohexylcarbamoyloxymethylphenyl)propionic acid *tert*-butyl ester




MS (ES) *m/e* 641 (M+1).

-358-

5


Example 662

3-(2-Cyclohexylcarbamoxyloxymethyl-4-{2-[5-methyl-2-(4-morpholin-4-ylphenyl)oxazol-4-yl]ethoxy}phenyl)propionic acid



Step A: 3-(2-Cyclohexylcarbamoxyloxymethyl-4-{2-[5-methyl-2-(4-morpholin-4-ylphenyl)oxazol-4-yl]ethoxy}phenyl)propionic acid *tert*-butyl ester

10



A 1 mL microvial under a nitrogen atmosphere was charged with 3-(4-{2-[2-(4-bromophenyl)-5-methyloxazol-4-yl]ethoxy}-2-cyclohexylcarbamoxyloxymethylphenyl)-propionic acid *tert*-butyl ester (0.30 g, 0.468 mmol; Example 39), anhydrous toluene (0.5 mL), and then morpholine (0.053 mL, 0.61 mmol). Tris(dibenzylideneacetone)-dipalladium(0) (0.004 g, 0.0044 mmol), 2-(di-*tert*-butylphosphine)biphenyl (0.006 g, 0.020 mmol), and sodium *tert*-butoxide (0.063g, 0.655 mmol) were added sequentially. The mixture was stirred at ambient temperature under a nitrogen atmosphere for 5 h and was poured into EtOAc (50 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and concentrated to a dark yellow oil. This crude product was purified using radial chromatography (EtOAc:hexanes 15:85 to 50:50) to give a yellow oil (0.058 g, 19%). ¹H NMR (400 MHz, CDCl₃) δ 1.07-1.18 (m, 3H), 1.29-1.40 (m, 1H), 1.42 (s, 9H), 1.52-1.70 (m, 4H), 1.91-1.93 (m, 2H), 2.34 (s, 3H), 2.46

20

-359-

- 5 (t, J= 8.1 Hz, 2H), 2.87 (t, J= 7.8 Hz, 2H), 2.94 (t, J= 6.8 Hz, 2H), 3.24 (t, J= 4.9 Hz, 4H), 3.48-3.50 (m, 1H), 3.86 (t, J= 4.9 Hz, 4H), 4.20 (t, J= 6.6 Hz, 2H), 5.07 (s, 2H), 6.79 (dd, J= 8.6, 2.7 Hz, 1H), 6.85-6.93 (m, 3H), 7.09 (d, J= 8.3 Hz, 1H), 7.87 (dd, J= 11.5, 2.7 Hz, 2H). MS (ES) *m/e* 648 (M+1).

- 10 Step B: 3-(2-Cyclohexylcarbamoyloxymethyl-4-{2-[5-methyl-2-(4-morpholin-4-ylphenyl)oxazol-4-yl]ethoxy}phenyl)propionic acid

A 15 mL round bottomed flask was charged with 3-(2-cyclohexylcarbamoyloxymethyl-4-{2-[5-methyl-2-(4-morpholin-4-ylphenyl)oxazol-4-yl]ethoxy}phenyl)propionic acid (0.058 g, 0.09 mmol) and CH₂Cl₂ (1 mL).

- 15 Trifluoroacetic acid (0.5 mL) was added, and the solution was stirred at ambient temperature under a nitrogen atmosphere for 1.5 h. The mixture was concentrated to give a yellow solid (0.052 g, 98%). ¹H NMR (400 MHz, CDCl₃) δ 1.03-1.11 (m, 3H), 1.18-1.30 (m, 2H), 1.45-1.50 (m, 1H), 1.60-1.63 (m, 2H), 1.82-1.85 (m, 2H), 2.38 (s, 3H), 2.52 (t, J= 7.6 Hz, 2H), 2.84 (t, J= 7.6 Hz, 2H), 3.04 (t, J= 5.6 Hz, 2H), 3.30 (t, J= 4.9 Hz, 4H),
20 3.39 (br s, 1H), 3.80 (t, J= 4.9 Hz, 4H), 4.17 (t, J= 5.9 Hz, 2H), 4.86 (br s, 1H), 4.99 (s, 2H), 6.70 (d, J= 8.3 Hz, 1H), 6.81 (d, J= 2.4 Hz, 1H), 7.03 (d, J= 8.3 Hz, 1H), 7.90 (d, J= 9.3 Hz, 2H). MS (ES) *m/e* 592 (M+1).


- The following Examples 663 to 664 are prepared by following a
25 substantially similar procedure as described in Example 662.

-360-

5

Example 663

3-(2-Cyclohexylcarbamoyloxymethyl-4-{2-[5-methyl-2-(4-phenylaminophenyl)oxazol-4-yl]ethoxy}phenyl)propionic acid




¹H NMR (400 MHz, CDCl₃) δ 1.11-1.42 (m, 5H), 1.56 (br s, 1H), 1.66-1.69 (m, 2H), 1.89-1.92 (m, 2H), 2.38 (s, 3H), 2.60 (t, J= 7 Hz, 2H), 2.91 (t, J= 7 Hz, 2H), 3.03 (t, J= 5.6 Hz, 2H), 4.22 (t, J= 6.1 Hz, 2H), 4.86 (br s, 1H), 5.06 (s, 2H), 6.78 (d, J= 7.8 Hz, 1H), 6.89 (s, 1H), 7.04-7.10 (m, 4H), 7.16 (d, J= 7.8 Hz, 2H), 7.34 (t, J= 7.8 Hz, 2H), 7.88 (d, J= 8.3 Hz, 2H). MS (ES) *m/e* 598 (M+1).

15

Example 664

3-[2-Cyclohexylcarbamoyloxymethyl-4-(2-{5-methyl-2-[4-(methylphenylamino)phenyl]oxazol-4-yl}ethoxy)phenyl]propionic acid



¹H NMR (400 MHz, CDCl₃) δ 1.02-1.35 (m, 5H), 1.56 (br s, 1H), 1.67-1.70 (m, 2H), 1.89 (br s, 2H), 2.41 (s, 3H), 2.60 (t, J= 7 Hz, 2H), 2.91 (t, J= 7 Hz, 2H), 3.09 (t, J= 7 Hz, 2H), 3.39 (s, 3H), 3.47 (br s, 1H), 4.24 (t, J= 7 Hz, 2H), 4.95 (br s, 1H), 5.06 (s, 2H), 6.81 (d, J= 8.3 Hz, 2H), 6.89 (s, 1H), 7.09 (d, J= 7.8 Hz, 1H), 7.23 (t, J= 7.6 Hz, 3H), 7.43 (t, J= 7.8 Hz, 3H), 7.87 (d, J= 8.8 Hz, 2H). MS (ES) *m/e* 612 (M+1).


25

-361-


5

Example 665

3-{2-Methoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



Step A: 3-{2-Methoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl) ethoxy] phenyl}
propionic acid *tert*-butyl ester



10

A 15 mL round bottomed flask was charged with 3-{2-hydroxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester (0.10 g, 0.23 mmol) and methyl iodide (0.21 mL, 2.3 mmol) in anhydrous DMF (1 mL) under a nitrogen atmosphere. The mixture was cooled in an ice bath and treated in one portion with NaH (0.018 g, 0.25 mmol, 60% oil dispersion). The reaction mixture was stirred for 2 h, concentrated, and diluted with EtOAc (40 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and concentrated. The crude product was purified using radial chromatography (EtOAc:hexanes 10:90 to 25:75) to give a yellow oil (0.060 g, 58%). MS (ES) *m/e* 648 (M+1).

20

Step B: 3-{2-Methoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}
propionic acid

A 15 mL round bottomed flask was charged 3-{2-methoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester (0.060 g, 0.13 mmol), CH₂Cl₂ (1.2 mL), and then trifluoroacetic acid (0.6 mL). The solution was stirred at ambient temperature under a nitrogen atmosphere for 16 h and was concentrated. The residue was diluted with CH₂Cl₂ (30 mL) and washed with brine. The organic layer was dried (Na₂SO₄), filtered, and concentrated to give a white solid (0.046 g, 89%). MS (ES) *m/e* 648 (M+1).

-362-

5 The following Examples 666 to 673 are prepared by following a substantially similar procedure as described in Example 665.


10 Example 666

3-{2-Benzylloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester

MS (ES) *m/e* 528 (M+1)

15 Example 667

3-{2-Benzylloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



MS (ES) *m/e* 472 (M+1).


20 Example 668

3-{2-Ethoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester

MS (ES) *m/e* 466 (M+1)

25 Example 669

3-{2-Ethoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



MS (ES) *m/e* 410 (M+1).

-363-

5

Example 670


3-{2-(4-*tert*-Butyl-benzyloxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester

MS (ES) *m/e* 584 (M+1)

10

Example 671

3-{2-(4-*tert*-Butyl-benzyloxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



MS (ES) *m/e* 410 (M+1).

15

Example 672


3-{2-(Biphenyl-4-ylmethoxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester

MS (ES) *m/e* 604 (M+1)

20

Example 673

3-{2-(Biphenyl-4-ylmethoxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid




25 MS (ES) *m/e* 410 (M+1).


5

Example 674

3-{2-*sec*-Butoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



10 Step A: 3-{2-Bromomethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid *tert*-butyl ester



15 A 100 mL round bottomed flask was charged with 3-{2-hydroxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester (1.0 g, 2.29 mmol), anhydrous THF (25 mL), and then triphenylphosphine (1.20 g, 4.57 mmol) and CBr₄ (1.52 g, 4.57 mmol). The yellow mixture was stirred at ambient temperature under a nitrogen atmosphere for 1 h and was poured into EtOAc (100 mL). The organic layer was washed with brine, dried (Na₂SO₄), and concentrated. The crude product was purified using radial chromatography (EtOAc:hexanes 15:85 to 25:75) to give a colorless oil (0.95 g, 83%). MS (ES) *m/e* 501 (M+1).

20

Step B: 3-{2-*sec*-Butoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid

25 A flame-dried 15 mL round bottomed flask was charged with 2-butanol (0.92 mL, 1.0 mmol), anhydrous DMF (1 mL), and then NaH (0.013 g, 0.2 mmol, 60% oil dispersion). The reaction mixture was cooled in an ice bath for 15 min, and 3-{2-bromomethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester ((0.050 g, 0.10 mmol) in anhydrous DMF (0.5 mL) was added. The reaction mixture was stirred under a nitrogen atmosphere for 2 h. A product mixture of free acid and *tert*-butyl ester formed. The reaction was poured into EtOAc (40 mL), washed with

-365-


- 5 brine (3x), dried (Na_2SO_4), and concentrated. The crude oil was then dissolved directly in CH_2Cl_2 (1 mL) and treated with TFA (1.5 mL). The solution was stirred at ambient temperature for 16 h and concentrated. The crude product residue was purified using radial chromatography (EtOAc:hexanes 15:85 to 1:1) to give a yellow oil (0.012 g, 27%). MS (ES) m/e 438 (M+1).

10

The following Examples 675 to 680 are prepared by following a substantially similar procedure as described in Example 674.

Example 675


- 15 3-{2-Isopropoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-phenyl}propionic acid



MS (ES) m/e 424 (M+1).

Example 676

- 20 3-{2-Cyclohexyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



MS (ES) m/e 464 (M+1).


25

-366-

5

Example 677

3-{2-Isobutoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid




MS (ES) *m/e* 438 (M+1).

10

Example 678

3-{2-Cyclohexylmethoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid




MS (ES) *m/e* 478 (M+1).

15

Example 679

3-{2-(Biphenyl-4-yloxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid




20 MS (ES) *m/e* 534 (M+1).

-367-

5

Example 680

3-{2-(3-Methylbutoxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid




MS (ES) *m/e* 452 (M+1).

10

Example 681

3-[4-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-(4-trifluoromethylphenoxy)methyl]phenyl]propionic acid




General procedure for parallel synthesis using the DynaVac Carousel apparatus. A 50 mL glass tube with screw cap and nitrogen inlet was charged with 3-{2-bromomethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid *tert*-butyl ester (0.040 g, 0.080 mmol), 4-trifluoromethylphenol (0.019 g 0.12 mmol), anhydrous DMF (0.5 mL), and then cesium carbonate (0.039 g, 0.12 mmol). The mixture was stirred at ambient temperature for 16 h. MS analysis of the reaction indicated formation of the intermediate ester product [MS (ES) *m/e* 582 (M+1)]. The reaction mixture was poured into ether (30 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and concentrated. The residue was diluted with CH₂Cl₂ (1 mL) and trifluoroacetic acid (0.25 mL) was added. The mixture was stirred at ambient temperature for 1.5 h and was concentrated. The crude product mixture was purified by mass-directed reverse phase HPLC to provide the title compound (0.045 g, 64%). MS (ES) *m/e* 526 (M+1).

-368-

5 The following Examples 682 to 691 are prepared by following a substantially similar procedure as described in Example 681.

Example 682


10 3-{2-(4-Fluorophenoxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



MS (ES) *m/e* 476 (M+1).

Example 683


15 3-{2-(3-Fluorophenoxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



MS (ES) *m/e* 476 (M+1).

Example 684

20 3-{2-(2-Fluorophenoxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



MS (ES) *m/e* 476 (M+1).


25

-369-

5

Example 685

3-{4-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-*p*-tolylloxymethylphenyl}propionic acid




MS (ES) *m/e* 472 (M+1).

10

Example 686


3-{4-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-*m*-tolylloxymethylphenyl}propionic acid



15 MS (ES) *m/e* 472 (M+1).

Example 687

3-{4-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-*o*-tolylloxymethylphenyl}propionic acid



20


MS (ES) *m/e* 472 (M+1).

-370-

5

Example 688

3-{2-(4-Methoxyphenoxy)methyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid




MS (ES) *m/e* 488 (M+1).

10

Example 689


3-{2-(Biphenyl-2-yloxy)methyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid



15 MS (ES) *m/e* 534 (M+1).

Example 690

3-{4-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-phenylsulfanylmethylphenyl}propionic acid



20


MS (ES) *m/e* 474 (M+1).

-371-

5

Example 691

3-{2-Benzenesulfonylmethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid




MS (ES) m/e 506 (M+1).


10

Example 692

3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(4-trifluoromethylphenoxy-methyl)-phenyl]-propionic acid



15 Step A: 3-[2-Bromomethyl-4-(*tert*-butyldiphenylsilanyloxy)phenyl]propionic acid *tert*-butyl ester




20 A 200 mL round bottomed flask was charged with 3-[4-(*tert*-butyldiphenylsilanyloxy)-2-hydroxymethylphenyl]propionic acid *tert*-butyl ester (3.03 g, 6.18 mmol), anhydrous THF (75 mL), and then triphenylphosphine (3.24 g, 12.4 mmol) and CBr₄ (4.10 g, 12.4 mmol). The yellow mixture was stirred at ambient temperature under a nitrogen atmosphere for 1 h and was concentrated. The residue was diluted with EtOAc (500 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and

-372-

- 5 concentrated to a solid. The crude product was purified using a Biotage medium pressure chromatography system (EtOAc:hexanes 10:90) to give a yellow oil (2.95 g, 86%), MS (ES) m/e 572 ($M+NH_4$).


Step B: 3-[4-(*tert*-Butyldiphenylsilyloxy)-2-(4-trifluoromethylphenoxy)methyl]phenyl]propionic acid *tert*-butyl ester

10



- A 50 mL round bottomed flask was charged with 3-[2-bromomethyl-4-(*tert*-butyldiphenylsilyloxy)phenyl]propionic acid *tert*-butyl ester (1.0 g, 1.81 mmol), anhydrous DMF (10 mL), and then 4-trifluoromethylphenol (0.44 g, 2.7 mmol) and cesium carbonate (0.88 g, 2.7 mmol). The reaction mixture was stirred at ambient temperature under a nitrogen atmosphere for 18 h and concentrated. The residue was diluted with EtOAc (100 mL). The organic layer was washed with brine (2x), dried (Na_2SO_4), and concentrated to a yellow oil. The crude residue was purified using radial (EtOAc:hexanes 15:85 to 50:50) to give a yellow oil (0.66 g, 58%). MS (ES) m/e 635 ($M+1$).
- 15
- 20

Step C: 3-[4-Hydroxy-2-(4-trifluoromethylphenoxy)methyl]phenyl]propionic acid *tert*-butyl ester



- A 100 mL round bottomed flask was charged with 3-[4-(*tert*-butyldiphenylsilyloxy)-2-(4-trifluoromethylphenoxy)methyl]phenyl]propionic acid *tert*-
- 25

-373-

5 butyl ester (0.65 g, 1.02 mmol), anhydrous THF (40 mL), and then tetrabutylammonium fluoride (3.1 mL, 3.1 mmol, 1.0 M in THF). The reaction was stirred under a nitrogen atmosphere for 1.5 h and was concentrated. The residue was diluted with EtOAc (100 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and concentrated to an orange oil. The crude oil was purified using radial chromatography (EtOAc:hexanes
10 15:85) to give a white solid (0.31 g, 77%). MS (ES) *m/e* 395 (M-1).

Step D: 3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(4-trifluoromethylphenoxy-methyl)-phenyl]-propionic acid

General procedure for parallel synthesis using the DynaVac Carousel

15 apparatus: A 50 mL glass tube with screw cap and nitrogen inlet was charged with 3-[4-hydroxy-2-(4-trifluoromethylphenoxy-methyl)phenyl]propionic acid *tert*-butyl ester (0.050 g, 0.146 mmol), toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methyloxazol-4-yl)ethyl ester (0.078 g, 0.18 mmol), anhydrous DMF (0.5 mL), and then cesium carbonate (0.072 g, 0.22 mmol). The mixture was stirred at ambient temperature for 16 h. MS
20 analysis of the reaction indicated formation of the ester intermediate MS (ES) *m/e* 658 (M+1). The reaction mixture was poured into ether (30 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and concentrated. The crude residue was diluted with CH₂Cl₂ (2 mL) and then trifluoroacetic acid (1 mL). The mixture was stirred at
25 ambient temperature for 2 h and concentrated under a stream of N₂. The crude product was purified by mass-directed reverse phase HPLC to provide the title compound (0.053 g, 71%). ¹H NMR (250 MHz, CDCl₃) δ 2.41 (s, 3H), 2.70 (t, J= 7.7 Hz, 2H), 2.95-3.05 (m, 4H), 4.28 (t, J= 6.6 Hz, 2H), 5.09 (s, 2H), 6.90 (dd, J= 8.5, 2.8 Hz, 1H), 7.01 (d, J= 2.6 Hz, 1H), 7.06 (d, J= 8.6 Hz, 2H), 7.21 (d, J= 8.5 Hz, 1H), 7.41-7.72 (m, 9H), 8.07 (d, J= 8.4 Hz, 2H). MS (ES) *m/e* 602 (M+1).


30

-374-

5 The following Examples 693 to 697 are prepared by following a substantially similar procedure as described in Example 692.

Example 693


10 3-[4-[2-(2-Cyclohexyl-5-methyloxazol-4-yl)ethoxy]-2-(4-trifluoromethylphenoxy)methyl]phenyl]propionic acid



MS (ES) *m/e* 532 (M+1).

Example 694

15 3-[4-[2-(5-Methyl-2-morpholin-4-yl-thiazol-4-yl)ethoxy]-2-(4-trifluoromethylphenoxy)methyl]phenyl]propionic acid



MS (ES) *m/e* 551 (M+1).


20

-375-

5

Example 695

3-[4-[2-(2-Biphenyl-3-yl-5-methyloxazol-4-yl)ethoxy]-2-(4-trifluoromethylphenoxy)methyl]phenyl]propionic acid




MS (ES) *m/e* 602 (M+1).

10

Example 696

3-[4-{2-[5-Methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}-2-(4-trifluoromethylphenoxy)methyl]phenyl]propionic acid



15

MS (ES) *m/e* 618 (M+1).

Example 697

3-[4-[4-Methyl-2-(4-trifluoromethylphenyl)thiazol-5-ylmethoxy]-2-(4-trifluoromethylphenoxy)methyl]phenyl]propionic acid

20


MS (ES) *m/e* 596 (M+1)

-376-

5


Example 698

3-{2-Benzyloxymethyl-4-[2-(2-cyclohexyl-5-methyl-oxazol-4-yl)-ethoxy]-phenyl}-
propionic acid



Step A: 3-[2-Benzyloxymethyl-4-(tert-butyl-diphenyl-silanyloxy)-phenyl]-propionic acid

10 tert-butyl ester




A 100 mL round bottomed flask was charged with 3-[4-(*tert*-butyl-diphenylsilanyloxy)-2-hydroxymethylphenyl]propionic acid *tert*-butyl ester (1.52 g, 3.10 mmol), anhydrous DMF (15 mL), and then benzyl bromide (1.84 mL, 15.5 mmol) under a nitrogen atmosphere. The solution was cooled to -10°C, and NaH (0.124 g, 3.10 mmol, 60% oil dispersion) was added. The mixture was stirred for 5 h, and was poured into EtOAc (150 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and concentrated to a brown oil. The crude oil was purified using radial chromatography (EtOAc:hexanes 2:98 to 5:95) to give a pale yellow oil (1.04 g, 58%). MS (ES) *m/e* 598 (M+NH₄).

15

20

-377-

5 Step B: 3-(2-Benzyloxymethyl-4-hydroxy-phenyl)-propionic acid *tert*-butyl ester

A 250 mL round bottomed flask was charged with 3-[2-benzyloxymethyl-4-(*tert*-butyl-diphenyl-silanyloxy)-phenyl]-propionic acid *tert*-butyl ester (1.03 g, 1.77 mmol), anhydrous THF (60 mL), and then tetrabutylammonium fluoride (5.3 mL, 5.3 mmol, 1.0 M in THF). The reaction mixture was stirred under a nitrogen atmosphere for 1 h, concentrated, and diluted with EtOAc (100 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and concentrated to give an oil. The crude oil was purified using radial chromatography (EtOAc:hexanes 10:90 to 50:50) to give a white solid (0.52 g, 86%). MS (ES) *m/e* 343 (M-1).

15

Step C: 3-{2-Benzyloxymethyl-4-[2-(2-cyclohexyl-5-methyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid


General procedure for the parallel synthesis of analogs using the DynaVac Carousel apparatus: A 50 mL glass tube with screw cap and nitrogen inlet was charged with 3-(2-benzyloxymethyl-4-hydroxy-phenyl)-propionic acid *tert*-butyl ester (0.050 g, 0.146 mmol), toluene-4-sulfonic acid 2-(2-biphenyl-4-yl-5-methyloxazol-4-yl)ethyl ester (0.078 g, 0.18 mmol), anhydrous DMF (0.5 mL), and then cesium carbonate (0.072 g, 0.22 mmol). The mixture was stirred at ambient temperature for 16 h. MS analysis of the reaction indicated that the intermediate ester was formed, MS (ES) *m/e* 534 (M+1). The crude reaction was poured into ether (30 mL), and washed with brine (2x), dried (Na₂SO₄), and concentrated *in vacuo*. The crude residue was dissolved in CH₂Cl₂ (2 mL) and trifluoroacetic acid was added (1.0 mL). The reaction mixture was stirred at ambient temperature for 2 h and was concentrated under a stream of N₂. The crude product was purified by mass-directed reverse phase HPLC to provide 0.053 g (71%) of 3-{4-[2-(2-Cyclohexyl-5-methyloxazol-4-yl)ethoxy]-2-phenoxyethylphenyl}propionic acid. MS (ES) *m/e* 478 (M+1).

-378-

5 The following Examples 699 to 704 are prepared by following a substantially similar procedure as described in Example 698.

10 Example 699


3-{2-Benzylloxymethyl-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl}propionic acid



MS (ES) *m/e* 497 (M+1).

15 Example 670


3-{2-Benzylloxymethyl-4-[2-(2-biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]phenyl}propionic acid



MS (ES) *m/e* 548 (M+1).

20 Example 671

3-{2-Benzylloxymethyl-4-[2-(2-biphenyl-3-yl-5-methyloxazol-4-yl)ethoxy]phenyl}propionic acid




MS (ES) *m/e* 548 (M+1).

-379-

5

Example 672

3-(2-Benzylloxymethyl-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl)propionic acid




MS (ES) *m/e* 564 (M+1).

10

Example 673


3-(2-Benzylloxymethyl-4-{2-[2-(4-butoxyphenyl)-5-methyloxazol-4-yl]ethoxy}phenyl)propionic acid



15 MS (ES) *m/e* 544 (M+1).

Example 674

3-(2-Benzylloxymethyl-4-{2-[5-methyl-2-(4-trifluoromethylphenyl)oxazol-4-yl]ethoxy}phenyl)propionic acid



20


MS (ES) *m/e* 540 (M+1).

-380-


5

Example 705

3-{4-[3-(Biphenyl-4-yloxy)-propoxy]-2-cyclohexylcarbamoyloxymethyl-phenyl}-
propionic acid



10 Step A: 3-{4-[3-(Biphenyl-4-yloxy)-propoxy]-2-cyclohexylcarbamoyloxymethyl-phenyl}-propionic acid *tert*-butyl ester



A mixture of 3-(2-cyclohexylcarbamoyloxymethyl-4-hydroxy-phenyl)-propionic acid *tert*-butyl ester (50 mg, 0.13 mmol; Example 652 Step D), 4-(3-bromopropoxy)-biphenyl (58 mg, 0.2 mmol; *Tetrahedron* 1994, 50, 3427, and potassium carbonate (53 mg, 0.39 mmol) in acetonitrile (6 mL) was heated at 80°C overnight. Ethyl acetate (20 mL) and H₂O (20 mL) were added. The organic layer was separated, and the aqueous layer was extracted with ethyl acetate (3x). The combined organic layers were dried (MgSO₄), filtered, and concentrated. The crude mixture was purified by silica gel column chromatography (hexane/ethyl acetate 4/1) to give the title compound (50 mg, 65%).

¹H-NMR (300 MHz, CDCl₃): δ 7.54 (m, 4H), 7.42 (t, 2H, *J*=7.6), 7.32 (d, 1H, *J*=8.4), 7.12 (d, 1H, *J*=8.3), 6.99 (d, 2H, *J*=8.7), 6.93 (d, 1H, *J*=2.2), 6.82 (dd, 1H, *J*=8.4, 2.7), 5.09 (s, 2H), 4.74 (d, 1H, *J*=7.5), 4.18 (m, 4H), 3.56-3.42 (m, 1H), 2.90 (t, 2H, *J*=7.8), 2.48 (t, 2H, *J*=7.8), 2.28 (qn, 2H, *J*=6.0), 1.95-1.88 (m, 2H), 1.75-1.08 (m, 8H), 1.44 (s, 9H).

25

-381-


- 5 Step B: 3-{4-[3-(Biphenyl-4-yloxy)-propoxy]-2-cyclohexylcarbamoyloxymethyl-phenyl}-propionic acid

Trifluoroacetic acid (0.032 mL, 0.42 mmol) was added to a solution of 3-{4-[3-(biphenyl-4-yloxy)-propoxy]-2-cyclohexylcarbamoyloxymethyl-phenyl}-propionic acid *tert*-butyl ester (50 mg, 0.085 mmol) in CH₂Cl₂ (5 mL) at room temperature. The reaction mixture was stirred overnight and concentrated under vacuum to give the title compound (41 mg, 90%). ¹H-NMR (200 MHz, CDCl₃): δ 9.4 (br s, 1H), 7.58-7.49 (m, 4H), 7.42 (m, 2H), 7.32 (m, 1H), 7.13 (d, 1H, *J*=8.3), 7.00-6.93 (m, 3H), 6.84 (dd, 1H, *J*=7.8, 2.7), 5.11 (s, 2H), 4.76 (br s, 1H), 4.17 (m, 4H), 3.57-3.35 (m, 1H), 2.95 (t, 2H, *J*=7.8), 2.63 (t, 2H, *J*=7.8), 2.27 (qn, 2H, *J*=6.0), 1.99-1.05 (m, 10H).


15

Example 706

3-{4-[3-(4-Benzoyl-phenoxy)-propoxy]-2-cyclohexylcarbamoyloxymethyl-phenyl}-propionic acid



- 20 Step A: 3-{4-[3-(4-Benzoyl-phenoxy)-propoxy]-2-cyclohexylcarbamoyloxymethyl-phenyl}-propionic acid *tert*-butyl ester



- A mixture of 3-(2-cyclohexylcarbamoyloxymethyl-4-hydroxy-phenyl)-propionic acid *tert*-butyl ester (50 mg, 0.13 mmol; Example 652 Step D), [4-(3-bromopropoxy)-phenyl]-phenyl-methanone (64 mg, 0.2 mmol, *Bull. Chem. Soc. Jpn.* 1990, 63, 1342) and potassium carbonate (53 mg, 0.39 mmol) in acetonitrile (6 mL) was heated at 80°C overnight. Ethyl acetate (20 mL) and H₂O (20 mL) were added. The organic layer was separated, and the aqueous layer was extracted with ethyl acetate (3x). The

-382-

5 combined organic layers were dried (MgSO_4), filtered, and concentrated. The crude mixture was purified by silica gel column chromatography (hexane/ethyl acetate 4/1) to give the title compound (50 mg, 62%). $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.74 (d, 2H, $J=8.5$), 7.68 (d, 2H, $J=7.7$), 7.48 (d, 1H, $J=8.4$), 7.40 (m, 2H), 7.05 (d, 1H, $J=8.5$), 6.90 (d, 2H, $J=8.6$), 6.85 (s, 1H), 6.74 (dd, 1H, $J=8.5$, 2.2), 5.04 (s, 2H), 4.68 (d, 1H, $J=7.3$),
10 4.17 (t, 2H, $J=5.9$), 4.09 (t, 2H, $J=5.9$), 3.43 (m, 1H), 2.82 (t, 2H, $J=7.8$), 2.40 (t, 2H, $J=7.8$), 2.21 (qn, 2H, $J=5.9$), 1.85 (m, 2H), 1.67-1.02 (m, 8H), 1.36 (s, 9H).

Step B: 3-{4-[3-(4-Benzoyl-phenoxy)-propoxy]-2-cyclohexylcarbamoyloxymethyl-phenyl}-propionic acid


15 Trifluoroacetic acid (0.031 mL, 0.4 mmol) was added to a solution of 3-{4-[3-(4-benzoyl-phenoxy)-propoxy]-2-cyclohexylcarbamoyloxymethyl-phenyl}-propionic acid tert-butyl ester (50 mg, 0.08 mmol) in CH_2Cl_2 (5 mL) at room temperature. The reaction mixture was stirred overnight and concentrated under vacuum to give the title compound. (39 mg, 88%). $^1\text{H-NMR}$ (200 MHz, CDCl_3): δ 7.83-7.72 (m, 4H), 7.57-
20 7.43 (m, 3H), 7.12 (d, 1H, $J=8.6$), 6.98-82 (m, 4H), 5.10 (s, 2H), 4.82 (br s, 1H), 4.23 (t, 2H, $J=5.9$), 4.15 (t, 2H, $J=5.9$), 3.57-3.36 (m, 1H), 2.94 (m, 2H), 2.62 (m, 2H), 2.28 (qn, 2H, $J=5.9$), 1.98-1.02 (m, 10H).

-383-


5

Example 708

2-{2-Methoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid




10 Step A: 2-(4-Benzyloxy-2-formyl-phenoxy)-2-methyl-propionic acid ethyl ester



5-Benzyloxy-2-hydroxy-benzaldehyde (Kappe, T.; Witoszynskyj, T. *Arch. Pharm.*, 1975, 308 (5), 339-346) (169 g, 741 mmol), ethyl bromoisobutyrate (164 mL, 1.11 mol), and cesium carbonate (240 g, 741 mmol) in dry DMF (600 mL) were heated at 80 °C for 15 h. Additional cesium carbonate (5 g) and ethyl bromoisobutyrate (20 mL) were added, and the reaction mixture was heated for 6 h. The reaction mixture was cooled, diluted with EtOAc (4 L), and washed with water (3 x 2 L). The organic layer was dried (Na₂SO₄) and concentrated to a brown oil. The crude product was recrystallized from EtOAc (150 mL) with hexanes until turbid to give the title compound as a pale yellow solid (210 g, 83%): mp 65 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.24 (t, 3H, J = 7.1 Hz), 1.62 (s, 6H), 4.23 (q, 2H, J = 7.1 Hz), 6.81 (d, 1H, J = 8.8 Hz), 7.10 (dd, 1H, J = 4.6, 9.0 Hz), 7.30-7.43 (m, 6H); MS (ES) m/e 343.1 [M+1].

Step B: 2-(4-Hydroxy-2-hydroxymethyl-phenoxy)-2-methyl-propionic acid ethyl ester



25


2-(4-Benzyloxy-2-formyl-phenoxy)-2-methyl-propionic acid ethyl ester (185 g, 540 mmol) in ethanol (700 mL) was treated with 10% Pd/C (205 g) and hydrogen

-384-

5 (60 psi) at 50 °C for 2 d. The mixture was filtered through Celite, washed with ethanol (1.5 L), and concentrated. The residue was recrystallized from EtOAc/hexanes to give the title compound (116 g). ¹H NMR (400 MHz, CDCl₃) δ 1.26 (t, 3H, *J* = 7.3 Hz), 1.52 (s, 6H), 2.14 (s, 3H), 4.23 (q, 2H, *J* = 7.3 Hz), 4.59 (brs, 2H), 6.61-6.68 (m, 2H), 6.77 (d, 1H, *J* = 2.8 Hz).


10

Step C: 2-{2-hydroxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid ethyl ester



A 25 mL round bottomed flask under a nitrogen atmosphere was charged with toluene-4-sulfonic acid 2-(5-methyl-2-phenyloxazol-4-yl)ethyl ester (0.77 g, 2.17 mmol), 2-(4-hydroxy-2-hydroxymethylphenoxy)-2-methylpropionic acid ethyl ester (0.5 g, 1.97 mmol), and absolute ethanol (10 mL). Potassium carbonate (0.54 g, 3.94 mmol, 325 mesh) was added, and the reaction was heated to 80° C for 12 h. The mixture was concentrated and the crude residue was diluted with EtOAc (75 mL). The organic layer was washed with brine (2x), dried (Na₂SO₄), and concentrated to an oil. The crude product was purified using radial chromatography (EtOAc:Hex 5:95 to 35:65) to give a colorless oil (0.17 g, 20%). MS (ES) *m/e* 452 (M+1).

Step D: 2-{2-Methoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid ethyl ester



A 15 mL round bottomed flask under a nitrogen atmosphere were charged with 2-{2-hydroxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid ethyl ester (0.075 g, 0.17 mmol), anhydrous DMF (1 mL), and then methyl iodide (0.16 mL, 1.7 mmol). The solution was cooled in an ice bath and was

-385-

5 treated with NaH (0.014 g, 0.34 mmol, 60% oil dispersion). The mixture was stirred for 2 h, poured into EtOAc (6 mL) and brine (10 mL), and acidified using dilute sulfuric acid. The organic layer was separated, dried (Na₂SO₄), and concentrated to an oil. The crude product was purified using radial chromatography (EtOAc:Hex 15:85) to give a colorless oil (0.039 g, 51%). MS (ES) *m/e* 454 (M+1).

10


Step E: 2-{2-Methoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid

A 25 mL round bottomed flask was charged with 2-{2-methoxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid ethyl ester (0.039 g, 0.087 mmol), ethanol (2 mL), and then aqueous 2N NaOH (0.22 mL, 0.44 mmol). The solution was heated at 55° C for 1 h. The mixture was concentrated, acidified using 5% H₂SO₄ (1.5 mL), and partitioned between CH₂Cl₂ (15 mL) and brine (15 mL). The organic layer was dried (Na₂SO₄), filtered, and concentrated and to give a white solid (0.024 g, 66%). MS (ES) *m/e* 426 (M+1).

20

Example 709

2-{2-Benzoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)-ethoxy]phenoxy}-2-methylpropionic acid




25 The above compound is prepared by following a substantially similar procedure as described in Example 708. MS (ES) *m/e* 502 (M+1).

-386-


5

Example 710

2-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid



Step A: 3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}-propionic acid ethyl ester



A 15 mL round bottomed flask under N₂ was charged 2-{2-hydroxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid ethyl ester (0.075 g, 0.17 mmol), cyclohexylisocyanate (0.13 mL, 1.0 mmol), anhydrous CH₂Cl₂ (0.5 mL), and then 1.0 N HCl in ether (0.086 mL, 0.086 mmol). The reaction mixture was stirred at ambient temperature for 18 h and was diluted with CH₂Cl₂ (10 mL). The organic layer was washed with brine, dried (Na₂SO₄) and concentrated to give 0.10 g of a crude oil, which was used directly in the next step. MS (ES) *m/e* 564 (M+1).

20

Step B: 3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}-propionic acid

To a 15 mL round bottomed flask was charged with 3-{2-cyclohexylcarbamoyloxymethyl-4-[2-(2-phenyloxazol-4-yl)ethoxy]phenyl}propionic acid ethyl ester (0.10 g, 0.17 mmol), ethanol (2 mL), and then 2N NaOH (0.48 mL, 0.96 mmol). The solution was heated at 55°C for 2 h. The mixture was concentrated, acidified using 5% H₂SO₄ (1.5 mL), and partitioned between CH₂Cl₂ (15 mL) and brine (15 mL).

25


-387-

- 5 The organic layer was dried (Na_2SO_4), filtered, and concentrated and was concentrated. The crude residue was submitted for mass-directed HPLC purification to give a white solid (0.058 g, 63%) . MS (ES) m/e 537 (M+1).

10 The following Examples 711 to 713 are prepared by following a substantially similar procedure as described in Example 710.

Example 711

2-{2-Isopropylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid




15

MS (ES) m/e 497 (M+1).

Example 712

20 2-{2-Benzylcarbamoyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid




MS (ES) m/e 545 (M+1).

-388-

5

Example 713

2-{2-(4-Fluorobenzylcarbamoyloxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid




MS (ES) m/e 563 (M+1).


10

Example 714

2-Methyl-2-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-m-tolyloxymethyl-phenoxy}-propionic acid



15 Step A: 2-{2-Bromomethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid ethyl ester



A 100 mL round bottomed flask was charged with 2-{2-hydroxymethyl-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenoxy}-2-methyl-propionic acid ethyl ester (1.0 g, 2.25 mmol), dissolved in anhydrous THF (75 mL), and then of triphenylphosphine (1.18 g, 4.50 mmol) and CBr_4 (1.49 g, 4.50 mmol). The mixture was stirred at ambient temperature under a nitrogen atmosphere for 1 h and was poured into EtOAc (135 mL). The organic layer was washed with brine (50 mL), dried (Na_2SO_4), and concentrated. The crude product was purified using radial chromatography (EtOAc:hexanes 15:85) to give a the title compound as a colorless oil (0.95 g, 76%).

- 5 Step B: 2-Methyl-2-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-*m*-tolylloxymethyl-phenoxy}-propionic acid


General procedure for the parallel synthesis of analogs using the DynaVac Carousel apparatus: A 50 mL glass tube with screw cap and nitrogen inlet was charged with 2-{2-bromomethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methyl-
10 propionic acid ethyl ester (0.040 g, 0.080 mmol), *m*-cresol (0.012 mL, 0.12 mmol), absolute ethanol (1 mL), and then potassium carbonate (0.022 g, 0.16 mmol; 325 mesh). The mixture was heated to 80° C for 4 h. MS analysis of the reaction indicated that 2-methyl-2-{4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-*m*-tolylloxymethylphenoxy}-propionic acid ethyl ester: MS (ES) *m/e* 530 (M+1) had formed. The reaction mixture
15 was treated with 2N NaOH (0.4 mL), heated at 55°C for 3 h, cooled, and concentrated. The residue was treated with 5N HCl (0.75 mL) and CH₂Cl₂ (1 mL) and was poured onto a 3-mL ChemElute cartridge to remove the aqueous layer. The cartridge was washed with CH₂Cl₂, and the solvent was removed *in vacuo*. The crude residue was purified by mass-directed reverse phase HPLC to provide the title compound (0.032 g, 38%). MS
20 (ES) *m/e* 502 (M+1).

The following Examples 715 to 723 are prepared by following a substantially similar procedure as described in Example 714.

25

Example 715

2-{2-(4-Fluorophenoxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid




MS (ES) *m/e* 506 (M+1).

30

Example 716

2-{2-(3-Fluorophenoxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid

-390-




5

MS (ES) *m/e* 506 (M+1).Example 717


2-{2-(2-Fluorophenoxy)methyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid

10

MS (ES) *m/e* 506 (M+1).Example 718

2-Methyl-2-{4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-*p*-tolylloxymethylphenoxy}propionic acid

15

MS (ES) *m/e* 502 (M+1).


20

-391-

5

Example 719

2-Methyl-2-{4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-*o*-tolylloxymethylphenoxy}propionic acid




MS (ES) *m/e* 502 (M+1).

10

Example 720


2-{2-(4-Methoxyphenoxy)methyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid



15 MS (ES) *m/e* 518 (M+1).

Example 721

2-Methyl-2-[4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]-2-(4-trifluoromethylphenoxy)methylphenoxy]propionic acid



20


MS (ES) *m/e* 556 (M+1).

-392-

5

Example 722

2-{2-(Biphenyl-2-yloxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid




MS (ES) *m/e* 564 (M+1).

10

Example 723


2-{2-(Biphenyl-4-yloxymethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy}-2-methylpropionic acid



15 MS (ES) *m/e* 564 (M+1).

Example 724

3-{2-(Benzoylamino-methyl)-4-[3-(biphenyl-4-yloxy)propoxy]phenyl} propionic acid



20

A solution of 4-(3-bromo-propoxy)-biphenyl (291 mg, 1.00 mmol; Preparation 12) and 3-[2-(benzoylamino-methyl)-4-hydroxy-phenyl]-propionic acid tert-


-393-

- 5 butyl ester (320 mg, 0.90 mmol; Preparation 21) in DMF (5 mL) was treated with K_2CO_3 (100 mg) and heated at 60°C for 48 h. The reaction mixture was diluted with water (50 mL) and extracted with EtOAc (3 x 20 mL). The combined organics were dried (Na_2SO_4), concentrated, and purified using silica gel chromatography (10-30% EtOAc/hexanes) to give 3-{2-(benzoylaminomethyl)-4-[3-(biphenyl-4-yloxy)propoxy]phenyl} propionic acid *tert*-butyl ester (450 mg). This material was treated with a mixture of CH_2Cl_2 (1.0 mL)/TFA (0.8 mL)/water (0.1 mL) at ambient temperature for 2 h. The reaction mixture was concentrated and dried under vacuum to afford the title product as a white solid (380 mg, 75%). MS (ES) m/e 510.1 (M+1).

- 15 The following Examples 725 to 757 are prepared by following a substantially similar procedure as described in Example 724.

Example 725


3-{2-(Benzoylaminomethyl)-4-[2-(4-phenoxy-phenoxy)ethoxy]phenyl} propionic acid



MS [ES] m/z 512 (M+1).

Example 726

3-{2-(Benzoylaminomethyl)-4-[2-(3-phenylbenzofuran-6-yloxy)ethoxy]-phenyl}-propionic acid




MS [ES] m/z 536 (M+1).

-394-

5

Example 727

3-{2-(Benzoylaminoethyl)-4-[2-(6-methoxynaphthalen-2-yloxy)ethoxy]phenyl}
propionic acid




MS [ES] m/z 500 (M+1).

10

Example 728


3-{2-[(Cyclobutanecarbonylamino)methyl]-4-[2-(4-phenoxyphenoxy)-ethoxy]phenyl}
propionic acid



15 MS [ES] m/z 490 (M+1).

Example 729

3-[4-[2-(Biphenyl-4-yloxy)ethoxy]-2-(isopropoxycarbonylaminoethyl)-phenyl]
propionic acid



20


MS [ES] m/z 473 (M+1).

-395-

5

Example 730

3-(4-[2-(Biphenyl-4-yloxy)ethoxy]-2-[[pyridine-2-carbonyl]amino]methyl}-phenyl)
propionic acid




MS [ES] m/z 497 (M+1).

10

Example 731


3-(4-[2-(Biphenyl-3-yloxy)ethoxy]-2-[[pyridine-2-carbonyl]amino]methyl}
phenyl)propionic acid



15 MS [ES] m/z 497 (M+1).

Example 732

3-(4-[2-(4-Phenoxy-phenoxy)ethoxy]-2-[[pyridine-2-carbonyl]amino]-methyl}phenyl)
propionic acid



20


MS [ES] m/z 501 (M+1).

-396-

5

Example 733

3-(4-[2-(3-Phenylbenzofuran-6-yloxy)ethoxy]-2-[[[(pyridine-2-carbonyl)-amino]methyl]phenyl] propionic acid




MS [ES] m/z 537 (M+1).

10

Example 734


3-(4-[2-(6-Methoxynaphthalen-2-yloxy)ethoxy]-2-[[[(pyridine-2-carbonyl)-amino]methyl]phenyl] propionic acid



15 MS [ES] m/z 501 (M+1).

Example 735

3-{2-(Benzoylaminoethyl)-4-[4-(biphenyl-4-yloxy)butoxy]phenyl} propionic acid:




20 MS [ES] m/z 524 (M+1).

-397-

5

Example 736

3-{2-(Benzoylaminoethyl)-4-[4-(biphenyl-3-yloxy)butoxy]phenyl} propionic acid:




MS [ES] m/z 524 (M+1).

10


Example 737

3-{2-(Benzoylaminoethyl)-4-[4-(4-phenoxyphenoxy)butoxy]phenyl} propionic acid:



MS [ES] m/z 540 (M+1).

15

Example 7383-{2-(Benzoylaminoethyl)-4-[4-(3-phenylbenzofuran-6-yloxy)butoxy]phenyl}-
propionic acid

MS [ES] m/z 564 (M+1).


20

-398-

5

Example 739

3-{2-(Isopropoxycarbonylaminomethyl)-4-[4-(4-phenoxyphenoxy)butoxy]phenyl}
propionic acid




MS [ES] m/z 522 (M+1).

10

Example 740


3-{2-(Isopropoxycarbonylaminomethyl)-4-[4-(3-phenylbenzofuran-6-yloxy)butoxy]phenyl} propionic acid



15 MS [ES] m/z 546 (M+1).

Example 741

3-(4-[4-(Biphenyl-3-yloxy)butoxy]-2-[[pyridine-2-carbonyl]amino]methyl)-phenyl
propionic acid



20


MS [ES] m/z 525 (M+1).

-399-

5

Example 742

3-(4-[4-(4-Phenoxyphenoxy)butoxy]-2-[[[(pyridine-2-carbonyl)amino]-methyl}phenyl)
propionic acid




MS [ES] m/z 541 (M+1).

10

Example 743


3-(4-[4-(3-Phenylbenzofuran-6-yloxy)butoxy]-2-[[[(pyridine-2-carbonyl)-amino]methyl}
phenyl) propionic acid



15 MS [ES] m/z 565 (M+1).

Example 744

3-(4-[4-(6-Methoxynaphthalen-2-yloxy)butoxy]-2-[[[(pyridine-2-carbonyl)-
amino]methyl}phenyl) propionic acid



20


MS [ES] m/z 529 (M+1).

-400-

5

Example 745

3-(4-[4-(Biphenyl-3-yloxy)butoxy]-2-[[2,5-dichlorothiophene-3-carbonyl-
amino]methyl]phenyl) propionic acid




MS [ES] m/z 599 (M+1).

10

Example 746

3-{2-(Benzoylaminomethyl)-4-[3-(biphenyl-3-yloxy)propoxy]phenyl} propionic acid:




MS [ES] m/z 510 (M+1).

15

Example 747

3-[4-[3-(Biphenyl-3-yloxy)propoxy]-2-(isopropoxycarbonylaminomethyl)-phenyl]
propionic acid




20 MS [ES] m/z 492 (M+1).

-401-

5

Example 748

3-(4-[3-(Biphenyl-4-yloxy)propoxy]-2-[[pyridine-2-carbonyl]amino]methyl}-phenyl)
propionic acid




MS [ES] m/z 511 (M+1).

10

Example 749


3-(4-[3-(Biphenyl-3-yloxy)propoxy]-2-[[pyridine-2-carbonyl]amino]methyl} phenyl)
propionic acid



15 MS [ES] m/z 511 (M+1).

Example 750

3-(4-[3-(6-Methoxynaphthalen-2-yloxy)propoxy]-2-[[pyridine-2-carbonyl]-
amino]methyl}phenyl) propionic acid



20


MS [ES] m/z 515 (M+1).

-402-

5

Example 751

3-(4-[3-(Biphenyl-4-yloxy)propoxy]-2-[[2,5-dichlorothiophene-3-carbonyl]-amino]methyl}phenyl) propionic acid




MS [ES] m/z 585 (M+1).

10

Example 752

3-{2-[[2,5-Dichlorothiophene-3-carbonyl]amino]methyl}-4-[3-(3-phenyl-benzofuran-6-yloxy)propoxy]phenyl} propionic acid




15 MS [ES] m/z 625 (M+1).

-403-

5

Example 753

3-[4-[3-(Biphenyl-4-yloxy)propoxy]-2-(1,3-dioxo-1,3-dihydroisoindol-2-ylmethyl)phenyl] propionic acid




MS [ES] m/z 536 (M+1).

10

Example 754


3-[4-[3-(Biphenyl-4-yloxy)propoxy]-2-(isopropoxycarbonylaminomethyl)-phenyl] propionic acid



15 MS [ES] m/z 492 (M+1).

Example 755

3-{2-(Benzoylaminomethyl)-4-[3-(4-phenoxyphenoxy)propoxy]phenyl} propionic acid




20 MS [ES] m/z 526 (M+1).

-404-

5

Example 756

3-{2-(Isopropoxycarbonylaminoethyl)-4-[3-(4-phenoxyphenoxy)propoxy]-phenyl}
propionic acid




MS [ES] m/z 508 (M+1).

10

Example 757


3-(4-[3-(4-Phenoxyphenoxy)propoxy]-2-[(pyridine-2-carbonyl)amino]-methyl}phenyl}
propionic acid



15 MS [ES] m/z 527 (M+1).


Example 758

(2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-acetic acid methyl ester



20


Step 1: 4-Bromo-3-oxo-pentanoic acid methyl ester



25 A solution of bromine (28.3 g, 0.177 mol) in chloroform (30 mL) was added dropwise over 2 h to a solution of methyl propionylacetate (23.5 g, 0.177 mol) in chloroform (155 mL) at 0-5 °C. The mixture was stirred for 30 min, and the cooling bath


- 5 was removed. The mixture was stirred for 18 h, and ice water (200 mL) was added. The organic layer was collected and washed with cold water (2 x 200 mL), 10% aqueous sodium thiosulfate (2 x 200 mL) and brine (200 mL). The filtered solution was dried (Na_2SO_4) and concentrated to 36.5 g of the title compound as a clear liquid.

- 10 Step 2: Biphenyl-4-carboxylic acid 3-methoxycarbonyl-1-methyl-2-oxo-propyl ester



- A mixture of biphenyl-4-carboxylic acid (800 g, 4.03 mol) in acetone (4.6 L) was treated with triethylamine (0.6 L, 4.3 mol, 1.07eq) dropwise over 13 min while
15 maintaining the temperature at 15-30 °C. 4-Bromo-3-oxo-pentanoic acid methyl ester (880 g, 4.21 mol, 1.04 eq) was added dropwise over 21 min at 15-30 °C. The mixture was stirred overnight at room temperature. Water (9.6 L) was added dropwise over 85 minutes at 15-30 °C. The mixture was stirred for 2 h. The precipitated product was collected by filtration and washed twice with water (1 L). The product was dried under
20 vacuum at 50 °C to afford 1291 g (98% yield, 96% HPLC purity) of the title compound.

- Step 3: Biphenyl-4-carboxylic acid 2-amino-3-methoxycarbonyl-1-methyl-allyl ester



- A mixture of biphenyl-4-carboxylic acid 3-methoxycarbonyl-1-methyl-2-oxo-propyl ester (1275 g, 3.9 mol, 1 eq) and ammonium acetate (640 g, 8.3 mol) in
25 ethanol (10 L) was heated with stirring at 70-75 °C until the keto ester compound is completely consumed (1-2 h). The mixture was then kept at 0-5 °C for 1.5 h. The precipitated solid was collected by filtration and washed with hexanes (2.5 L). The product was dried overnight under vacuum at 50 °C to obtain 1244 g (90% yield, 98%
30 HPLC purity) of the title compound.

- Step 4: (2-Biphenyl-4-yl-5-methyl-oxazol-4-yl)-acetic acid methyl ester

A mixture of biphenyl-4-carboxylic acid 2-amino-3-methoxycarbonyl-1-methyl-allyl ester (566 g, 1.74 mol, 1 eq) and ammonium acetate (283 g, 3.67 mol) in glacial

-406-


5 acetic acid (11.3 L) was heated at reflux for 2 h, cooled, and concentrated. The residue was coevaporated with toluene (2 x 2.5 L) and EtOAc (2.5 L). The mixture was diluted with EtOAc (6.6 L) and transferred with EtOAc (2.2 L) to a bottom outlet separation flask. The mixture was washed twice with water (2.2 L), saturated aqueous NaHCO₃ (1.1 L), and brine (2 x 2.2 L). The organic layer was dried (Na₂SO₄, 550 g), filtered with the aid of EtOAc (1.1 L), and concentrated. The residue was dissolved in isopropanol (2 L) at 50°C and allowed to cool overnight at room temperature. The mixture was kept at 0°C to 5°C for 1 h. The precipitated solid was broken up and collected by filtration. The solid was washed with cold isopropanol (4 x 0.55 L) and dried overnight in a vacuum oven at 50°C to yield 428 g of the title compound (72% from Step 2).

15

-407-

5 WHAT IS CLAIMED IS:

1. A compound of formula I,



and pharmaceutically acceptable salts, solvates, hydrates or stereoisomers thereof,
wherein:

- 10
- n^1
- is 2, 3, 4 or 5;

V is a bond or O;

X is CH_2 or O;

p is 0 or 1;

m is 1-4;

15

Y^1 is: $Y^{1a} - \text{Ar} -$ wherein,



is: aryl or heteroaryl,

wherein aryl and heteroaryl are optionally substituted with one or more groups

20

independently selected from the group consisting of:

hydrogen, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, halo, haloalkyl and haloalkyloxy; Y^{1a} is: hydrogen, $(\text{C}_0\text{-C}_3)\text{alkyl-aryl}$,

25

 C(O)-aryl ,

heteroaryl,

cycloalkyl,

heterocycloalkyl,

aryloxy,

30

 $\text{NR}^5(\text{CH}_2)_m\text{OR}^5$,

-408-

- 5 aryl-Z-aryl,
 aryl-Z-heteroaryl,
 aryl-Z-cycloalkyl,
 aryl-Z-heterocycloalkyl,
 heteroaryl-Z-aryl,
10 heteroaryl-Z-heterocycloalkyl or
 heterocycloalkyl-Z-aryl,
 wherein aryl, cycloalkyl, aryloxy, heteroaryl, and heterocycloalkyl are optionally
 substituted with one or more substituents independently selected from the group
 consisting of:
- 15 halo,
 hydroxyl,
 nitro,
 cyano,
 C₁-C₆ alkyl,
20 C₁-C₆ alkoxy optionally substituted with N(R⁵)₂,
 haloalkyl,
 N(R⁵)₂,
 N[C(O)R⁵]₂,
 N[S(O)₂R⁵]₂,
25 NR⁵S(O)₂R⁵,
 NR⁵C(O)R⁵,
 NR⁵C(O)OR⁵,
 C(O)N(R⁵)₂,
 C(O)OR⁵ and
30 C(O)R⁵;

- Z is: a bond,
 -oxygen-
 -C(O)NR⁵-
35 -NR⁵C(O)-,
 -NR⁵C(O)O-,

-409-

- 5 -C(O)-,
 -NR⁵-,
 -[O]_p(CH₂)_m-,
 -(CH₂)_m[O]_p-,
 -NR⁵(CH₂)_m- or
 10 -(CH₂)_mNR⁵-;

Y² and Y³ are each independently:

- hydrogen,
 C₁-C₆ alkyl or
 15 C₁-C₆ alkoxy;

- Y⁴ is: (C₁-C₃)alkyl-NR⁵C(O)-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-NR⁵C(O)-(C₂-C₅)alkenyl-Y⁷,
 (C₁-C₃)alkyl-NR⁵C(O)-(C₂-C₅)alkynyl-Y⁷;
 20 (C₁-C₃)alkyl-NR⁵C(O)O-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-NR⁵C(O)NR⁵-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-NR⁵C(S)NR⁵-(C₀-C₅)alkyl-Y⁷,
 (C₀-C₃)alkyl-C(O)NR⁵-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-OC(O)NY¹⁰Y¹¹,
 25 (C₁-C₃)alkyl-NY¹⁰Y¹¹,
 (C₁-C₃)alkyl-O-(C₀-C₅)alkyl-Y⁷,
 (C₁-C₃)alkyl-S-(C₀-C₅)alkyl-Y⁷ or
 CN;

- 30 Y⁷ is: hydrogen,
 aryl,
 heteroaryl,
 C₁-C₁₂ alkyl,
 C₁-C₆ alkoxy,
 35 cycloalkyl,
 heterocycloalkyl,

-410-

5 aryloxy,
C(O)-heteroaryl or
SR⁶,
wherein alkyl, aryl, aryloxy, alkoxy, heteroaryl, cycloalkyl, and heterocycloalkyl
are optionally substituted with one or more groups independently selected from
10 R⁷:

Y¹⁰ and Y¹¹ are each independently:
hydrogen,
aryl,
15 heteroaryl,
C₁-C₁₀ alkyl,
cycloalkyl,
SO₂(R⁶); or
Y¹⁰ and Y¹¹ together are a 5- to 10-membered heterocycloalkyl ring or
20 heterocycloalkyl ring fused with aryl, and the heterocycloalkyl ring optionally
containing one or more heteroatoms selected from N, O or S; and wherein,
aryl, heteroaryl, heterocycloalkyl and alkyl are optionally substituted with one or
more substituents independently selected from R⁷;

25 R⁵ is: hydrogen or C₁-C₆ alkyl;


R⁶ is: hydrogen,
C₁-C₁₀ alkyl,
cycloalkyl,
30 aryl, or
heteroaryl,
wherein alkyl, cycloalkyl, aryl and heteroaryl are optionally substituted with one
or more substituents independently selected from R⁷;

35 R⁷ is: halo,
nitro,
oxo,
cyano,

-411-

5. hydroxyl,
benzyl,
phenyl,
phenoxy,
heteroaryl,
10 C(O)R⁶,
C₁-C₁₀ alkyl,
C₁-C₆ alkoxy,
C₁-C₆ haloalkyl,
C₁-C₆ haloalkyloxy,
15 O(CH₂)_m-phenyl,
(CH₂)_mOC(O)-aryl,
C(O)OR⁵,
S(O)₂R⁵,
S(O)₂N(R⁵)₂,
20 SR⁵ or
N(R⁵)₂,
wherein phenyl and phenoxy are optionally substituted with one or more groups
independently selected from halo or trifluoromethyl.

2. A compound represented by the following structural formula Ia:



Ia

or a pharmaceutically acceptable salt, hydrate or solvate thereof, wherein:

- Y¹ is an unsubstituted or substituted group selected from the group consisting of:
aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aryl-C₁-C₄ alkyl, heteroaryl-C₁-C₄ alkyl,
30 cycloalkyl-C₁-C₄ alkyl, and t-butyl;

Y² is selected from the group consisting of:

H, C₁-C₁₀ alkyl, cycloalkyl, (C₁-C₁₀ alkyl)-Y⁵, O-Y⁶;

-412-

5 Y^5 is selected from the group consisting of:

an aryl, substituted aryl group, $-COR^4$, $-COOR^4$, $-CONR^6R^7$, $-CSR^4$, and $-C(S)NR^6R^7$;

Y^6 is selected from the group consisting of:

an aliphatic group, a substituted aliphatic group, an aryl, substituted aryl group,
 $-COR^4$, $-COOR^4$, $-CONR^6R^7$, $-CSR^4$, and $-C(S)NR^6R^7$;

10 R^4 , R^6 and R^7 are each independently selected from the group consisting of:

H, an aliphatic group, a substituted aliphatic group, an aryl group and a substituted aryl group;

Y^3 is selected from the group consisting of:

H, aliphatic, substituted aliphatic, aryl, substituted aryl, heteroaryl, substituted heteroaryl,
 15 and $(C_1-C_{10} \text{ alkyl})-R^8$; R^8 is selected from the group consisting of aryl, substituted aryl, heteroaryl, and substituted heteroaryl; and

R^5 is selected from the group consisting of:

H, aliphatic group, a substituted aliphatic group, heteroaryl, substituted heteroaryl, an aryl, a substituted aryl, and $(C_1-C_{10} \text{ alkyl})-R^9$;

20 R^9 is selected from the group consisting of

aryl, substituted aryl, heteroaryl, and substituted heteroaryl, aminoalkyl, and cycloalkyl;

V is a bond or O;

X is CH_2 or O;

Y^4 is selected from the group consisting of:

25 $-(C_1-C_3)\text{alkyl}-O-W-Y^7$, $-C(O)NY^8Y^9$, $-(C_1-C_3)\text{alkyl}-NY^{10}Y^{11}$, and
 $-(C_1-C_3)\text{alkyl}N(Y^{13})W-(C_0-C_5)\text{alkyl}-Y^{14}$;

W is selected from the group consisting of:

a bond, $-CONY^{12}$, $-C(O)-$, $-OCH_2-$, C_1-C_6 alkyl, $-CO_2-$, $-CHOY^{15}$, $-CSNY^{16}$ and
 $-SO_2-$;

30 Y^7 is selected from the group consisting of:

aryl, substituted aryl, heteroaryl, substituted heteroaryl, aliphatic, branched aliphatic and substituted (C_1-C_{10}) alkyl;


Y^8 , Y^9 , Y^{10} , Y^{11} , Y^{12} , Y^{13} , Y^{14} , Y^{15} , and Y^{16} are each independently selected from the group consisting of:

35 aryl, substituted aryl, heteroaryl, substituted heteroaryl, aliphatic, branched aliphatic and substituted (C_1-C_{10}) alkyl; and


n^1 is 2, 3, 4 or 5.

-413-

- 5 3. The compound of Claim 1, wherein Y^{1a} is selected from the group consisting of: aryl, heteroaryl, cycloalkyl, heterocycloalkyl, aryloxy,




4. The compound of Claim 1 or 2, wherein the compound is represented by the following structural formula,




10 wherein E is O or S.

- 5 5. The compound of Claim 4, wherein the compound is represented by the following structural formula,



wherein q is 1 or 2; and each R⁵ is independently hydrogen or methyl.


6. The compound of Claim 4, wherein the compound is represented by the following structural formula,



wherein q is 1 or 2; and each R⁵ is independently hydrogen or methyl.


-415-

- 5 7. The compound of Claim 4, wherein the compound is represented by the following structural formula,




wherein q is 1 or 2; and each R⁵ is independently hydrogen or methyl.

8. The compound of Claim 4, wherein the compound is represented by the following structural formula,



wherein q is 0 or 1; and each R⁵ is independently hydrogen or methyl.


9. The compound of Claim 4, wherein the compound is represented by the following structural formula,



wherein q is 1 or 2; and each R⁵ is independently hydrogen or methyl.


-416-

- 5 10. The compound of Claim 4, wherein the compound is represented by the following structural formula,




wherein q is 1 or 2; and each R⁵ is independently hydrogen or methyl.

- 10 11. The compound of Claim 4, wherein the compound is represented by the following structural formula,



wherein q is 1 or 2; and each R⁵ is independently hydrogen or methyl.

12. The compound of Claim 4, wherein the compound is represented by the following structural Formula,




15


wherein q is 1 or 2; and each R⁵, Y² and Y³ are independently hydrogen or methyl.

-417-

- 5 13. The compound of Claim 4 represented by the following structural formula,




wherein Y^{1a} is optionally substituted phenyl, naphthyl,




- 10 and Z is a bond, oxygen, -NH-, -N(CH₃)-, -NHC(O)- or -C(O)NH-.

14. The compound of Claim 4 represented by the following structural formula,



wherein Y^{1a} is optionally substituted phenyl, naphthyl or


- 15 Z is a bond, oxygen, -NH-, -N(CH₃)-, -NHC(O)- or -C(O)NH-.



and


-418-

- 5 15. The compound of Claim 4 represented by the following structural formula,



wherein Y^{1a} is optionally substituted aryl, heteroaryl, heterocycloalkyl, heteroaryl-Z-heterocycloalkyl or heteroaryl-Z-aryl.


- 10 16. The compound of Claim 1 represented by the following structural formula,



wherein q is 1 or 2; and each R^5 is independently hydrogen or methyl.


-419-

- 5 17. The compound of Claim 1 represented by the following structural formula,



wherein q is 1 or 2; and each R^5 is independently hydrogen or methyl.


- 10 18. The compound of Claim 1 represented by the following structural formula,



wherein,

Y^{1a} is hydrogen, aryl, heteroaryl, or aryloxy; q is 1 or 2; and n^1 is 2, 3, or 4.


19. The compound of Claim 1 represented by the following structural formula,




wherein,

Y^{1a} is hydrogen, aryl, heteroaryl or aryloxy; q is 1 or 2; and n^1 is 2, 3, or 4.

20. The compound of Claim 1 or 2 represented by a following structural formula,









21. The compound of Claim 1 or 2 represented by a following structural formula,














5

22. A compound selected from the group consisting of:







No.	Compound	Name
1		3-{2-(Diphenylacetylaminomethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
2		3-{2-[(2-Cyclopropylacetylaminomethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
3		3-{2-[(3-Methoxybenzoylamino)methyl]-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
4		3-{2-[(Biphenyl-2-carbonyl)amino]methyl}-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
5		3-(4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-[(2,5-dichlorothiophene-3-carbonyl)amino]methyl}phenyl) propionic acid
6		3-{2-(Isopropoxycarbonylaminomethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid







No.	Compound	Name
7		3-{2-(1,3-Dioxo-1,3-dihydroisoindol-2-ylmethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
8		3-{4-[2-(5-Methyl-2-phenyloxazol-4-yl)ethoxy]-2-[(3-phenylureido)methyl]phenyl} propionic acid
9		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid
10		3-{2-(Isopropoxycarbonylaminomethyl)-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid
11		3-{2-(Benzoylaminomethyl)-4-[3-(biphenyl-4-yloxy)propoxy]phenyl} propionic acid
12		3-(4-[3-(Biphenyl-4-yloxy)propoxy]-2-[[pyridine-2-carbonyl]amino]methyl} phenyl} propionic acid

No.	Compound	Name
13		3-{2-Benzylcarbamoyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
14		3-{2-Benzylcarbamoyl-4-[2-(2-biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]phenyl} propionic acid
15		3-{4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-cyanophenyl} propionic acid
16		3-[4-[2-(2-Biphenyl-3-yl-5-methyloxazol-4-yl)ethoxy]-2-(1,3-dioxo-1,3-dihydroisoindol-2-ylmethyl)phenyl] propionic acid
17		3-{2-(2-Isopropoxycarbonylaminoethyl)-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
18		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid







No.	Compound	Name
19		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(3-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
20		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(6-phenylpyridin-3-yl)thiazol-4-yl]ethoxy}phenyl) propionic acid
21		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(2-isopropoxycarbonylaminoethyl)phenyl] propionic acid
22		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(isobutoxycarbonylaminomethyl)phenyl] propionic acid
23		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(cyclopentyloxycarbonylaminomethyl)phenyl] propionic acid







-425-

No.	Compound	Name
24		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[2-(4-isopropoxyphenyl)-5-methyloxazol-4-yl]ethoxy}phenyl) propionic acid
25		3-(2-Benzylcarbamoyl-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
26		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-morpholin-4-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
27		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-piperidin-1-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
28		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-pyrimidin-2-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
29		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-pyrazin-2-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid








No.	Compound	Name
30		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(6-phenoxy pyridin-3-yl)thiazol-4-yl]ethoxy}phenyl) propionic acid
31		3-{2-Cyclohexylcarbamoxyloxymethyl-4-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid
32		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(4-phenylaminophenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
33	 <p>ClH</p>	3-(4-{2-[5-Methyl-2-(6-phenylpyridin-3-yl)thiazol-4-yl]ethoxy}-2-[[pyridine-2-carbonyl)amino]methyl}phenyl) propionic acid HCl salt
34		3-{4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-[(3-methylbutyrylamino)methyl]phenyl} propionic acid
35		3-{2-(Isopropoxycarbonylaminomethyl)-4-[2-(5-methoxy-2-phenyloxazol-4-yl)ethoxy]phenyl} propionic acid

No.	Compound	Name
36		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(6-phenoxy pyridin-3-yl)oxazol-4-yl]ethoxy}phenyl) propionic acid
37		3-{4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-cyclohexylcarbamoyloxymethyl phenyl} propionic acid
38		3-{2-Cyclohexylcarbamoyloxymethyl-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid
39		3-(2-Cyclohexylcarbamoyloxymethyl-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
40		3-(2-Cyclohexylcarbamoyloxymethyl-4-{2-[5-methyl-2-(4-morpholin-4-ylphenyl)oxazol-4-yl]ethoxy}phenyl) propionic acid
41		3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[3-(tetrahydropyran-4-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid

No.	Compound	Name
42		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(cyclopropylmethoxycarbonylamino-methyl)phenyl] propionic acid
43		3-{2-(Cyclopropylmethoxycarbonylamino-methyl)-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid HCl salt
44		3-[4-[2-(2-Biphenyl-4-yl-5-methyloxazol-4-yl)ethoxy]-2-(cyclobutoxycarbonylamino-methyl)phenyl]propionic acid HCl salt
45		3-{2-(Cyclobutoxycarbonylamino-methyl)-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid HCl salt
46		3-[4-{2-[2-(4-Butyrylamino-phenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylamino-methyl)phenyl] propionic acid
47		3-{2-(Isopropoxycarbonylamino-methyl)-4-[2-(5-methyl-2-{4-[(pyridine-2-carbonyl)-amino]-phenyl}-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid

No.	Compound	Name
48		3-(4-[2-(2-Biphenyl-4-yl)-5-methyloxazol-4-yl]ethoxy)-2-[(pyrazine-2-carbonyl)amino]methyl]phenyl] propionic acid
49		3-[4-{2-[2-(3-Cyclohexylcarbamoylphenyl)-5-methyloxazol-4-yl]ethoxy}-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid
50		3-(2-(Isopropoxycarbonylaminomethyl)-4-{2-[5-methyl-2-(2-phenoxyphenyl)oxazol-4-yl]ethoxy}phenyl] propionic acid
51		3-(2-Cyano-4-{2-[5-methyl-2-(4-phenoxyphenyl)oxazol-4-yl]-ethoxy}-phenyl] propionic acid
52		3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(pyridin-2-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid
53		3-[4-[2-(2-Biphenyl-4-yl)-5-methyloxazol-4-yl]ethoxy]-2-(4-trifluoromethylphenoxy)methyl]phenyl] propionic acid

No.	Compound	Name
54		3-{2-(Isobutoxycarbonylaminomethyl)-4-[2-(5-methyl-2-morpholin-4-ylthiazol-4-yl)ethoxy]phenyl} propionic acid
55		3-[2-(Isopropoxycarbonylaminomethyl)-4-(2-{5-methyl-2-[4-(pyrimidin-2-yloxy)phenyl]oxazol-4-yl}ethoxy)phenyl] propionic acid
56		3-[4-[2-(2-Biphenyl-4-yl-5-methoxyoxazol-4-yl)ethoxy]-2-(isopropoxycarbonylaminomethyl)phenyl] propionic acid
57		3-(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[[pyridine-2-carbonyl]-amino]-methyl}-phenyl)-propionic acid
58		3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(2,4,5-trifluoro-benzoylamino)-methyl]-phenyl}-propionic acid
59		3-{2-[(2,4-Difluoro-benzoylamino)-methyl]-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
60		3-(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[[thiophene-2-carbonyl]-amino]-methyl}-phenyl)-propionic acid

No.	Compound	Name
61		3-(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[[thiophene-2-carbonyl]-amino]-methyl}-phenyl)-propionic acid
62		3-{2-(Butyrylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
63		3-{2-[(Cyclobutanecarbonyl)-amino]-methyl}-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
64		3-{2-(Benzyloxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
65		3-{2-(tert-Butoxycarbonylamino-methyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid
66		3-{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-2-[(2-phenoxy-acetyl)-amino]-methyl}-phenyl}-propionic acid
67		3-{2-[(Cyclopentanecarbonyl)-amino]-methyl}-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid

5 23. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and at least one compound of Claims 1-22 or a pharmaceutically acceptable salt, solvate or hydrate thereof.

 24. A pharmaceutical composition comprising (1) a compound of Claim 1 or 2, or a pharmaceutically acceptable salt, solvate, hydrate or stereoisomer thereof; (2) a second therapeutic agent selected from the group consisting of insulin sensitizers, sulfonylureas, biguanides, thiazolidinediones, α -glucosidase inhibitors, insulin secretagogues, insulin, antihyperlipidemic agents, plasma HDL-raising agents, HMG-CoA reductase inhibitors, statins, acyl CoA:cholesterol acyltransferase inhibitors, antiobesity compounds, antihypercholesterolemic agents, fibrates, vitamins and aspirin; 10 and (3) a pharmaceutically acceptable carrier.

 25. A method of modulating a peroxisome proliferator activated receptor (PPAR), comprising the step of contacting the receptor with at least one compound of Claims 1-22, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

20 26. The method of Claim 25, wherein the peroxisome proliferator activated receptor is an alpha-receptor.

 27. The method of Claim 25, wherein the peroxisome proliferator activated receptor is a gamma-receptor.

25 28. A method for treating or preventing a peroxisome proliferator activated receptor-gamma mediated disease or condition comprising the step of administering an effective amount of at least one compound of Claim 1 or 2.

 29. A method for lowering blood-glucose comprising the step of administering an effective amount of at least one compound of Claim 1 or 2.

30 30. A method of treating or preventing disease or condition selected from the group consisting of hyperglycemia, dyslipidemia, Type II diabetes, Type I diabetes, hypertriglyceridemia, syndrome X, insulin resistance, heart failure, diabetic dyslipidemia, hyperlipidemia, hypercholesteremia, hypertension, obesity, anorexia bulimia, anorexia nervosa, cardiovascular disease and other diseases where insulin resistance is a component, comprising the step of administering an effective amount of at 35 least one compound of Claim 1 or 2.

5 31. A method of treating or preventing diabetes mellitus in a mammal comprising the step of administering to a mammal a therapeutically effective amount of at least one compound of Claim 1 or 2.

 32. A method of treating or preventing cardiovascular disease in a mammal comprising the step of administering to a mammal a therapeutically effective
10 amount of at least one compound of Claim 1 or 2, or a pharmaceutically acceptable salt, solvate, hydrate or stereoisomer thereof.

 33. A method of treating or preventing syndrome X in a mammal, comprising the step of administering to the mammal a therapeutically effective amount of at least one compound of Claim 1 or 2, or a pharmaceutically acceptable salt, solvate,
15 hydrate or stereoisomer thereof.

 34. A method of treating or preventing disease or condition selected from the group consisting of hyperglycemia, dyslipidemia, Type II diabetes, Type I diabetes, hypertriglyceridemia, syndrome X, insulin resistance, heart failure, diabetic dyslipidemia, hyperlipidemia, hypercholesteremia, hypertension, obesity, anorexia
20 bulimia, anorexia nervosa, cardiovascular disease and other diseases where insulin resistance is a component, comprising the step of administering an effective amount of at least one compound of Claim 1 or 2 and an effective amount of second therapeutic agent selected from the group consisting of: insulin sensitizers, sulfonylureas, biguanides, thiazolidinediones, α -glucosidase inhibitors, insulin secretagogues, insulin,
25 antihyperlipidemic agents, plasma HDL-raising agents, HMG-CoA reductase inhibitors, statins, acyl CoA:cholesterol acyltransferase inhibitors, antiobesity compounds, antihypercholesterolemic agents, fibrates, vitamins and aspirin.

 35. Use of a compound of Claim 1 or 2 and pharmaceutically acceptable salt, solvate, hydrate or stereoisomer thereof, for the manufacture of a
30 medicament for the treatment of a condition modulated by a PPAR.

INTERNATIONAL SEARCH REPORT

 Inte 1st Application No
 PCT/US 02/15143

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 7 A61K31/42 C07D263/32 A61P3/10 C07D231/12 A61K31/422
 A61K31/426 A61K31/427 C07D233/58 C07D277/24 C07D277/38
 C07C271/24 C07C271/22 C07C233/87 C07C233/63 C07D413/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 02 18355 A (BROOKS DAWN ALISA ;CONNOR SCOTT EUGENE (US); DOMINIANNI SAMUEL JAM) 7 March 2002 (2002-03-07) page 213; example 50	1-35
P, Y	WO 02 16332 A (BROOKS DAWN ALISA ;GODFREY ALEXANDER GLENN (US); MANTLO NATHAN BRY) 28 February 2002 (2002-02-28) claims definition of R2/R3	1-35
Y	WO 01 16120 A (DOMINIANNI SAMUEL J ;MATTHEWS DONALD P (US); MICHELLYS PIERRE YVES) 8 March 2001 (2001-03-08) claims definition of R2/R3	1-35

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

3 September 2002

Date of mailing of the international search report

12/09/2002

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Kollmannsberger, M

INTERNATIONAL SEARCH REPORT

 Inter Application No
 PCT/US 02/15143

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 7 C07D417/12 C07D413/14 C07D413/04 C07D413/10 C07D417/14
 C07D417/04 C07D213/81 C07D307/83

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01 00566 A (ZANDT MICHAEL C VAN ; INST FOR PHARMACEUTICAL DISCOV (US)) 4 January 2001 (2001-01-04) claim 1 see definition of R1-R4 and R7; page 77 -page 78; examples 43,44 page 1, line 16 - line 29 ----	1-35
E	EP 1 216 980 A (EISAI CO LTD) 26 June 2002 (2002-06-26) page 135 -page 137; examples 771-785 page 1, paragraph 1 -page 2, paragraph 18 page 48 -page 50	1-35
Y	& WO 01 25181 A (EISAI CO LTD) 12 April 2001 (2001-04-12) -----	1-35



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

3 September 2002

Date of mailing of the international search report

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Kollmannsberger, M

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 02/15143

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 25-34 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. Appl. Application No

PCT/US 02/15143

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0218355	A	07-03-2002	AU	8465801 A	13-03-2002
			WO	0218355 A1	07-03-2002
WO 0216332	A	28-02-2002	AU	8466001 A	04-03-2002
			WO	0216332 A1	28-02-2002
WO 0116120	A	08-03-2001	AU	7073400 A	26-03-2001
			EP	1206457 A1	22-05-2002
			WO	0116120 A1	08-03-2001
			US	6417212 B1	09-07-2002
WO 0100566	A	04-01-2001	AU	5886900 A	31-01-2001
			BR	0011928 A	09-04-2002
			CN	1364153 T	14-08-2002
			CZ	20014637 A3	17-04-2002
			EP	1198451 A2	24-04-2002
			NO	20016272 A	17-01-2002
			WO	0100566 A2	04-01-2001
			US	6420426 B1	16-07-2002
EP 1216980	A	26-06-2002	AU	7449900 A	10-05-2001
			EP	1216980 A1	26-06-2002
			WO	0125181 A1	12-04-2001